

# Elsberry Plant Materials Center 2002 Annual Technical Report



Plant Solutions for  
Conservation Needs



Plant  
Materials  
Program

Visit us: <http://Plant-Materials.nrcs.usda.gov> or <http://www.mo.nrcs.usda.gov>

**ELSBERRY PLANT MATERIALS CENTER  
2002**

***Advisory Committee***

**Roger A. Hansen, State Conservationist, Missouri, Chairman  
Leroy Brown, State Conservationist, Iowa  
William J. Gradle, State Conservationist, Illinois**

***Resource Personnel***

**Charles R. Freeland, State Resource Conservationist, Missouri  
(January 1, 2002 – June 29, 2002)  
R. Dwaine Gelnar, State Resource Conservationist, Missouri  
(October 6, 2002-December 31, 2002)  
James E. Ayen, State Resource Conservationist, Iowa  
Richard G. Hungerford, Jr., State Resource Conservationist, Illinois**

***Plant Materials Specialist***

**Jerry U. Kaiser**

***Plant Materials Personnel***

**Jimmy Henry, Plant Materials Center Manager  
Steven B. Bruckerhoff, Conservation Agronomist  
Pamela K. Stewart, Secretary  
Ronald L. Cordsiemon II, Biological Science Technician  
Donald D. Tapley, Agricultural Science Research Technician**



**2002  
Annual Technical Report  
Elsberry Plant Materials Center  
Elsberry, Missouri**

<i>Introduction/History</i> .....	5
<i>Plant Materials Center Operations</i> .....	6
<i>Climatic Data</i> .....	7
<i>Tours, Visitors, and Meetings</i> .....	9
 <i>Study Activities:</i>	
<b><u>29I093R</u></b> – Miscellaneous Herbaceous Plant Evaluation .....	10
<b><u>29I097G</u></b> – Assembly and Evaluation of Big Bluestem, <i>Andropogon gerardii</i> .....	13
<b><u>29I101J</u></b> – Assembly and Evaluation of Arrowwood, <i>Viburnum dentatum</i> , L. ....	19
<b><u>29I107G</u></b> – Assembly and Evaluation of Eastern Gamagrass, <i>Tripsacum dactyloides</i> , L. ....	21
<b><u>29I108G</u></b> – Assembly and Evaluation of Low Growing Rhizomatous Switchgrass for Use in Waterways, Filter Strips and Other Conservation Uses .....	27
<b><u>29I110J</u></b> – Assembly and Evaluation of Chokecherry, <i>Prunus virginiana</i> .....	33
<b><u>29A116W</u></b> – Evaluation of Miscellaneous Trees and Shrubs .....	41
<b><u>29I124G</u></b> – Production of Native Iowa Ecotypes of Grasses and Forbs for Roadside, Critical Areas, and All Other Vegetative Plantings Where Native Grasses and Forbs are Now Being Planted .....	45
<b><u>29A128J</u></b> – Flowering Dogwood, Interagency Study Between Department of Interior, National Parks, Service, National Capital Region and Department of Agriculture .....	50
<b><u>29I132O</u></b> – Miscellaneous Wetland Plant Evaluation .....	51
<b><u>29I134J</u></b> – Assembly and Evaluation of Eastern Redcedar, <i>Juniper virginiana</i> L. ....	55
<b><u>29I135J</u></b> – Assembly and Evaluation of Hazelnut, <i>Corylus americana</i> , Walt.. ....	69
<b><u>29I136J</u></b> – Assembly and Evaluation of Wild Plum, <i>Prunus americana</i> , Marsh. ....	77
<b><u>29A137O</u></b> – Wetland/Riparian Propagation, Establishment, and Demonstration ...	90

Continued: ...

<b><u>29I141G</u></b> – Assembly and Evaluation of Little Bluestem, <i>Schizachyrium scoparium</i> , Michx. ....	100
<b><u>29I142G</u></b> – Production of Native Missouri Ecotypes of Grasses, Legumes, and Forbs for Roadsides, Critical Areas, and All Other Vegetative Plantings Where Native Plants are Now Being Planted .....	120
<b><u>29I143G</u></b> – Seed Coating/Seeding Rates Study .....	129
<b><u>MOPMC-P-001, WO, WL, WE</u></b> – Assembly, Evaluation and Selection of Bur Oak, <i>Quercus macrocarpa</i> , Michx. ....	159
<b><u>MOPMC-P-002, WE, WL</u></b> – Assembly, Evaluation and Selection of False Indigo Bush, <i>Amorpha fruticosa</i> , L. ....	161
<b><u>MOPMC-P-003, PA, WL</u></b> – Evaluation and Release of Eastern Gamagrass, <i>Tripsacum dactyloides</i> , L. ....	164
<b><u>MOPMC-T-0104</u></b> – Native Plant Identification .....	168
<b><u>MOPMC-T-0105, PA</u></b> – Compatibility Study Using Native Warm Season and Cool Season Grasses with Native Legumes and Forbs .....	169
<b><u>MOPMC-T-0106, BU</u></b> – Collection and Evaluation of Native Cool Season Grasses and Sedges for Filter Strips .....	174
<b><u>MOPMC-P-0107, PA, WL</u></b> – Evaluation and Release of Big Bluestem, <i>Andropogon gerardii</i> , L. ....	176
<b><u>MOPMC-T-0208-PA</u></b> – Testing Warm Season Grasses for Forage Quality .....	177
<b><u>MOPMC-P-0209, PA, WL</u></b> – Evaluation and Release of <i>Paspalum</i> Species .....	179
Released From the Elsberry PMC .....	181
Studies/Projects at the Elsberry PMC 1958 through the Present .....	183
Herbaceous and Woody Seed and Plant Production 2002 .....	190

## **Introduction**

The Elsberry Plant Materials Center (PMC) was established in June 1934. The Center is located approximately 60 miles northwest of St. Louis, Missouri, on Highway 79. It includes 243 acres of various soil types.

The Elsberry PMC primarily serves Illinois, Iowa and Missouri, however, it makes significant contributions to other states in the Midwest region.

The mission of the NRCS Plant Materials Program is to develop and transfer plant materials and plant technology for the conservation of natural resources. In working with a broad range of plant species, including grasses, forbs, trees, and shrubs, the program seeks to address priority needs of field offices and land managers in both public and private sectors. Emphasis is focused on using native plants as a healthy way to solve conservation problems and protect ecosystems.

The objectives of the Elsberry PMC and of the plant materials program is to assemble, test, select and develop improved plants; and to develop reliable techniques for successfully establishing and maintaining plants for conservation uses.

Of particular importance is finding suitable plants for wetland situations, high traffic areas, wildlife food and habitat, farmstead and field windbreaks, windbarriers, pastures, landscape and beautification, roadside restoration, biofuel concerns, riparian plantings, woodland, erosion control on cropland and etc.

Each of the three states served by the Center has identified their plant materials problems, needs and priorities. PMC activities are directed toward meeting the needs and priorities set forth in the states' long-range plans.

## **History**

The Elsberry Plant Materials Center was established in June 1934, which makes it the oldest Center in the nation. During the Center's earlier existence it produced 10,000,000 seedlings for use in windbreaks during the dust bowl era. As early as 1939 the Center began searching for plants to respond to specific conservation problems. The Center is located approximately 60 miles northwest of St. Louis, Missouri, on Highway 79. It includes 243 acres of land of which 60 percent is bottomlands and 40 percent is uplands.

## **Plant Materials Center Operations**

The Center's operations are carried out in accordance with policies set forth in the National Plant Materials Handbook.

Guided by the Center's Multi-Year Business Plan, plant species are collected (mainly local field collections [95%]). Other collections come from locations within the species range in the United States. Center personnel then prepare the seed/plant for planting. Each collection is given an identification number (accession) and planted in a uniform nursery. Initial evaluation data is recorded on such factors as seedling emergence and vigor, rate of growth, disease and insect resistance, and ability to spread. Also recorded are date and amount of bloom, seed production, winter hardiness, and foliage characteristics. Selections are made and seed increased for advanced evaluation plantings. Field plantings are then conducted to determine plant performance and soil and climatic adaptation throughout its intended area of use. Evaluations are made comparing selected candidate accessions with "standards of comparison" such as cultivars or varieties that are already in the commercial market, or other species used for the same purpose.

After several years (10-15) of evaluation, selected accessions are cooperatively released with the USDA-Agricultural Research Service (ARS), State Agricultural Experiment Stations, Conservation Commissions, Universities, Departments of Transportation, and/or other interested agencies. The Center releasing a named variety is responsible for maintaining the breeder and foundation seed. These fields undergo annual inspections by the Missouri Seed Improvement Association to insure that seed is available to commercial producers and ultimately to the public for solving conservation problems.

Additional avenues have been established and used by the Plant Materials discipline to release plants to the commercial market: Source Identified, Selected and Tested. These three new avenues provide a quicker release of plants as compared to cultivar release (10-15 years).

The Elsberry Plant Materials Center has released 68 plants during its 68-year history. Sixty-two of the total numbers of plants released are natives.

**CLIMATIC DATA – CALENDAR YEAR 2002**

**TEMPERATURE (Fahrenheit)**

<b>Month</b>	<b>70 Year Monthly High Average</b>	<b>Year 2002 Monthly High Average</b>	<b>Year 2002 Monthly High Departure</b>	<b>70 Year Monthly Low Average</b>	<b>Year 2002 Monthly Low Average</b>	<b>Year 2002 Monthly Low Departure</b>
<b>January</b>	38.1	45.23	+7.13	18.11	25.10	+6.99
<b>February</b>	43.22	47.21	+3.99	22.37	25.93	+3.56
<b>March</b>	53.79	50.68	-3.11	36.65	29.45	-7.20
<b>April</b>	66.66	68.10	+1.44	41.91	45.00	+3.09
<b>May</b>	76.56	73.00	-3.56	56.92	51.26	-5.66
<b>June</b>	85.44	86.27	-.83	71.62	64.30	-7.32
<b>July</b>	89.69	91.26	-1.57	64.56	68.65	+4.09
<b>August</b>	87.65	87.94	+.29	62.40	66.55	+4.15
<b>September</b>	80.43	81.87	+1.44	54.14	57.67	+3.53
<b>October</b>	69.48	62.35	-7.13	43.05	44.06	+1.01
<b>November</b>	50.31	51.22	+.91	32.05	30.73	-1.32
<b>December</b>	42.03	43.87	+1.84	22.68	27.19	+4.51

<b>2002</b>	
<b>Last Killing Frost</b>	April 23
<b>First Killing Frost</b>	Nov 2
<b>Number of Frost-Free Days</b>	207

## CLIMATIC DATA – CALENDAR YEAR 2001

### Precipitation (Inches)

<b><u>Month</u></b>	<b><u>72 Year Average</u></b>	<b><u>2002 Total</u></b>	<b><u>Departure</u></b>
<b>January</b>	1.87	2.53	+.66
<b>February</b>	1.99	1.24	-.75
<b>March</b>	3.16	3.21	+.05
<b>April</b>	3.70	4.46	+.76
<b>May</b>	4.10	13.97	+9.87
<b>June</b>	3.78	3.15	-.63
<b>July</b>	3.41	.79	-2.62
<b>August</b>	3.35	4.29	+.94
<b>September</b>	3.30	1.29	-2.01
<b>October</b>	3.00	3.89	+.89
<b>November</b>	2.87	.61	-2.26
<b>December</b>	2.46	1.83	-.63
<b>Year Total</b>	37.04	41.26	+4.22

## **Tours, Visitors, and Meetings**

The Elsberry Plant Materials Center was visited by 309 registering guests. These individuals represented many walks of life, foreign and domestic, students, farmers, ranchers, researchers and other professionals.

They came individually and in formal groups. All were interested in one or more aspects of our dynamic soil and water conservation program.

The following groups are representative of the interest in the Elsberry Plant Materials Program. Not all individuals are included in this listing.

<u>Groups</u>	<u>Date 2001</u>	<u>No. of Participants</u>
Grass-Based Dairy Group Meeting	January 15-17	5
Lost Creek Watershed Meeting	February 19	12
Bowling Green Group of Missouri Department of Conservation	February 21	7
Clarence Cannon Memorial Watershed Meeting	February 25	15
ECAP Ethanol Group Meeting	March 21	14
Grow Native Group Meeting	April 9	9
PMC Annual Tour	June 12	45
Grace Hill Americorps St. Louis	June 13	7
Prairie Conference Meeting	June 25	30
Missouri Ecotype Meeting	July 16	10
Preconstruction Group Meeting	September 3	8
NEMO Scottish Rite Group	October 1	44
Missouri PM Committee Meeting	October 1-3	22
State Conservationists' Advisory Committee Meeting	October 22	6
North Technical High School	October 30	10
EMP Planning Meeting (MDC)	December 16-17	19
MDC St. Louis Wildlife Meeting	December 18	10

**Study: 29I093R**

**Study Title:** Miscellaneous Herbaceous Plant Evaluation.

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

Plants arrive at the Plant Materials Center (PMC) from many sources and for many different purposes. Most of the plants are assigned to a specific study. Plants are also received that are not tied to a specific study. These can be from other PMC's for area of adaptation or plants in advanced stages of evaluation. Plants are received from individuals who are interested in an unfamiliar species or a plant with unusual characteristics. Many species exist on the center that are not involved with an active study addressing a specific problem.

**Problem:**

Keeping track of numerous miscellaneous plants around the PMC without an organized evaluation system became inefficient. This study organizes miscellaneous plant material coming into the center for evaluation.

**Objective:**

To evaluate winter hardiness, insect and disease resistance, and vigor of plants for climatic adaptation. Plants brought in for other specific reasons like forage production, landscape beautification, shoreline stabilization, etc., will be evaluated accordingly.

**Procedure:**

As miscellaneous plants are received at the center, they are assigned an accession number and as much background information as available or necessary are documented. The accession is then assigned a location for planting that best suits its needs for evaluation. Plants are evaluated as necessary. Many plants are left for plant identification sessions or demonstrations for several years.

**Discussion:****1984-1990**

This study was initiated in April 1984 in the PMC pipeline area. There are approximately 150 different accessions of the following species of plants: indiagrass, switchgrass, big bluestem, purpletop, little bluestem, buffalograss, wheatgrass, fescue, timothy, ryegrass, redtop, orchardgrass, kura clover, blackeyed susan, and lespedeza. Factors involved in evaluations dealt with area of adaptation.

## 1991-1994

Approximately 75 accessions were added during 1991. Forty of them were warm season grasses used in three FEP (Field Evaluation Planting) variety studies: 29A111G, 29A118G, and 29A127G. Twenty-six were accessions of common cool season grasses and legumes used for pasture and hay in the three state area. These were commonly used for plant identification sessions.

## 1995-1998

The accessions added in 1997 are being looked at for forage. They include 'Steadfast' birdsfoot trefoil, 'Mandan' Canada wildrye, and several bermudagrasses including 'Hardy' and 'OK-74-12-6'. Also zoysia grass, centipedegrass, and buffalograss from the Fort Leonard Wood Wear Tolerance Study are being looked at for adaptation. Several big bluestem accessions from Study 29I097G are being evaluated as landscape plants.

## 1999

The accessions added in 1999 are a Lincoln County Missouri collection of Virginia wildrye and a Crawford County Missouri collection of Virginia wildrye variation geneses. These species are being looked at for shade tolerance for riparian areas and cover crop for tree plantings.

## 2000

No new accessions were added in 2000. Two species that are getting the most interest are the Lincoln County accessions of Virginia wildrye and 'Tufcote' bermudagrass.

The Lincoln County accession of Virginia wildrye is a shade tolerant cool season grass that has potential for a covercrop for woody plantings as well as a possible buffer species along riparian areas. This accession should be in commercial production and available soon.

The 'Tufcote' bermudagrass accession was tested at Fort Leonard Wood for wear tolerance and showed very good potential. It could be used on playgrounds, sports fields, lawns, as well as having potential for high livestock use areas. This species is not native and does show potential for spreading so it should not be planted in areas where it could escape and cause problems.

## 2001

Three new species of native legumes were added in 2001. Native legumes are seldom used in mixtures with warm season grasses planted for pastures primarily because of their cost, lack of availability, and lack of knowledge on which ones will perform best in a mixture. The following species were planted for observational evaluation: goats rue, *Tephrosia virginiana*; sensitive brier, *Schrankia uncinata*; and Sampson's snakeroot, *Orbexilium pedunculatum*.

The Lincoln County Missouri collection of Virginia wildrye, accession 9083169, has shown excellent vigor and seed production. Forage quality is comparable to tall fescue, spring green-up earlier than tall fescue and seedhead emergence is approximately two weeks later than tall fescue. This accession is scheduled for release in 2002.

## 2002

One new collection was planted in the miscellaneous block. Accession 9083240, western wheatgrass, *Pascopyrum smithii*, was planted as greenhouse plugs May 10, 2002. This material was collected in Audrain County, Missouri.

The Lincoln County Missouri collection of Virginia wildrye, accession 9083169, was released as a selected class and given the name Cuivre River. The Cuivre River selection has early vigorous growth that is earlier than tall fescue. Booting occurred at the end of May to the first week of June at Elsberry. This is approximately two weeks later than tall fescue.

Although Cuivre River was released as a selection and only limited testing has been done, its anticipated uses are wildlife food/cover, plant diversity in wetland and riparian plantings, covercrop for woody plantings, erosion control, and forage.

Cuivre River has not been tested for grazing but forage clippings were taken at different stages of growth and compared to tall fescue clippings from adjacent plots. Forage quality of the Cuivre River selection compared favorably to tall fescue as indicated by data below.

Clipping Date	Percent Protein		Percent ADF		Percent NDF	
	<u>TF</u>	<u>VWR</u>	<u>TF</u>	<u>VWR</u>	<u>TF</u>	<u>VWR</u>
4/24/02		27		26		47
5/30/01	9	12	40	34	61	60
10/11/01	15	15	31	34	52	55
11/15/01	20	17	22	24	37	44

TF = tall fescue, VWR = Cuivre River Virginia wildrye, ADF = acid detergent fiber, NDF = neutral detergent fiber

**Study: 29I097G**

**Study Title:** Assembly and Evaluation of Big Bluestem, *Andropogon gerardii* Vitman.

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

Big bluestem is a tall, warm-season, perennial, native grass with stiff, erect culms; flattened and keeled sheaths; membranous ligules; and flat or folded leaf blades. Big bluestem has developed a very efficient spreading root system that may reach depths of 5-8 feet (150-200 cm). Big bluestem reaches a mature height of 3-4 feet (90-120 cm) in northern latitudes, and 6-8 feet (180-240 cm) or more in the southern part of its natural range. Although short rhizomes may be present, it usually makes a bunch type growth. Big bluestem is composed of many ecotypes with a wide range of adaptation to soil and climate. Big bluestem is one of the most widespread and important forage grasses of the North American tallgrass prairie region. It is usually associated with one or more of the other three dominant species, Indiangrass (*Sorghastrum nutans* (L) Nash.), switchgrass (*Panicum virgatum* L.), and little bluestem (*Schizachyrium scoparium* (Michx.) Nash.) Big bluestem occurs on subirrigated lowlands, nearly level to gently undulating glacial till plains, overflow sites, level swales and depressions, residual and glacial uplands, and stream terraces and bottomlands along rivers and tributaries. The abundant, leafy forage is palatable to all classes of livestock.

**Problem:**

There is a need for an adapted variety of big bluestem for pasture and range seedings, surface mine reclamation, critical area planting, recreational area development and other conservation uses in Arkansas and Southern Missouri.

**Objective:**

The objective is to assemble, evaluate, develop and cooperatively release an adapted variety and/or varieties of big bluestem for conservation use in the following Major Land Resource Areas: 116A, 116B, 117, 118, and 119.

**Cooperators:**

USDA-NRCS Plant Materials Center at Elsberry, Missouri and the USDA-NRCS Plant Materials Center at Booneville, Arkansas.

**Assembly:**

The assembly consists of vegetative materials from adapted ecotypes throughout Northwestern Arkansas and Southwestern Missouri Major Land Resource Areas: 116A, 116B, 117, 118, and 119. Collection dates were between November 9 and 13, 1987. Four collection sites per county within the geographic area of collection were made. The number of sites was determined by the size of the county. The study plan supplement lists the states and the number of sites per county.

**Procedure:**

Four collections per county in the targeted Major Land Resource Areas were requested. The intent was to get a broad genetic base of plant material; therefore, the site selection attempt was to get as diverse sampling as practical when selecting superior big bluestem plants in the field. If a county had more than one Major Land Resource Area, collections were made in each area. Collections were from typical locations, which included natural grasslands (range), relic areas, and road right-of-ways. Avoided areas were those that may have been artificially seeded. Where possible, collections came from diverse soil textural types, such as sandy and silty; or range site groupings such as: (1) Run-in sites represented by overflow, or subirrigated; (2) normal upland sites represented by sandy, silty or clayey. Six subsamples (6" x 6" x 8" deep) were collected vegetatively at each site.

The samples were transported in material provided by the Plant Materials Center that included cartons, plastic bags, accession data sheets, and instructions for handling.

Plant Materials Center personnel picked up the cartons containing the samples at designated central locations within each administrative area in November 1987.

Transplanting procedures included temporary storage and handling. The samples were first assigned accession numbers and placed in temporary storage. On February 15, 1988, each subsample was transplanted into separate containers and maintained under controlled greenhouse conditions. The plants were then divided between two locations, Elsberry, Missouri and Booneville, Arkansas Plant Materials Centers, and established in space plant initial evaluation nurseries.

**Discussion:****1987-1989**

A total of 370 accessions (collections) of big bluestem were initially collected during November, 1987, from the targeted areas: 194-Missouri; 85-Arkansas; 82-Oklahoma; and 8-Illinois. Individual plantlets were separated, transplanted into cone-tainers, and grown out in Forrest Keeling Nursery's greenhouse from February until May 1988. More than 4400 individual plantlets were transplanted into a space plant nursery with two replications and six plants per replication. The nursery is located in Field #14 at the PMC and was planted June 1988. The entire nursery was irrigated three times weekly in 1988 to insure good survival. Data collected in 1988 was mostly survival. Data collected in 1989 included survival, vigor, disease resistance, plant size, foliage size, and abundance and visual seed production. Accessions from each state were selected from the above criteria. The numbers selected from each state were as follows: Arkansas-14, Missouri-46, and Oklahoma-13. Table #1 shows the 73 accessions selected from the initial space plant nursery located in Field #14 on the PMC. These plants were vegetatively removed from the initial evaluation nursery in November 1989.

**1990-1991**

The plants selected in 1989 were transplanted into cone-tainers and grown out in the greenhouse that winter. These plants were planted in an isolated crossing block in Field #1 on May 23, 1990. Fifteen bulk pounds of clean seed were harvested in 1991.

### **1992-1993**

The seed harvested in 1991 was sorted by weight and grown in cone-tainers in the greenhouse from January until April. Approximately 500 plants were planted in Field #7 in April and May 1992 for further evaluation.

Beginning in July 1993, the great flood began flooding approximately 86 acres on the PMC. The area where this planting was located was completely inundated with approximately eight feet of water. Just prior to the flooding of this site (July 8, 1993), the PMC staff uprooted 62 selections of big bluestem and re-established them to an upland site on the PMC (Field #8).

### **1994-1996**

The nursery block established in Field #8 in July 1993 was evaluated for forage quality and quantity, seed production, plant maturity differences, and disease and insect resistance. Twenty-eight of the 62 plants were selected and allowed to cross. Seed from this crossing block is a composite of the original 73 accessions collected and is the breeders' block for the new accession 9078831. Seed was harvested in 1995 and 1996 and a seed increase plot will be established in 1997. The Booneville PMC also has made their selection and both will be included in the advanced evaluation.

### **1997-1998**

The diversity in the original nursery block containing all 370 accessions is tremendous. There is a lot of variation within this species. The need for plant diversity for prairie restoration led to the release of the source-identified composite of all 370 accessions. This composite was given the accession number 9062323 and given the name OH-370 which stands for a composite of 370 collections made from the Ozark Highlands of Southern Missouri, Northern Arkansas, Eastern Oklahoma, and Southern Illinois. This plant was released in April 1997.

A 0.4-acre increase planting of 9078832 was planted May 22, 1997, in Field # 6. This planting was established in a conventional seedbed in 36" rows. The first year the planting produced 10 pounds bulk clean seed and in 1998 it produced 27 pounds bulk clean seed. The 1998 seed tested poorly but it is not known why. When seed becomes available from the Arkansas PMC the study will begin an advanced evaluation to compare the new accession, 9078831 with available varieties and also the accession Booneville has selected out of the original assembly of 370 collections.

The original planting was again evaluated the spring of 1997 looking for a tall, stiff stemmed, upright plant to use in wind barriers. Wind erosion is a problem in the flat and sandy crop fields in the bootheel area of Missouri. Switchgrass windbarriers are being tried in areas where field windbreaks using trees are not acceptable. Big bluestem was requested by the Missouri plant materials committee as an additional species to go along with switchgrass since the nursery is still intact. Five accessions (Table #2) were selected and increased vegetatively in the greenhouse and transplanted into an isolation block in Field #4. This block contained 126 plants and of those, 34 plants were selected to represent the crossing block that will serve as the breeders' block for a wind barrier selection. The final accessions represented in this block are 9065960, 9056913, and 9056914.

Selections were also made for landscape and beautification (Table # 3). These selections were transplanted into the rod row initial evaluation area for further evaluation.

## **1999**

The increase plot of 9078831 was expanded in 1999 but did not develop as the 1997 original increase plot did. This accession is scheduled for release as a pre-varietal selection in 2000 if enough seed is available and field plantings are successful.

The wind barrier selection block was again evaluated in 1999 and narrowed down to a single accession, 9066960 (Table #2).

No additional selections were made for landscape plants in 1999 (Table #3).

## **2000**

The increase plot of 9078831 was again expanded in 2000 but again was very slow to germinate. Seed was sent for testing and the sample contained a high percentage of dormant seed. This prevarietal selection was scheduled to be released in 2000 and given the name OZ 70 that stands for Ozark Highland composite of 70 collections. The release has been delayed until a solution can be found for its high seed dormancy.

Seed was harvested from the wind barrier block and an increase planting will be made in 2001.

## **2001**

The increase plot of 9078831 (OZ 70) was again expanded in 2001 but this year it was planted the first week of March to allow for stratification. Seed harvested in 2000 was used in the planting because seed less than one year old appears to have more dormancy than seed that has had time in storage. The portion of the plot that was planted in 2001 established well and even produced a small amount of seed the first year.

Seed harvested from the wind barrier accession was propagated in the greenhouse and transplanted into an evaluation nursery. The evaluation nursery has approximately 250 plants on a three-foot grid. These plants will be evaluated for two additional years for height, biomass production and lodging. This plant will be released as a tall, stiff stemmed selection.

## **2002**

Field testing has shown possible problems with establishment of OZ 70 big bluestem. A trial was started using replicated plots to compare the establishment of OZ 70 with Rountree big bluestem. First year data indicates that Rountree establishes quicker with higher stand density than OZ-70. It also indicated that the winter dormant plots (planted March 14, 2002) of OZ 70 were better than the spring planted plots (planted June 21, 2002). This was reversed with the Rountree. This information supports the high seed dormancy problem indicated in seed tests. These plots will be monitored one more year to see if the slow establishment has to do with the long-term density of the plots.

A comparison between new seed and one-year-old seed is planned for 2003. Seed tests indicate a problem with seed dormancy in new seed. Storage for one year could help rectify this problem.

**Study 29I097G - Assembly and Evaluation of Big Bluestem,  
*Andropogon gerardii*, Vitman.**

**Table #1**

**Accessions Selected for Crossing Block**

<u>Collector</u>	<u>State</u>	<u>County</u>	<u>Accession Number</u>	<u>MLRA</u>	<u>Soil</u>
Levonna S. Vekman	Arkansas	Faulkner	9056956	118	Leadville
Mark L. Kennedy	Arkansas	Fulton	9056968	116A	Geesville
Luther O. Shaw	Arkansas	Izard	9056920	116A	Mako
NRCS-Field Office	Arkansas	Logan	9056964	118	Taff
NRCS-Field Office	Arkansas	Madison	9056962	118	Leadvale
Stephen T. Ford	Arkansas	Madison	9056945	117	Nixa-SL
John Y. Harrington	Arkansas	Madison	9056923	116A	Estate-SC
John Y. Harrington	Arkansas	Madison	9056952	116A	Estate-SC
Lane L. Gentry	Arkansas	Perry	9056922	119	Clebit
John D. Kopf	Arkansas	Scott	9056936	119	Carnasaw
Jeremy R. Funk	Arkansas	Sharp	9056914	116A	Gepp
NRCS-Field Office	Arkansas	White	9057058	118, 134	
NRCS-Field Office	Arkansas	White	9057060	118,134	
Robert S. Garner	Arkansas	Yell	9056908	119,118	Clebit-FSL
H. Dan Philbrick	Missouri	Barry	9056832	116B	
Dudley W. Kaiser	Missouri	Benton	9056840	116B	Bardley
NRCS-Field Office	Missouri	Camden	9056724	116A	Gateway
William K. Quage	Missouri	Cedar	9056800	116B	Hector
Patricia A. Beneke	Missouri	Cole	9056821	115	Goutewood
Patricia A. Beneke	Missouri	Cole	9056806	115	Gateway
Melodie Marshall	Missouri	Crawford	9056820	116B	
Melodie Marshall	Missouri	Crawford	9056886	116B	
Melodie Marshall	Missouri	Crawford	9056767	116B, 116A	Lebanon
Myron C. Hartzell	Missouri	Dent	9056773	116B	Coulstone
Myron C. Hartzell	Missouri	Dent	9056763	116B	Lebanon
John L. Lumb	Missouri	Douglas	9056833	116B	Doniphan
Art Kitchen	Missouri	Franklin	9056855	115	Crider
Art Kitchen	Missouri	Franklin	9065771	115	Union
NRCS-Field Office	Missouri	Gasconade	9056848	116B	Gladden
Clayton P. Robertson	Missouri	Gasconade	9056875	116B	
H. Lane Thurman	Missouri	Greene	9056716	116B	Chirty Silt Loam
NRCS-Field Office	Missouri	Hickory	9056839	116A	
Stanley Lamb	Missouri	Iron	9056774	116A	Midco
Howard Combes	Missouri	Howell	9056753	116A	Doniphan
Joe H. Everett	Missouri	Jefferson	9056842	115	GL
NRCS-Field Office	Missouri	LaCled	9056741	116A	Cherty Silt Loam
Kees VanderMer	Missouri	LaCled	9056791	116A	Union
Cecile Allen	Missouri	Lawrence	9056709	116B	Viraton
Ron R. McMurtrey	Missouri	McDonald	9056719	116A	
Larry E. Lewis	Missouri	Miller	9056732	116B	SIL
Larry E. Lewis	Missouri	Miller	9056868	116B	SIL
Henry E. Knipker	Missouri	Moniteau	9056890	116B	Glensted
Mary Beth Roth	Missouri	Morgan	9056831	116B	

**Study 29I097G – Assembly and Evaluation of Big Bluestem, *Andropogon gerardii*, Vitman.**

**Table #1 - continued**

<u>Collector</u>	<u>State</u>	<u>County</u>	<u>Accession Number</u>	<u>MLRA</u>	<u>Soil</u>
Mary Beth Roth	Missouri	Morgan	9056837	116B	
Stephen E. Robbins	Missouri	Organ	9056770	116A	
William R. Dilbeck	Missouri	Polk	9056828	116B	
NRCS-Field Office	Missouri	Pulaski	9056746	116A	Wilderness
Clarence Wagy	Missouri	Reynolds	9056701	116A	
Charles E. Johnson	Missouri	Ripley	9056895	116A	
Charles E. Johnson	Missouri	Ripley	9056894	116A	
Steve Wall	Missouri	Shannon	9056762	116A	
Claude A. Peifer	Missouri	Ste. Genevieve	9056819	116B	Bloomsdale
Edward L. Templeton	Missouri	St. Francois	9056845	116A	Crider
Carl Wehrman and Dude Davidson	Missouri	Taney	9056712	116A	Clarksville
Jeff A. Lamb	Missouri	Texas	9056728	116A	Goss
NRCS-Field Office	Missouri	Wayne	9056854	116A	
Patrick L. Adams	Missouri	Washington	9056817	116A	Silty Clay Loam
Patrick L. Adams	Missouri	Washington	9056870	116A	Silty Clay Loam
John N. Emerson	Missouri	Webster	9056737	116B	
Dan D. Divine	Missouri	Wright	9056733	116B	
Andrew R. Inman	Oklahoma	Adair	9056996	117	Hector Complex
Billy D. Dudley	Oklahoma	Cherokee	9057010	116A, 117	Newtonia
Billy D. Dudley	Oklahoma	Cherokee	9057016	116A, 117	Talpa-Rock
Kenneth W. Swift	Oklahoma	Choctaw	9057025	112	Muskogee SL
Warren R. Sanders	Oklahoma	Coal	9057005	119	Boham
Steve D. Clark	Oklahoma	Latimer	9057014	118, 119	Stigler SL
Robert E. Blackman	Oklahoma	Mayes	9056995	112, 116A	Hector
Sam L. Viles	Oklahoma	McIntosh	9057035	118	Karma SL
Patrick I. Bogart	Oklahoma	Okmulgee	9057032	112, 118	Taloka SL
Patrick I. Bogart	Oklahoma	Okmulgee	9057037	112, 118	Taloka SL
NRCS-Field Office	Oklahoma	Ottawa	9057030	116A, 112	ETA-SL
William R. Bin	Oklahoma	Pushmataho	9957052	119	Bosville
William R. Bin	Oklahoma	Pushmataho	9057046	119	Bernow FSL

**Wind Barrier Selection Isolation Block**

**Table #2**

<u>Collector</u>	<u>State</u>	<u>County</u>	<u>Accession Number</u>	<u>MLRA</u>	<u>Soil</u>
	Arkansas	Logan	9056960	118	Laedvale

**Study 29I097G – Assembly and Evaluation of Big Bluestem, *Andropogon gerardii*, Vitman.**

**Landscape Selection Rod Row Area**

**Table #3**

<u>Collector</u>	<u>State</u>	<u>County</u>	<u>Accession Number</u>	<u>MLRA</u>	<u>Soil</u>
Clarence Wagy	Missouri	Carter	9056703	N116A	Opequon
Clarence Wagy	Missouri	Reynolds	9056708	N116A	Clarksville
Myron Hartzell	Missouri	Dent	9056812	116A	Elsah
Kenneth W. Swift	Oklahoma	Latimer	9057025	119	Freestone Variant - Bernow Variant Complex
	Oklahoma	McCurtain	9057049	1336	Kinta Clay Loam
Dennis W. Shirk	Missouri	Maries	9056877	116A	Lebanon
Larry B. Cash	Arkansas	Carroll	9056934	116A	Nixa

**Study: 29I101J**

**Study Title:** Assembly and Evaluation of Arrowwood, *Viburnum dentatum* L.

**Study Leader:** Henry, J.

**Introduction:**

Arrowwood is an upright bushy shrub to five meters; bracts are glabrous, becoming gray; leaves suborbicular to ovate, 3-8 cm long, short acuminate, rounded or subcordate, coarsely dentate, glabrous and lustrous above, glabrous beneath or bearded in the axils of the veins, with 6-10 pairs of veins; petiole 1-2.5 cm long; cymes slender stalked, 5-8 cm across, glabrous; stamens longer than corolla. Flowers are globose-ovoid, 6 mm long, blue-black.

**Problem:**

There is a need for developing arrowwood for use as wildlife food and habitat in the three states being served by the center.

**Objective:**

The objective is to assemble, comparatively evaluate, select and release an adapted cultivar of arrowwood.

## **Discussion:**

### **1988-1992**

Collections were requested from the three-state service area but only nine were made. There was concern regarding the correct species being collected because of its rare occurrence in the service area according to the literature reviewed. The collections were stratified and placed in the greenhouse for germination but none germinated.

### **1993**

One hundred and fifty plants were obtained with a field collection origin in the state of Iowa. These plants were planted in Field #7e in May 1993. All plants were surviving in good to excellent condition up to the time of the great flood of 1993.

Approximately eight and a half feet of floodwater inundated this planting. Once the floodwaters receded, it became apparent that the entire planting was destroyed.

More plants will be sought for possible replacing in 1994 or 1995.

### **1994**

This project was reestablished April 25, 1994 in Field #11e at the PMC. There was no seed from native collections available at this time so six accessions of plant materials were purchased from nursery production stock. Three accessions were named and three were common stock with origins from Iowa and Illinois. The summer of 1994 experienced several significant dry periods and although they were hand watered several times, some replanting of the smaller plants was necessary.

### **1995-1996**

The planting was evaluated for survival, height, spread, and form. Survival of five of the six accessions was excellent. The Iowa source was established with smaller plants but had only about 60% survival.

### **1997-1999**

Accession 9062310, origin Iowa; source, Forrest Keeling Nursery, was selected based on the following characteristics: seed production, insect and disease resistance and form. Seed of this accession was harvested in 1997, 1998 and 1999 and propagated in the PMC greenhouse. These plants will be used in field plantings in Iowa starting in the spring of year 2003. Plans are to release this accession as a selected class germplasm in year 2004-2005.

### **2000**

Plans were to release accession 9062310, arrowwood in year 2001 but because of the need for field planting evaluations to support this release; the release date will need to be

put off until at least 2004 or 2005. Nine ounces of clean seed were harvested from the planting located in Field #11 on the PMC on July 19, 2000. Seed was matured and had begun to shatter at the time of harvest. This accession will be evaluated in field plantings only in the state of Iowa.

The source of this accession (9062310) of arrowwood is Floyd County, Iowa near Charles City.

### 2001

The selected accession of arrowwood (9062310) produced a medium amount of seed this year (0.33 pound). The seed was harvested on July 9, 2001 from a planting located in Field 11 on the PMC. This accession will be placed in field plantings only in the state of Iowa in 2002.

### 2002

Accession 9062310 from Floyd County Iowa was selected from the *Viburnum dentatum* L., arrowwood assembly. Seed was harvested from this selection on July 22, 2002. This selection produced 1.3 pounds of clean seed. The following is a listing of seed production by year:

<u>Year of Harvest</u>	<u>Amount of Seed Harvested</u>
2000	9.00 ounces
2001	0.33 pound
2002	1.30 pounds

**Study:** 29I107G

**Study Title** - Assembly and Evaluation of Eastern Gamagrass, *Tripsacum dactyloides*, L.

**Study Leader:** Bruckerhoff, S. B.

#### **Introduction:**

Eastern gamagrass, *Tripsacum dactyloides* L., is a tall warm season perennial grass found from Florida to Texas and Mexico, north and west to Massachusetts, New York, Michigan, Illinois, Missouri, Iowa and Nebraska. Eastern gamagrass grows in large clumps with thick rhizomes, broad flat leaves, the staminate and pistillate flowers in separate parts of the same many-flowered spikes. The pistillate spikelets are solitary and occur in hollowed portions on opposite sides of the thickened hard joints of the lower part of the rachis; this pistillate portion breaks up at maturity into several one-seeded joints. The staminate spikelets are two-flowered and in pairs on one side of a continuous rachis. Eastern gamagrass occurs on prairies, open limestone slopes, borders of woods and thickets, fields, and along roadsides and railroads. Refer to literature review.

**Problem:**

Eastern gamagrass is high quality forage with few available varieties and none of local origin in the PMC service area. There is need for a better-adapted variety of eastern gamagrass for pasture and range seedings, silage production, recreational area development and other conservation uses in the Midwestern and Eastern states for summer forage and vegetation.

**Objectives:**

The objective is to assemble, evaluate (identify superior plants), develop and release an adapted variety and or varieties of eastern gamagrass for conservation use in Missouri, Iowa, Illinois, Indiana and Ohio.

**Procedure:**

The assembly consists of vegetative material from adapted ecotypes primarily from the three-state service area. Additional collections came from Indiana, Ohio, Tennessee, Kentucky, and eastern Nebraska. The targeted collection area included the following Major Land Resource Areas: 103 (south), 104 (south), 105 (south), 106 - 115, 121, 122, 125, 126, 128, 131 (north), and 134 (north). Four collections from four different sites per county were requested. When possible, collections should come from different soil textural types.

Vegetative collections were taken from natural prairie stands or prairie remnants. The intent was to get a broad genetic base of plant material; therefore, attempting to get as diverse sampling as is practical when selecting superior eastern gamagrass plants in the field. Vegetative collections were taken from typical natural areas, prairies, borders of woods, thickets, and along roadsides and railroads. Areas that may have been seeded were avoided.

The samples were collected when the plant was dormant in the fall, divided into plantlets in the winter and placed into square open bottom containers and grown out in the greenhouse. Twelve plants per accession were planted.

The plants were planted in a randomized complete block with three replications. Each plot had three plants and all plants were planted on four-foot centers. A border row was planted around the three replications. This study was planted into a clean tilled seedbed with recommended fertility and weed control. Plants were evaluated for survival, vigor, height, spread, disease and insect resistance, lodging, amount of seed production, plant phenology, forage quantity, and regrowth.

## **Discussion:**

### **1989-1990**

The collection of samples went very well the fall of 1989. Two hundred forty-three samples were collected over a seven-state area. The primary area of collection was Missouri, Iowa, and Illinois with the majority coming from Missouri. Other states sending collections were Nebraska, Tennessee, Indiana, and Virginia.

During February 1990, each sample was cut apart and planted into 2 7/8-inch square by 5 1/2-inch tall open bottom containers for root development by air pruning. Twelve plants of each accession were planted and grown out in the greenhouse. The week of May 7, 1990, the plants were transplanted into a randomized complete block with three replications and three plants per replication. Extra plants were used for the border rows. The study was established at the PMC in Field #7F.

### **1991-1992**

The planting was evaluated several times throughout 1991. Evaluations were made for survival, vigor, disease and insect resistance, amount of seed production, plant phenology, lodging, and size, height, width, and amount of foliage.

The planting was again evaluated in 1992 with an emphasis on amount of regrowth after clipping and late season vigor.

### **1993**

The planting was evaluated in 1993 but was also destroyed by the flood. Before the planting was inundated with approximately eight feet of floodwater, PMC personnel were able to vegetatively remove 45 accessions that were rated the best and replanted them (July 2, 1993) to an upland site. The 45 accessions (Table #1) were selected based on their performance documented with three years of evaluation data. The plants were transplanted during a poor time of year but with irrigation they all survived.

### **1994-1996**

The 45 best accessions were evaluated for forage quality and quantity, phenology, and number of chromosomes. Selections of the top five to ten accessions will be made in early 1997 from data taken in 1995 and 1996 (Table # 2). The plants will be increased in the greenhouse and planted into a crossing block in 1997.

### **1997-1998**

Based on the evaluations of the 45 plants that were saved, the best 13 (Table # 2) were increased in the greenhouse and planted in Field # 6. There was only one plant per accession of these 45 plants that were evaluated, so additional plants were planted for future consideration.

The top four rated diploids, 9061911, 9061984, 9061991, and 9061948 were increased vegetatively in the greenhouse and planted in an isolation block in Field # 7F. This block will be harvested and used as a breeder block for a possible varietal release. Seed from this block will be used to start an increase planting and to also start a new evaluation

nursery for recurrent selection. The accession 9061911 was also established in an isolation block by itself as the top diploid and will be compared against the composite. The accession 9061924 was also planted in an isolation block and will be evaluated as a possible northern source as it was the best northern collection and might be best suited for Northern Missouri and Southern Iowa.

Increase plots of the two top rated tetraploids, 9061944 and 9062018, were also established from vegetative material started in the greenhouse.

### **1999**

The composite of the four top rated diploids (9061911, 9061984, 9061991, and 9061948) were assigned the accession number 9083214. Seed was harvested in July and will be used for advanced testing and to also start an increase (foundation) field. Seed was also harvested from the following increase plots: 9061911, 9061924, 9061944, and 9061984.

### **2000**

An increase (foundation) field was planted May 15, 2000, for accession 9083214 using stratified seed. The planting was small and will be expanded in 2001. It did not produce seed in 2000 and was also thin. Accessions 9083214 (composite of the four best diploids), 9061911 (the best diploid), and 9061924 (best northern diploid) were propagated in the greenhouse for use in the advanced study of eastern gamagrass with Agricultural Research Service (ARS) in Woodward, Oklahoma (study MOPMC-P-003-PA, WL). The two best tetraploids (1944 and 9062018) were also propagated in the greenhouse but did not germinate. Seed was harvested from the breeders' blocks of all the above mentioned accessions.

### **2001**

The increase (foundation) field for the accession 9083214 was expanded in 2001 but the stand was thin the first year. The seed was wet treated for stratification and planted April 18, 2001. Two rows of plants propagated in the greenhouse from stratified seed were planted alongside the increase planting. These plants were transplanted in mid April and performed poorly early due to cool weather.

A crossing block in Field #6 consisting of eight diploid accessions was also harvested in 2001. This block contained accession numbers 9061991, 9061948, 9062005, 9062085, 9061937, 9061911, 9061924, and 9061984. Seed from this cross will be tested in study MOPMC-P-003-PA, WL. This composite was assigned the accession number 9083237. Plants from seed grown from this composite will be planted in an evaluation nursery at the PMC.

### **2002**

An evaluation nursery of the composite 9083237 was started in Field #13 at the PMC. Additional seed from the crossing block in Field #6 was treated and propagated in the greenhouse while selecting for quick emergence.

**Study 29I107G -Selected Accessions of Eastern Gamagrass**

**Table #1**

<u>Collector</u>	<u>State</u>	<u>County</u>	<u>Accession Number</u>
Patrick L. Adams	Missouri	Clinton	9061968
Christopher C. Bordon	Illinois	Calhoun	9062012
William L. Brouk	Missouri	Benton	9061948
Dennis J. Browning	Missouri	Daviess	9061896
Dennis J. Browning	Missouri	Daviess	9061897
Paul Frey	Missouri	Dallas	9062082
Paul Frey	Missouri	Dallas	9062085
Darin W. Gant	Missouri	Stoddard	9061991
C. Mark Green	Missouri	Christian	9062032
Kenneth N. Gruber	Missouri	Rodaway	9061924
Terry A. Gupton	Tennessee	Roane	9034521
Robert T. Hagedorn	Missouri	Johnson	9061940
Thomas J. Hagedorn	Missouri	Pettis	9061911
Montie b. Hawks	Missouri	DeKalb	9061970
Montie B. Hawks	Missouri	DeKalb	9061971
Lynn A. Jenkins	Missouri	Newton	9062005
Lynn A. Jenkins	Missouri	Newton	9062006
David V. Johnson	Missouri	Worth	9061957
Arthur P. Kitchen	Missouri	Franklin	9062071
Viletta F. Langston	Missouri	Stone	9062034
Bob McClenny	Virginia		9034551
Steve A. McMillin	Missouri	Butler	9061994
D. Scott Patterson	Missouri	Cass	9061944
Al Peifer	Missouri	Perry	9061995
Lisa A. Ptasnik	Illinois	Massac	9062015
Lisa A. Ptasnik	Illinois	Massac	9062018
Shepherd Farms	Missouri		9061869
Shepherd Farms	Missouri		9062048
Shepherd Farms	Missouri		9062089
James E. Sturn	Missouri	Mercer	9061892
Edward L. Templeton	Missouri	St. Francois	9061999
Edward L. Templeton	Missouri	St. Francois	9062002
USDA-NRCS-Quicksand-PMC	Tennessee	Anderson	9034501
USDA-NRCS-Quicksand-PMC	Tennessee	Anderson	9034502
USDA-NRCS-Quicksand-PMC	Tennessee	Anderson	9034503
USDA-NRCS-Quicksand-PMC	Tennessee	Anderson	9034504
Curtis W. Walker	Missouri	Andrew	9061923
Stan Wall	Missouri	Shannon	9061992
Stan Wall	Missouri	Shannon	9061984
Ed J. Weilbacher	Illinois	Randolph	9062010
David L. White	Iowa	Wayne	9061876
Melvin Womack	Indiana	DuBois	9062069
Darrel D. Wright	Nebraska	Pawnee	9061887
David L. Wright	Missouri	Hickory	9061906
David L. Wright	Missouri	Hickory	9061937

**Study 29I107G - Assembly and Evaluation of Eastern Gamagrass, *Tripsacum dactyloides*, L.**

Top Rated Accessions							Table #2
Percent Protein							
Accession	Ploidy	Percent Protein			Regrowth_3/	Regrowth	
Number	Level	5/3/96	6/27/96	7/19/96	8/27/96	10/15/96	
9061911	Diploid	17.2	12.0	7.5	11.0	5.9	
9061984	Diploid	19.4	11.7	9.3	13.5	8.1	
9061991	Diploid	17.3	11.1	9.3	11.1	8.2	
9061948	Diploid	17.3	11.4		13.2	7.5	
9062005	Diploid	17.3	11.7	8.6	11.7	9.5	
9061924	Diploid	17.0	10.3	7.2	11.6	7.8	
9062085	Diploid	16.9	11.0	7.0	9.4	8.8	
9061937	Diploid	18.8	14.1	6.9	13.0	6.5	
Pete	Diploid	11.6	7.0	5.3	11.0	5.2	
9061944	Tetraploid	15.6	10.1	8.8	11.7	7.6	
9062018	Tetraploid	18.4	9.4	7.0	11.0	8.7	
9061994	Tetraploid	16.0	10.0	6.3	11.0	9.1	
9061999	Tetraploid	18.2	13.3	7.7	12.2	9.0	
9062032	Tetraploid	16.7	11.6	9.0	10.2	9.4	
Accession	First	_1/	_2/		_3/	_4/	
Number	Seedhead Emergence	Forage Quantity	Vigor	Forage Height (ft)	Forage Regrowth	% Seed Fertility	
9061911	6/16/96	1	1.3	5.0	1	59.6	
9061984	6/16/96	1	1.6	5.3	2	41.5	
9061991	6/24/96	1	2.0	5.0	1	66.9	
9061948	6/8/96	2	2.0	5.0	2	71.7	
9062005	6/8/96	2	2.8	4.9	4	82.7	
9061924	6/10/96	2	1.9	4.0	1	75.9	
9062085	6/1/96	5	1.9	4.3	3	83.3	
9061937	6/1/96	3	3.0	4.5	4	85.2	
9061944	6/24/96	3	2.1	4.8	1	76.4	
9062018	7/1/96	2	2.3	4.3	3	59.6	
9061994	7/1/96	3	2.7	4.4	3	67.6	
9061999	6/24/96	3	2.9	4.4	4	68.4	
9062032	6/24/96	2	2.1	4.7	3	67.7	
_1/ Forage quantity was a visual 1 to 9 rating with 1 being the best.							
_2/ Vigor was a visual 1 to 9 rating of overall condition of the plant with 1 being the best.							
This is an average of 10 evaluations throughout the growing season.							
_3/ All plants were clipped to an 8 inch height on 7/22/96 and plants were rated for amount of regrowth on a 1 to 9 scale. Samples of regrowth were sent in for analysis.							
_4/ Percent of 400 seed that are viable; 100 seeds harvested four times at one week intervals.							

**Study: 29I108G**

**Study Title:** Assembly and Evaluation of Low Growing, Rhizomatous Switchgrass, *Panicum virgatum* L. for Use in Waterways, Filter Strips and Other Conservation Uses.

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

Switchgrass is a warm-season, perennial, native grass. Plants are usually green or glaucous, with numerous scaly creeping rhizomes. Culms are erect, tough and hard, one to two meters rarely to three meters tall; sheaths glabrous; blades 10-60 centimeters long, three to 15 millimeters wide, flat glabrous, or sometimes pilose above or near the base, rarely pilose all over; panicle 15-50 centimeters long; acuminate; first glume clasping, two-thirds to three-fourths as long as the spikelet. Switchgrass frequents a wide variety of habitat, usually sunny including dry or moist prairies, moist seepage of rocky glades and buff escarpments, gravel bars of streams, open woods and along railroad tracks.

**Problem:**

There is a need for an adapted variety of a dense low growing, strongly rhizomatous switchgrass for use in waterways, filter strips, and for other conservation uses in Missouri, Illinois, Iowa, and adjacent states.

**Objective:**

The objective is to assemble, select, and develop a dense low growing strongly rhizomatous switchgrass, with good seedling vigor and seed characteristics, for use in waterways and streambank corridors.

**Procedure:**

The assembly consists of the collection of vegetative material from adapted ecotypes in Iowa, Illinois, and Missouri. The targeted collection area includes the following Major Land Resource Areas: 102b, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 131, and 134. Five collections from each NRCS administrative area were requested.

Vegetative collections were taken from natural prairie stands, prairie remnants or individual short growing plants growing in areas that are seasonally wet like a waterway. Total height of the plant was to be no more than three feet.

The samples were collected when the plant was dormant in the fall, divided into plantlets in the winter and placed into square open bottom containers and grown out in the greenhouse. Twelve plants per collection were grown out in the greenhouse.

The plants were planted into a randomized complete block with three replications. Each plot had three plants and all plants were planted on a four-foot spacing. A border row was planted around the three replications. This study was planted into a clean tilled seedbed with recommended fertility and weed control. Plants were evaluated for survival, vigor, height, and spread that included rhizomatous characteristics, disease and insect resistance, lodging, and seed production.

## **Discussion:**

### **1990-1991**

The collections of *Panicum virgatum* L., low growing highly rhizomatous switchgrass was initiated in November 1990 and extended through 1991. One hundred eighteen collections were obtained from Major Land Resource Areas 102B-116, 131 and 134 in Missouri, Illinois and Iowa. The total number of collections received was 22-Illinois; 28-Iowa and 68-Missouri. All collections were assigned accession numbers and stored in a cool damp building.

### **1992-1993**

The collections were vegetatively propagated in cone-tainers and placed in the greenhouse in January 1992. These plants were then transplanted in Field #7c on the PMC on June 9, 1992, in a randomized complete block with three replications. Baseline evaluations were taken this year; survival, spread, height, and number of panicles per plant. More detailed evaluations were scheduled for succeeding years.

Beginning in July 1993, the great flood began inundating the area where this project was located. Prior to the flooding of this site (July 2, 1993), additional evaluations were started and 67 accessions were vegetatively moved to an upland site on the PMC for continued evaluation. Table #1 lists the selected accessions, origins, and collectors.

### **1994-1995**

Evaluations were continued on the 67 accessions during 1994 and 1995. The original planting in Field #7c that was flooded in 1993 was also checked for survivors. The planting was flooded by as much as eight feet of water for almost eight weeks. Nine plants were found that showed life and were dug up and moved to an upland site. These nine plants represented three accessions (Table #2).

Five accessions were selected out of the block of 67 for a short growing rhizomatous type. The five accessions (Table #3) were allowed to cross and seed was harvested and grown out in the greenhouse. The five accessions were also dug and increased in the greenhouse in containers.

### **1996**

The five selected accessions (Table #3) were planted into a crossing block June 26, 1996. Half the block was from clonal material from each of the five accessions and the other half was from seed harvested from each of the five plants that were allowed to cross with each other. The accessions of each half of the planting were replicated five times with five plants per replication. Unwanted plants will be eliminated and the remainder of the block will be used for seed increase.

## **1997-1998**

The three accessions (Table #2) of flood tolerant switchgrass were vegetatively increased in the greenhouse. Approximately 250 plants were transplanted April 1997 in Field #7. This is now the breeders' block for the accession 9083170 that is a composite of the three accessions listed in Table #2. Seed was harvested from this plot the first year and used to start a small increase plot in 1998. A small amount of seed was harvested from this increase plot the first year. It is also planned to increase the size of this plot in 1999.

The low growing switchgrass block containing five accessions (Table #3) was again evaluated in 1997. Thirty-five plants were selected from the block of 250. Selected plants were allowed to cross and produce seed. This seed was also used to start an increase field in 1998. This small increase plot produced minimal seed the first year. Seed was again harvested from the 35 plants in 1998 and will be used to make the increase plot size bigger in 1999. The 35 selected plants are the breeder's block for the new accession 9083172 that is a composite of the five accessions in Table #3.

## **1999**

The increase plot of flood tolerant switchgrass, accession 9083170, was expanded in May 1999. This planting did not do well, possibly poor seed germination combined with a very dry summer. Weed control was also poor. Establishment of field plantings was also poor. Expanding the increase plot will again be planned for 2000. Seed was harvested from the breeder's block and the 1998-increase plot. This seed was small due to dry weather.

The increase plot of low growing switchgrass, accession 9083172, was also expanded in May 1999. This planting also did poorly, again possibly poor seed germination combined with a very dry summer. Weed control was also poor. Field testing will begin when seed becomes available. Expanding the increase plot will be planned for year 2000. Seed was harvested from the original 35-plant breeder's block and also the increase field. This seed was also small due to dry weather.

## **2000**

Increase plots of the flood tolerant switchgrass, accession 9083170, and the low growing switchgrass, accession 9083172, were again planted in 2000. These plantings were very sparse and slow to establish. The plantings made in 1999 contained some plants with minimal seed produced. Plantings will again be tried in 2001 with more stratification.

## **2001**

The increase plots of the low growing switchgrass, accession 9083172, that were planted in 1998 and 1999 have filled in and produced seed. The plots planted in 2000 and 2001 have failed. This accession appears to have high seed dormancy and combined with excessive weed competition caused poor establishment. An increase planting is planned for 2002 on an upland site with less weed problems.

The increase plots of the flood tolerant switchgrass, accession 9083170 that was planted in 1998 produced seed in 2001. The 1999 planting was very thin and the 2000 and 2001 plantings have failed. This accession appears to have high seed dormancy. Another increase planting is planned in 2002 with additional stratification.

Accession 9062244 was observed in the nursery block in field eight as having high forage production (very leafy), medium height, and late maturity. Protein analysis of a sample taken was 15.6%. This plant was increased in the greenhouse from vegetative material and planted into a 200-plant nursery in 2000. Unwanted plants were rogued out and seed was harvested in 2001. Plants that germinate quicker from the heaviest seed will be placed in an evaluation nursery in 2002.

## **2002**

The low growing switchgrass, accession 9083172, increase plots had limited seed production in 2002. The 17.3-pound bulk seed produced will be used in the field-planting program for advanced testing. An additional 1.5 acres increase field was planted in 2002. No seed was harvested the establishment year from this plot.

The flood tolerant switchgrass, accession 9083170, increase plots also had limited seed production in 2002. The 32.5-pound bulk seed produced will be used in the field-planting program. Due to an extremely wet spring, no additional seed increase field was planted in 2002.

The medium height forage type switchgrass, accession 9062244, was propagated in the greenhouse and plants were selected for quick establishment and seedling vigor. These plants were transplanted into an evaluation nursery in Field #1 at the PMC.

## Study 29I108G-Selected Accessions of Low Growing Switchgrass

Table #1

<u>Accession #</u>	<u>State</u>	<u>County</u>	<u>MLRA</u>	<u>Collector Name</u>
9062155	Iowa	Louisa	108	Dean L. Pettit
9062157	Iowa	Cherokee	107	Lon Allan
9062158	Iowa	Clay	103	John P. Vogel
9062160	Iowa	Freemont	107	NRCS F. O.
9062163	Iowa	Hamilton	103	Dana C. Holland
9062165	Iowa	Woodbury	107	John P. Vogel
9062166	Iowa	Monona	107	Michael J. Kuera
9062178	Iowa	Muscatine	108	Douglas S. Johnson
9062181	Illinois	Champaign	108	Leon W. Wendt
9062188	Illinois	Macoupin	108	Ivan N. Dozier
9062189	Illinois	Macoupin	115	Ivan N. Dozier
9062190	Illinois	Macoupin	108	Ivan N. Dozier
9062195	Illinois	Carroll	105	Raymond J. Hudak
9062196	Illinois	Carroll	105	Raymond J. Hudak
9062205	Missouri	Barton	112	Jerry L. Cloyed
9062207	Missouri	Bates	112	Robert D. Bouland
9062208	Missouri	Pettis	116A	Thomas J. Hagedorn
9062209	Missouri	Christian	116A	C. Mark Green
9062211	Missouri	Ozark	116A	Carroll W. Foster
9062212	Missouri	Johnson	112	Robert T. Hagedorn
9062213	Missouri	Madison	116A	Sandra L. Lewis
9062214	Missouri	Ste. Genevieve	116B	Renee L. Phillips
9062215	Missouri	Oregon	116A	Stephen E. Robbins
9062216	Missouri	Shannon	116A	Steve Wall
9062217	Missouri	Reynolds	116A	Clarence W. Wagy
9062218	Missouri	Christian	116A	C. Mark Green
9062219	Missouri	Perry	116B	Claude E. Peifer
9062220	Missouri	Reynolds	116A	Clarence W. Wagy
9062221	Missouri	Dade	116B	Todd E. Mason
9062222	Missouri	Morgan	116B	James A. Maberry
9062223	Missouri	Franklin	116B	Arthur P. Kitchen
9062224	Missouri	Cedar	116B	Kim C. Ehlers
9062225	Missouri	Christian	116A	C. Mark Green
9062227	Missouri	Ozark	116	Carroll W. Foster
9062228	Missouri	Texas	116	Jeff A. Lamb
9062229	Missouri	Texas	116	Jeff A. Lamb
9062234	Missouri	Saline	107	Wayne E. McReynolds
9062237	Missouri	Ray	107	James M. Rehmsmeyer
9062238	Missouri	Worth	109	David A. Stevens
9062239	Missouri	Sullivan	109	Stuart A. Lawson
9062240	Missouri	DeKalb	109	Wm. A. Throckmorton

<b>Table #1 - continued</b>				
<u>Accession #</u>	<u>State</u>	<u>County</u>	<u>MLRA</u>	<u>Collector Name</u>
9062242	Missouri	DeKalb	109	Wm. A. Throckmorton
9062243	Missouri	Buchanan	107	Rodney Saunders
9062244	Missouri	Dent	116	Myron C. Hartzell
9062246	Missouri	Sullivan	109	Stuart A. Lawson
9062247	Missouri	Buchanan	107	Rodney Saunders
9062248	Missouri	Sullivan	109	Stuart A. Lawson
9062250	Missouri	Nodaway	109	Kenton L. Macy
9062251	Missouri	Worth	109	David A. Stevens
9062252	Missouri	Daviess	109	James A. Sturm
9062253	Missouri	Daviess	109	James A. Sturm
9062254	Missouri	Maries	116A	Dennis W. Shirk
9062255	Missouri	Maries	116B	Dennis W. Shirk
9062256	Missouri	Maries	116A	Dennis W. Shirk
9062257	Missouri	Maries	116A	Dennis W. Shirk
9062259	Missouri	Shannon	116A	Steve Wall
9062261	Missouri	Shannon	116A	Steve Wall
9062265	Missouri	Sullivan	109	Stuart A. Lawson
9062267	Missouri	Gentry	109	Gary J. Barker
9062268	Missouri	Platte	107	Terry A. Breyfogle
9062269	Missouri	Sullivan	109	Stuart A. Lawson
9062270	Missouri	Platte	107	Terry D. Breyfogle
9062271	Iowa	Page	104	Kevin J. McCall
9062272	Illinois	Fayette	104	Brad S. Simcox
9062274	Iowa	Madison	108/109	Larry Beeler/Tom Oswald
9062193	Illinois	Fayette	113	Brad S. Simcox

**Selected Accessions of Wet Tolerant Switchgrass**

**Table #2**

<u>Accession #</u>	<u>State</u>	<u>County</u>	<u>MLRA</u>	<u>Collector Name</u>
9062193	Illinois	Fayette	113	Brad S. Simcox
9062213	Missouri	Madison		Sandra L. Lewis
9062235	Missouri	Miller	116	Matt L. Burcham

**Final Accessions Selected for Low Growing Switchgrass**

**Table #3**

<u>Accession #</u>	<u>State</u>	<u>County</u>	<u>MLRA</u>	<u>Collector Name</u>
9062205	Missouri	Barton	112	Jerry L. Cloyed
9062225	Missouri	Christian	116A	C. Mark Green
9062252	Missouri	Daviess	109	James A. Sturm
9062255	Missouri	Maries	116B	Dennis W. Shirk
9062257	Missouri	Maries	116A	Dennis W. Shirk

**Study No. 29I110J**

**Study Title:** Assembly and Evaluation of Choke cherry, *Prunus virginiana* L.

**Study Leader:** Henry, J.

**Introduction:**

Choke cherry is one of the most widely distributed native tall shrubs or small trees in North America. It occurs from Newfoundland south to Georgia and west to California and British Columbia. In the Midwest its habitat includes moist sites in open areas, along fencerows, roadsides, borders of woods as well as sandy or rocky hillsides and ravines. Three varieties have been described: var. *virginiana* in the eastern United States, var. *melanocarpa* in the west, and var. *demissa* along the Pacific Coast. Some forms have yellow rather than dark red or black fruit. The leaves of var. *melanocarpa* are thicker and cordate rather than oval, oblong or obovate as in var. *virginiana*. The fruit is less astringent.

Adaptive characteristics of choke cherry includes fast growth, dependable fruit crops, tolerance to harsh climatic extremes, and the ability to grow in a wide variety of soil types.

**Problem:**

There is a need for developing a cultivar/selection of choke cherry for use as wildlife food and habitat in the three states served by the Center.

**Objectives:**

The objective is to assemble, comparatively evaluate, select, and release adapted cultivars/ selections of choke cherry.

**Discussion**

**1989-1992**

Seed collection was initiated in 1989 and 11 collections were made before the State Conservationists' Advisory Committee put the study on hold in 1992. The reason for placing this study on hold was the lack of personnel at the PMC to carry out the work involved with new studies. The intent was to make 40-50 collections from the three-state service area to be placed in a randomized complete block planting.

**1993-1996**

The project remained in an inactive status until 1996. At this time a decision was reached to germinate the seed that was collected earlier. Based on the viability of this seed collection, it may become necessary to recollect this species.

**1997-1998**

Seed collections of choke cherry were stratified and placed in the greenhouse for germination (March 1997). A total of 15 collections were made but only 11 germinated. Enough plants of the 11

collections were obtained to initiate a randomized complete block planting with twelve replications. This planting was made on June 23, 1998 in Field #6 on the PMC.

### 1999-2001

Table #1 lists the accessions of choke cherry collected, collector's name, state, county, MLRA, and soil type. Table #2 reflects the plants' performance for 1999, 2000 and 2001. Plans are to continue evaluations for survival, fruit production, height, spread, insect and disease resistance and vigor until selection(s) are made. Several accessions produced light to heavy fruit production. An Eastern tent caterpillar *Malacosoma americanum* infestation was noticed throughout this planting (all accessions) in years 2000 and 2001, however there was no serious damage recorded on any accession in this assembly. A solution of Malathion (one tablespoon per gallon of water) was sprayed on all plants. Control was almost instant in both years.

### 2002

Evaluations for this study were made on April 25, July 11 and October 9. The following characteristics were documented: vigor, insect and disease resistance, height, spread, and fruit production. The eastern tent caterpillar, *Malacosoma americanum*, infested this planting again this year. No chemical (Malathion) was applied this year in order to determine the extent of damage caused by these insects. The following tables reflect the above mentioned evaluations along with accession information (Table #1)

**Table #1 Accession Information**

<b>Collector</b>	<b>State</b>	<b>County</b>	<b>MLRA's</b>	<b>Soil</b>	<b>Accession</b>
R. W. Nuboer	Illinois	Carroll	111	Seaton Silt Loam	9008107
R. W. Nuboer	Illinois	Whiteside	108	Silt Loam	9057068
R. W. Nuboer	Illinois	Carroll	111	Fayette Silt Loam	9057069
R. E. Szafoni	Illinois	McLean	108	Unknown	9057089
W. D. Glass	Illinois	Iroquois	110	Sandy Loam	9057143
J. R. Heim	Illinois	Ogle	108	Unknown	9057162
J. P. Vogel	Iowa	Woodbury	107	Kennebec	9057181
J. P. Vogel	Iowa	Woodbury	107	Kennebec Silt Loam	9068669
Maggie Cole	Illinois	Cook	110	Unknown	9068542
Jimmy Henry	Missouri	Lincoln	115	Menfro Silt Loam	9008147
J. R. Heim	Illinois	Lee	108	Martinsville Silt	9068587
Maggie Cole	Illinois	Cook	110		9068660
Maggie Cole	Illinois	Cook	110		9008157
Nancy Pals	Illinois	Coles	108		9068667
Bart C. Pals	Illinois	Effingham	113		9068183
William A Throckmorton	Missouri	DeKalb	109	Lamoni	9068668
Kent A. Boyles	Illinois	Tazewell	108	Stronghurst Silt Loam	9068664
Louis Byford	Missouri	Atchison	107	Napier Silt Loam	9068658

Study 29I110J - Assembly and Evaluation of Chokecherry														Rating for Vigor & Ins/Dis = 1-Excellent, 9-Poor										Table #2					
Row	Acc#	9068664				9008107				9008157				9069664				9068664				9068660							
1	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	1	3	3	5	3	8	7	6	2	3	3	6	2	3	3	4	2	3	3	6	2	2	2	4	2	2	2	4
	Ins/Dis	1	2	2	7	2	8	7	7	1	3	5	4	1	2	4	5	1	1	4	6	1	3	4	5	1	3	4	5
	Height	1.5	2.0	2.5	7.3	2.0	2.5	3.0	5.0	1.7	2.3	2.9	6.0	2.0	2.6	3.2	5.8	1.9	2.5	3.0	6.5	2.0	2.5	3.0	6.1	2.0	2.5	3.0	6.1
	Fruit Prod	0	0	5	0	0	0	4	0	0	0	7	0	0	8	5	0	0	0	7	6	0	6	3	0	0	6	3	0
Row	Acc#	9068664				9008107				9008157				9068660				9068183											
1	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	2	3	3	dead	3	6	5	dead	2	7	7	4	dead	0	0	0	dead	0	0	0	0	0	0	0				
	Ins/Dis	1	2	3		3	6	6		1	8	6	3	0	0	0	0	0	0	0	0	0	0	0	0				
	Height	1.9	2.6	3.3		1.8	2.0	2.5		1.8	2.4	3.2	6.1	0	0	0	0	0	0	0	0	0	0	0	0				
	Fruit Prod	0	6	3		0	7	6		0	8	4	6	0	0	0	0	0	0	0	0	0	0	0	0				
Row	Acc#	9008107				9068658				9008107				9068183				9068667				9068669							
2	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	1	2	2	7	2	4	4	6	1	4	3	6	1	3	3	2	2	3	3	6	1	2	2	7	1	2	2	7
	Ins/Dis	1	4	5	8	1	4	7	5	1	4	6	6	1	4	7	2	3	4	6	8	2	4	6	9	2	4	6	9
	Height	1.5	2.2	2.9	5.1	1.3	1.7	2.2	7.2	2.0	2.4	3.0	5.6	1.5	2.3	2.8	7.9	2.0	2.6	3.0	6.5	2.5	2.8	3.2	7.6	2.5	2.8	3.2	7.6
	Fruit Prod	0	7	5	0	0	0	6	6	0	6	6	0	0	7	7	3	7	5	3	3	6	4	2	0	6	4	2	0
Row	Acc#	9068664				9008157				9068660				9068668				9008107											
2	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	1	1	1	1	1	3	3	1	4	5	5	7	2	1	1	1	4	8	7	6	4	8	7	6				
	Ins/Dis	1	4	7	9	2	4	7	3	4	4	7	6	1	4	6	3	4	5	6	4	4	5	6	4				
	Height	1.7	2.7	3.5	10.3	1.4	2.7	3.5	10.6	1.5	2.0	2.4	4.5	1.8	2.5	3.5	9.6	1.0	1.4	1.9	7.4	1.0	1.4	1.9	7.4				
	Fruit Prod	6	4	2	0	0	6	3	5.2	0	7	7	4	7	4	3	7	0	8	6	3	0	8	6	3				
Row	Acc#	9068668				9008107				9068658				9068668				9008183				9068660							
3	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	1	2	2	3	1	1	1	4	5	5	4	4	1	1	1	3	5	7	6	6	1	2	2	5	1	2	2	5
	Ins/Dis	3	1	4	4	2	2	4	4	5	5	5	5	2	2	5	4	6	6	6	6	3	3	6	3	3	3	6	3
	Height	1.5	2.3	3.0	8.7	2.4	2.7	3.2	7.3	1.5	3	5.5	8.5	1.8	2.7	3.3	8.7	1.0	1.5	2.0	2.5	2.0	2.5	3.5	6.5	2.0	2.5	3.5	6.5
	Fruit Prod	7	6	6	3	0	8	8	6	0	0	8	5	7	7	6	3	0	8	8	8	0	8	7	5	0	8	7	5

Study 29I110J - Assembly and Evaluation of Chokecherry - cont.														Rating for Vigor & Ins/Dis = 1-Excellent, 9-Poor								Table #2			
Row	Acc#	9068667				9068664				9068658				9068669				9068664							
3	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	1	1	1	3	2	7	6	4	2	3	3	4	2	4	4	6	2	5	5	5				
	Ins/Dis	1	1	4	3	2	6	6	4	1	2	6	5	3	2	5	6	5	6	6	4				
	Height	2.0	2.3	3.2	7.3	1.4	1.9	2.2	8.3	1.7	2.4	3.1	8.5	1.5	2.2	2.8	6.8	1.3	1.8	2.3	6.3				
	Fruit Prod	7	5	5	2	0	8	6	0	0	8	6	5	7	7	6	0	0	8	6	7				
Row	Acc#	9068664				9008157				9008107				9068658				9068668				9068669			
4	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	2	6	6	7	2	4	4	2	5	4	5	3	3	3	3	3	1	1	1	1	3	5	3	6
	Ins/Dis	3	6	6	5	3	3	3	1	2	1	3	6	5	2	4	6	2	2	6	2	3	6	6	4
	Height	1.6	2.2	2.7	3.7	2.0	2.3	3.0	8.8	1.7	2.6	3.4	7	1.0	1.4	2.9	7	2.0	2.5	4.0	10.5	2.5	2.9	3.5	5.5
	Fruit Prod	0	0	7	0	8	6	6	7	0	8	6	7	7	5	3	7	6	4	2	2	0	6	4	2
Row	Acc#	9068660				9068667				9068664				9068183				9008157							
4	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
cont	Vigor	2	4	4	4	1	3	3	4	1	3	3	7	3	3	3	dead	2	4	4	2				
	Ins/Dis	2	3	5	3	2	4	4	4	1	4	5	5	3	2	4		3	2	5	1				
	Height	1.5	1.9	2.7	6.8	1.5	1.9	3.3	6.5	1.5	2.2	3.0	3.7	1.8	2.4	3.0		1.8	2.2	2.8	8.8				
	Fruit Prod	0	0	4	2	6	4	4	2	0	7	4	0	7	6	3		0	6	3	7				
Row	Acc#	9068668				9068667				9068664				9068660				9008157				9068183			
5	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	2	4	5	4	1	2	4	3	2	3	4	3	2	3	3	4	1	2	2	4	2	5	5	dead
	Ins/Dis	3	5	6	3	3	2	6	3	4	4	7	4	2	4	7	4	2	2	6	3	2	5	6	
	Height	2.0	2.3	3.1	7.8	1.5	2.2	3.1	7.1	1.5	1.9	3.0	8.8	2.5	2.9	3.6	7.1	1.8	2.7	3.3	7.3	2.0	2.4	3.1	
	Fruit Prod	7	4	3	2	7	5	5	6	0	8	7	8	0	7	7	8	0	6	4	3	0	0	7	
Row	Acc#	9068668				9068669				9068658				9008107				9068660							
5	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	1	2	2	4	2	5	5	4	3	3	3	2	4	4	4	5	3	8	6	5				
	Ins/Dis	3	2	5	3	2	6	7	7	2	4	5	3	2	2	6	5	3	8	7	6				
	Height	1.8	2.3	3.0	7.8	1.5	1.8	2.3	4.8	1.5	1.8	2.4	7.8	1.0	1.4	2.1	7.1	1.0	1.3	2.0	5.6				
	Fruit Prod	8	7	7	2	7	6	6	0	0	8	8	0	0	7	7	5	0	7	6	5				

Study 29I110J - Assembly and Evaluation of Chokecherry - cont.														Rating for Vigor & Ins/Dis = 1-Excellent, 9-Poor								Table #2			
Row	Acc#	9008107				9068664				9068660				9068669				9068658				9008157			
6	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	2	5	5	7	2	3	3	7	1	3	4	1	1	2	2	6	2	4	4	3	2	3	4	6
	Ins/Dis	1	4	5	8	1	2	4	5	1	3	4	1	2	4	6	6	1	4	7	7	1	3	5	6
	Height	2.0	2.4	3.0	4.6	1.3	1.8	2.5	6.5	1.0	1.4	2.9	10.7	1.3	1.7	3	5.2	1.3	1.9	2.5	9.4	1.9	2.5	3.5	6.9
	Fruit Prod	0	7	4	3	0	7	5	0	0	5	4	8	6	4	3	0	7	4	3	6	8	6	6	0
Row	Acc#	9008107				9008183				9068668				9068667				9008183							
6	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	4	8	6	7	1	3	3	3	1	3	3	5	2	2	2	3	3	9	3	4				
	Ins/Dis	2	8	6	8	3	2	3	3	1	4	5	6	3	4	6	3	3	3	3	3				
	Height	1.4	1.8	2.7	4.6	2.5	2.9	3.1	4.9	2.0	2.4	3.0	4	2.0	2.5	3.5	8.6	1.5	1.7	2.8	3.5				
	Fruit Prod	0	7	6	3	0	7	7	6	8	7	6	0	7	5	4	1	6	6	6	6				
Row	Acc#	9068664				90008107				9068664				9008183				9068667				9068668			
7	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	3	4	3	6	2	3	3	5	2	3	3	2	6	8	6	7	4	4	4	dead	2	1	1	1
	Ins/Dis	2	3	5	4	2	2	5	3	2	4	5	6	4	2	7	7	2	3	5		2	3	5	2
	Height	2.0	2.4	3.2	5.3	2.0	2.4	3.0	6.5	1.9	2.5	3.3	9.9	1.0	1.5	2.0	3.3	1.3	1.6	2.4		1.5	2.1	3	11.1
	Fruit Prod	6	4	3	4	0	8	7	3	0	8	6	3	7	6	3	2	6	5	5		7	5	5	2
Row	Acc#	9068658				9068669				9008157				9068660				9068664							
7	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	4	4	4	4	4	7	6	8	2	2	2	5	3	4	2	4	3	6	5	2				
	Ins/Dis	4	4	3	3	4	7	3	4	3	3	6	4	3	4	6	2	2	7	5	6				
	Height	2	4	5.5	7.2	1.3	1.7	2.5	1.8	2.0	2.4	3.2	8.1	1.5	2.2	3	7.1	2	2.4	2.9	9.9				
	Fruit Prod	3	3	2	2	5	5	5	0	0	8	7	0	0	7	5	4	0	0	6	3				
Row	Acc#	9068668				9068669				9068667				9068664				9068668				9008107			
8	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	2	3	3	dead	4	44	2	6	3	3	2	8	4	4	4	6	3	4	5	dead	5	5	5	9
	Ins/Dis	2	3	6		4	3	6	6	2	6	5	6	4	6	6	4	2	2	5		4	5	6	9
	Height	1.5	2.3	2.9		1.3	1.5	2.3	5.8	2.9	2.4	3.2	3.2	2.0	2.4	3.1	5.3	1.9	2.3	3.0		1.5	2.0	2.7	3.8
	Fruit Prod	7	4	3		3	3	3	0	6	4	3	0	0	0	6	0	0	8	7		0	8	7	-

Study 29I110J - Assembly and Evaluation of Chokecherry - cont.														Rating for Vigor & Ins/Dis = 1-Excellent, 9-Poor								Table #2			
Row	Acc#	9088157				9068660				9008138				9068658				9008107							
8	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	6	5	4	4	4	3	4	6	6	7	7	7	4	4	4	6	3	5	5	6				
	Ins/Dis	4	4	3	3	3	3	7	8	6	8	6	6	5	4	7	5	3	7	7	9				
	Height	2.3	5	6.8	8	2.0	2.2	2.9	6	1	1.3	2	2.5	2.2	2.6	3.4	6.1	1.5	2.0	2.5	6.3				
	Fruit Prod	0	0	8	8	0	7	7	8	0	7	6	6	0	8	8	7	0	7	6	0				
Row	Acc#	9008107				9068668				9068667				9068660				9008183				9068658			
9	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	3	2	3	5	4	3	2	3	3	4	4	dead	4	5	3	7	5	9	7	6	3	8	6	dead
	Ins/Dis	3	2	7	6	3	3	6	2	2	5	5		2	5	6	6	3	9	9	3	4	9	9	
	Height	1.5	2.3	2.9	7.2	1.5	2.1	3.0	5.5	1.6	2.2	2.8		1.5	2.0	2.6	5	2.0	2.3	3	5.6	1.3	1.6	2.3	
	Fruit Prod	0	6	6	0	7	4	3	0	6	4	4		0	8	7	4	0	8	7	7	0	8	7	
Row	Acc#	9068664				9008107				9068669				9008157				9008157							
9	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	3	6	6	9	4	6	5	5	9	9	9	9	2	3	3	2	3	5	5	5				
	Ins/Dis	2	7	7	8	3	8	7	6	9	9	9	9	2	2	5	3	2	5	5	3				
	Height	2.0	2.3	2.8	2.8	2.0	2.3	3.1	5.1	1.9	3.6	4	4.3	1.5	2.3	3.0	8.4	1.5	2.0	2.6	6.6				
	Fruit Prod	0	8	6	0	0	0	5	5	0	0	0	0	0	8	5	3	0	6	4	5				
Row	Acc#	9068669				9068658				9008107				9068183				9068667				9068669			
10	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	4	3	3	4	6	4	4	dead	4	4	4	3	7	9	3	4	5	5	4	6	6	4	4	9
	Ins/Dis	3	3	3	4	7	4	3		3	5	5	2	7	9	9	6	3	6	6	8	5	4	5	8
	Height	2.0	2.4	3.1	8.3	1.5	2.4	3.2		2.0	2.4	3.0	7.4	1.5	1.8	2.7	5.2	2.0	2.4	3.1	5.1	2.2	2.5	3.3	5.3
	Fruit Prod	0	7	4	0	0	8	5		8	6	6	8	0	0	7	0	0	6	6	8	8	7	5	0
Row	Acc#	9068664				9008157				9068660				9068668				9068660							
10	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
Cont	Vigor	4	4	4	5	3	4	3	9	3	2	2	5	8	8	8	8	3	3	3	3				
	Ins/Dis	2	5	6	4	3	6	6	9	2	3	5	4	8	8	9	9	2	4	5	3				
	Height	1.4	1.9	2.4	6.3	2.0	2.4	3.2	2.7	1.5	2.0	3.0	7.5	2	3.5	4	4.6	2.2	2.7	3.2	5.8				
	Fruit Prod	0	5	5	8	8	8	6	0	0	7	5	8	0	0	8	8	6	6	4	2				

Study 29I110J - Assembly and Evaluation of Chokecherry - cont.										Rating for Vigor & Ins/Dis = 1-Excellent, 9-Poor										Table #2									
Row	Acc#	9068664				9068664				9068668				9068660				9008107				9008157							
11	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	5	8	8		2	2	2		4	3	3		0	0	0		4	1	1		3	2	2					
	Ins/Dis	3	7	6		1	1	2		3	3	4		0	0	0		3	2	4		4	2	4					
	Height	2.0	2.2	3.0		2.3	2.7	3.2		1.5	2.0	2.7		0	0	0		2.0	2.5	3.1		2.0	2.7	3.5					
	Fruit Prod	0	5	5		0	5	5		0	0	7		0	0	0		8	7	7		8	8	6					
Row	Acc#	9008183				9068667				9068658				9068669				9008182											
11	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002								
cont	Vigor	4	7	7		3	6	6		4	5	4		6	8	8		4	8	7									
	Ins/Dis	1	8	7		2	7	6		5	8	7		7	8	7		2	8	8									
	Height	1.5	1.8	2.4		2.0	2.4	3.0		1.5	2.4	3.0		1.8	2.4	3.0		2.0	2.4	3.2									
	Fruit Prod	0	0	6		0	7	7		0	7	6		8	7	6		0	6	5									
Row	Acc#	9008157				9068669				9068664				9068667				9068183				9008107							
12	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	3	4	4		3	8	8		3	3	3		4	2	2		2	2	3		4	1	3					
	Ins/Dis	3	2	4		2	8	7		2	2	5		4	2	3		2	2	4		2	3	5					
	Height	2.0	2.5	3.0		1.5	1.8	2.5		2.0	2.6	3.2		1.3	1.9	2.7		1.5	2.0	2.8		1.5	1.9	2.7					
	Fruit Prod	8	7	7		8	7	6		0	6	6		0	7	7		0	0	5		8	6	6					
Row	Acc#	9068658				9068108				9008167				9008147				9068664											
12	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002								
cont	Vigor	2	1	1		5	4	4		4	2	2		4	3	4		3	4	4									
	Ins/Dis	2	2	4		7	5	6		3	1	5		2	3	5		2	3	6.0									
	Height	2.0	2.5	3.9		1.3	1.7	2.2		1.8	2.4	3.1		2.2	2.8	3.4		1.5	2.3	3.0									
	Fruit Prod	0	7	5		7	7	5		7	6	4		6	6	4		0	5	5									
Row	Acc#	9068668				9068108				9008167				9068108				9008147				9008157							
13	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	3	4	4		4	1	1		3	8	7		3	3	3		4	5	6		6	7	7					
	Ins/Dis	2	2	5		3	2	5		2	8	7		3	2	6		3	7	7		3	8	6					
	Height	1.5	2.0	2.5		1.5	2.0	3.0		1.5	1.8	2.4		2.0	2.6	3.4		1.4	1.8	2.6		1.3	1.6	2.1					
	Fruit Prod	0	8	7		7	6	4		7	6	4		7	7	6		6	6	5		8	8	5					

Study 29I110J - Assembly and Evaluation of Chokecherry - cont.										Rating for Vigor & Ins/Dis = 1-Excellent, 9-Poor										Table #2					
Row	Acc#	9068108				9008167				9008147				9068108				9008107							
13	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	3	4	3		4	3	4		3	4	4		3	5	4		3	5	5					
	Ins/Dis	2	2	4		3	5	6		2	3	6		2	6	6		4	3	5					
	Height	1.5	2.0	2.5		1.4	1.9	2.6		2.0	2.6	3.0		1.5	1.9	2.7		1.5	2.4	2.9					
	Fruit Prod	7	6	4		7	6	4		6	6	5		7	6	5		8	7	7					
Row	Acc#	9008157				9008107				9068668				9068660				9068664				9008107			
14	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002
	Vigor	0	0	0		4	2	2		4	6	5		4	3	3		3	3	3		3	2	1	
	Ins/Dis	0	0	0		3	4	5		4	4	5		3	3	5		4	5	7		2	1	3	
	Height	0	0	0		1.5	2.5	3.0		2.2	2.8	3.3		1.4	2.0	2.7		1.5	2.0	2.6		1.5	2.3	3.0	
	Fruit Prod	0	0	0		8	6	6		0	7	6		6	6	5		0	5	5		8	6	5	
Row	Acc#	9008157				9008183				9068669				9068668				9068660							
14	Year	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002	1999	2000	2001	2002				
cont	Vigor	3	4	5		3	6	6		4	6	4		4	2	2		7	4	5					
	Ins/Dis	2	5	7		2	5	6		2	5	7		3	2	5		7	4	5					
	Height	2.0	2.3	3.1		1.4	1.9	2.5		1.5	2.2	3.0		1.5	2.2	2.9		1.0	1.4	2.2					
	Fruit Prod	8	8	6		0	6	5		8	6	6		0	8	7		6	6	4					

**Study No. 29A116W**

**Study Title:** Evaluation of Miscellaneous Trees and Shrubs.

**Study Leader:** Henry, J.

**Introduction:**

The evaluation of woody plant materials on the USDA-NRCS Elsberry Plant Materials Center began in 1989. Since that time plants have been added for multiple purposes. The evaluations of these plant materials have been in cooperation with the USDA-ARS, Plant Introduction Station, Ames, Iowa; Missouri Department of Conservation; and other plant materials centers.

**Problem:**

Trees and shrubs are needed to provide for windbreaks, recreation, and multipurpose use in the Midwest Region and provide multiple wildlife benefits throughout the three-state area. New selections, collections and public and private releases need to be evaluated as potential conservation species.

**Objective:**

The objectives of this study are to assemble and evaluate woody plant materials (both collections in the wild and also released cultivars) for conservation uses, area of adaptation, and to select and increase limited quantities of promising woody plants for advanced evaluation. Superior accessions or those exhibiting unique characteristics will be placed in field evaluations and field plantings in the three-state area being served by the PMC.

**Assembly:**

Plant materials of various woody species representing many species have been planted on the PMC. The sources include other PMC's, commercial nurseries, and other agencies.

**Discussion:****1994-2002**

This study is a long-term ongoing evaluation of miscellaneous trees and shrubs that were not part of a collection made over a broad area. Some new species will be planted yearly. Although this study was started in 1989, it includes some species from past studies. Presently there are 29 different species included. Twenty-two of the 29 are exhibiting 100 percent survival. Five species have failed to survive. For more information regarding plant performances refer to Table #2.

The trees and shrubs in this study are often utilized during plant identification courses held at the Center.

Table #1 reflects the species included in this assembly, accession numbers, sources and date planted. Table #2 reflects the plants' performance for years 1990-1992, 1998-2002.

Study 29A116W

Table #1 List of species included in study.

<b>Common Name</b>	<b>Genus</b>	<b>Species</b>	<b>Accession Number</b>	<b>Alternate No.</b>	<b>Source</b>	<b>Date Planted</b>
Densehead Mountain ash	<i>Sorbus</i>	<i>alnifolia</i>		7761	F.K. Nursery	11/65
Ruby redosier dogwood	<i>Cornus</i>	<i>stolonifera</i>	443229		Big Flats PMC	5/89
Late lilac	<i>Syringa</i>	<i>villosa</i>	9006228		Bismarck PMC	5/89
Redstone cornelian cherry dogwood	<i>Cornus</i>	<i>mas</i>	9055585		Elsberry PMC	5/89
Roselow sargent crabapple	<i>Malus</i>	<i>sargentii</i>	477986		Roselake PMC	5/89
Elsmo lacebark elm	<i>Ulmus</i>	<i>parvifolia</i>	9004438		Asia	5/89
Blueleaf honeysuckle	<i>Lonicera</i>	<i>korolkowi</i>	9062152		Nebraska	5/89
Birch	<i>Betula</i>	<i>species</i>	502295		Ames, IA	4/90
Willow oak	<i>Quercus</i>	<i>phellos</i>		4723	Ames, IA	4/90
Fragrant epaulettetree	<i>Pterostyrax</i>	<i>hispidia</i>		A80779	Ames, IA	4/90
Bradford pear	<i>pyrus</i>	<i>calleryana</i>		19173	Ames, IA	4/69
Prairie rose	<i>Rosa</i>	<i>setigera</i>	495616		Ames, IA	4/90
Ural falsepirea	<i>Sorbaria</i>	<i>sorbifolia</i>		7778	Ames, IA	4/90
Weeping Lilac	<i>Syringa</i>	<i>pekinensis</i>	478008		Ames, IA	4/90
Flameleaf sumac	<i>Rhus</i>	<i>copallina</i>		7764	Ames, IA	4/90
Western paper birch	<i>Betula</i>	<i>occidentalis</i>	495882		Ames, IA	4/90
Amur honeysuckle	<i>Lonicera</i>	<i>mackii</i>	477998		Ames, IA	4/90
Mountain ash	<i>Sorbus</i>	<i>reducta</i>		A-8371	Ames, IA	4/90
Blackhaw	<i>Viburnum</i>	<i>prunifolium</i>		2813	Ames, IA	4/90
Largeleaf dogwood	<i>Cornus</i>	<i>macrophylla</i>		10178	Ames, IA	4/90

<b>Common Name</b>	<b>Genus</b>	<b>Species</b>	<b>Accession Number</b>	<b>Alternate No.</b>	<b>Source</b>	<b>Date Planted</b>
Border privet	<i>Ligustrum</i>	<i>obtusifolium</i>	477010		Ames, IA	4/90
Willow oak	<i>Quercus</i>	<i>phellos</i>		4724	Ames, IA	4/90
Arrowwood	<i>Viburnum</i>	<i>dentatum</i>			Elsberry, MO	4/90
Redbud	<i>Cercis</i>	<i>canadensis</i>	496399		Ames, IA	5/91
Birch	<i>Betula</i>	<i>species</i>	14942		Ames, IA	5/91
Wichita osageorange	<i>Maclura</i>	<i>pomifera</i>			Kansas	5/91
Denmark osageorange	<i>Maclura</i>	<i>pomifera</i>			Denmark, IA	6/92
Magenta	<i>Malus</i>	<i>species</i>	514275		Roselake PMC	4/93
Ocean view beach plum	<i>Prunus</i>	<i>maritima</i>	518824		Cape May PMC	5/93
Sandy rugosa rose	<i>Rosa</i>	<i>rugosa</i>			Cape May PMC	5/93
Wildwood bayberry	<i>Myrica</i>	<i>pennsylvanica</i>	548966		Cape May PMC	5/93
Wildwood bayberry	<i>Myrica</i>	<i>pennsylvanica</i>	434150		Cape May PMC	5/93
Wildwood bayberry	<i>Myrica</i>	<i>pennsylvanica</i>	548964		Cape May PMC	5/93
Ocean view beach plum	<i>Prunus</i>	<i>maritima</i>	518822		Cape May PMC	5/93
Ocean view beach plum	<i>Prunus</i>	<i>maritima</i>	518823		Cape May PMC	5/93
Oahe hackberry	<i>Celtis</i>	<i>occidentalis</i>	476982		Bismarck PMC	5/93
King Red Russian olive	<i>Elaeagnus</i>	<i>angustifolia</i>	434029		NPMC	5/93



**Study: 29I124G**

**Study Title:** Production of Native Iowa Ecotypes of Grasses and Forbs for Roadside, Critical Areas, and All Other Vegetative Plantings Where Native Grasses and Forbs are Now Being Planted.

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

Well-adapted native grass, legume, and forb plantings offer many advantages as low cost sustainable vegetative cover for management of soil and water resources. Native plant communities resist noxious weed invasion, provide excellent erosion control, and generally require relatively low maintenance.

These characteristics make them an excellent selection for use in roadside plantings, critical areas, long term land retirement programs, and all other vegetative plantings where mono-cultures of native grasses are being planted. This is especially true along public transportation right-of-ways. These transportation corridors constitute a major land resource and management problem in the state of Iowa. Based on 1987 NRI data, over one million acres of Iowa land are devoted to rural transportation.

Proper vegetation management along these corridors is an important element in controlling soil loss and unwanted weedy plant species. Many of these acres are now seeded to introduced cool-season grass and legume species which are often invaded by noxious weeds requiring extensive mowing or herbicide treatment programs. These management techniques are expensive and can also result in additional water quality problems where herbicides are used extensively.

Managing or re-seeding these acres to promote native grasses, legumes, and forbs offers a low cost environmentally sound approach to roadside vegetation management. Herbicide use, soil erosion, and most mowing can be reduced significantly where a vigorous native grass, legume, and forb mixture dominates a roadside right-of-way. In addition, these goals are consistent with on-going NRCS programs designed to improve ground and surface water quality, reduce soil loss and increase wildlife habitat.

**Problem:**

Many adapted native species are either currently not commercially available or available only in very limited quantities. When native species are available, the origin is often from considerable distance away and adaptation can be a concern. The species that are available are often as a 'Variety' that has been developed for pasture and hay. These are generally high forage producing and more vigorous than wild collections of seed that have not been through an evaluation and breeding program. Seed of local origin that have not been improved or selected for superior forage yield is more likely to remain in a prairie mixture without crowding out other species and become a monoculture. There is a need for additional native grass, legume, and forb species for use in roadside and other types of conservation plantings.

**Objective:**

The objective of this study is to accelerate the collection and increase of selected native grass, legume, and forb species through a cooperative program between the University of Northern Iowa (UNI), USDA Natural Resources Conservation Service (NRCS), and the Iowa Roadside Integrated Vegetation Management Program (IRVM).

**Cooperators:**

The USDA Natural Resources Conservation Service, Plant Materials Center; the University of Northern Iowa; and the Integrated Roadside Vegetation Management Office

**Procedures:**

The state of Iowa was divided into three zones: North, Central, and South (Table #1). Seed collected from within each zone was kept separate from the other zones. The IRVM office organized seed collections from each zone. Collections were made from native prairie remnants throughout each zone striving for a relatively equal and representative collection. Seed from each collection site was inventoried by location and a small portion was started in the greenhouse at UNI and transplanted into plots. The remainder of the seed was sent to the PMC, cleaned, and seeded for increase plots. Seed from the plots at UNI was hand harvested and also used to start increase plots or mixed with additional seed and became available to seed growers. When enough seed becomes available, the species is released as 'Source Identified' Germplasm from the zone in which it was collected. Source identified seed has not been improved by evaluation and selection or plant breeding procedures.

**Discussion:**

The study officially started October 1, 1990, at the beginning of fiscal year 1991 with agreements signed. Seed collections had started earlier in the year and seed was available for increase plots the spring of 1991. Most of the plots started from 1991 to 1993 were destroyed in the flood the summer of 1993. Plant re-establishment started in 1994 and new plots have been started each year. Progress of species released to growers as 'Source Identified' Germplasm can be seen in Table #2.

**2000**

New increase plots established in 2000 were *Liatris asper*, rough blazing star; *Monarda fistulosa*, horsemint; and *Lobelia siphilitica*, great blue lobelia. Surflan was used for weed control and the horsemint was not resistant.

New plant releases for 2000 were Northern Iowa Germplasm Big Bluestem, Northern Iowa Germplasm Tall Dropseed, Northern Iowa Germplasm Roundhead Lespedeza, and Southern Iowa Germplasm Prairie Blazing Star.

## 2001

There were no new plant releases through the plant materials program in 2001 but seed of previous releases was allocated to growers. Initial seed increase is now in production at the new UNI Native Roadside Vegetation Center at the University of Northern Iowa, Cedar Falls, Iowa. A new plot of southern Iowa June grass was established at the PMC from plants started in the greenhouse. This species exhibits very slow growth and a serious problem is weed control.


## 2002

There were no new increase plots established in 2002. Seed production and allocation to growers continued on previously established plots.

New plant releases for 2002 were Northern, Central and Southern Iowa Germplasm New England Aster, Northern and Southern Iowa Germplasm Pale Purple Coneflower, Southern and Central Iowa Germplasm Rigid Goldenrod, and Southern Iowa Germplasm Tall Dropseed. For a complete list of Iowa ecotype releases and all other releases through the PMC program, see the section, "Releases from the Elsberry PMC", page 175 of this publication.

### IOWA ECOTYPE ZONE MAP

**Table #1**



<b>Study 29I124G-Production of Native Iowa Ecotypes of Grasses and Forbs for Roadside, Critical Areas, and All Other Vegetative Plantings Where Native Grasses and Forbs are Now Being Planted (UNI).</b>				
				<b>Table #2</b>
<b>Common Name</b>		<b>Accession</b>		
<b>Genus/Species</b>	<b>Zone</b>	<b>Number</b>	<b>Status of Accession</b>	<b>Status of Increase Plot</b>
Big bluestem	1	9068614	Released in 2000	in production
<i>Andropogon gerardii</i>	2	9068615	Released in 1998	in production
	3	9068616	Released in 1999	in production
Sideoats grama	1	9062278	Released in 1994	in production
<i>Bouteloua curtipendula</i>	2	9062279	Released in 1994	in production
	3	9062280	Released in 1994	in production
Purple prairie clover	1	9068608	Released in 1998	Increase plot planted in 1998
<i>Dalea purpurea</i>	2	9068609	Planned release in 2002	Increase plot planted in 1999
	3	9068610	Planned release in 2002	increase plot planted in 1999
Pale purple coneflower	1	9068611	Released in 2002	in production
<i>Echinacea pallida</i>	2	9068612	Released in 1998	in production
	3	9068613	Released in 2002	in production
Canada wildrye	1	9062275	Released in 1994	in production
<i>Elymus canadensis</i>	2	9062276	Released in 1994	out of production
	3	9062277	Released in 1994	out of production
Rattlesnake master	1	9068602	Released in 1998	in production
<i>Eryngium yuccifolium</i>	2	9068603	Released in 1999	in production
	3	9068604	Released in 1999	in production
Oxeye false sunflower	1	9068605	Released in 1997	in production
<i>Heliopsis scaberrima</i>	2	9068606	Released in 1996	in production
	3	9068607	Released in 1997	in production
Junegrass	1	9068620	Planned release in 2003	
<i>Loeloria macrantha</i>	2	9068621	Planned release in 2003	
	3	9068622	Planned release in 2003	increase plot planted in 2001
Round-head bushclover	1	9062281	Released in 1999	in production
<i>Lespedeza capitata</i>	2	9062282	Released in 1996	in production
	3	9062283	Released in 1997	in production
Rough blazing star	1	9068684	Planned release in 2004	increase plot planted in 2000
<i>Liatris asper</i>	2	9068685	Planned release in 2003	increase plot planted in 2000
	3	9068686	Planned release in 2003	increase plot planted in 2000
Blazing star	1	9068626	Released in 1999	in production
<i>Liatris pycnostachya</i>	2	9068627	Released in 1999	in production
	3	9068628	Released in 2000	in production

Study 29I124G - Native Iowa Ecotypes				Table #2 - continued
Common Name		Accession		
Genus/Species	Zone	Number	Status of Accession	Status of Increase Plot
Horsemint	1	9068678		increase plots planted in 2000
<i>Monarda fistulosa</i>	2	9068679		increase plots planted in 2000
	3	9068680		increase plots planted in 2000
Little bluestem	1	9062319	Released in 1999	in production
<i>Schizachyrium</i>	2	9062320	Released in 1997	in production
<i>scoparium</i>	3	9062321	Released in 1999	in production
Compassplant	1	9068675		
<i>Silphium laciniatum</i>	2	9068676		
	3	9068677		
Stiff goldenrod	1	9068617	Released in 1998	in production
<i>Solidago rigida</i>	2	9068618	Released in 2002	in production
	3	9068619	Released in 2002	in production
Indiangrass	1	9062316	Released in 1997	in production
<i>Sorghastrum nutans</i>	2	9062317	Released in 1996	in production
	3	9062318	Released in 1998	in production
Tall dropseed	1	9062313	Released in 2000	in production
<i>Sporobolus compositus</i>	2	9062314	Released in 1996	in production
	3	9062315	Released in 2002	in production
New England aster	1	9068681	Released in 2002	increase plot planted in 1999
<i>Aster novae angliae</i>	2	9068682	Released in 2002	increase plot planted in 1999
	3	9068683	Released in 2002	increase plot planted in 1999
Butterfly milkweed	1	9068687		
<i>Asclepias tuberosa</i>	2	9068688		
	3	9068689		
Blue lobelia	1	9068696	Planned release in 2003	increase plot planted in 2000
<i>Lobelia siphilitica</i>	2	9068697	Planned release in 2003	increase plot planted in 2000
	3	9068698	Planned release in 2003	increase plot planted in 2000
Switchgrass	1	9068705		
<i>Panicum virgatum</i>	2	9068706		
	3	9068707		
Golden alexanders	1	9068702		
<i>Zizia aurea</i>	2	9068703		
	3	9068703		

**Study: 29A128J**

**Study Title:** *Cornus florida* L., Flowering Dogwood Interagency Study Between Department of Interior, National Parks Service, National Capital Region (NRC) and the Department of Agriculture.

**Study Leader:** Henry, J.

**Introduction:**

Flowering dogwood is probably Missouri's favorite spring flowering tree. It is Missouri's state tree. It is a rather small tree, rarely over 30 feet high and over six to eight inches in diameter; however, in 1867 a dogwood six feet in circumference was reported in Pemiscot County, Missouri. It is commonly an understory tree to many species of oak and hickory in the hardwood forests. Besides being of great value for ornamental purposes, flowering dogwood has special wood characteristic that makes it irreplaceable for certain products. Because of its high resistance to shocks, the wood is being used almost exclusively for weaving shuttles and spool and bobbin heads. It is also being used in golf club and mallet heads and in jeweler's blocks.

**Objectives:**

- A. Clean (depulp) and condition seed collections and keep accession records on individual ecotypes.
- B. Establish at Elsberry PMC, an area free of dogwood anthranose, 12 to 15 plants from three specified parks for a period of 30 to 40 years.
- C. Provide, upon request, a report on the status of the plants maintained by NRCS.
- D. Provide a study coordinator for all activities performed by NRCS under the terms of the Interagency Agreement.
- E. Provide seed to the NRCS upon request.

**Discussion:**

**1994 - 1999**

As of the date this report was written there has only been one accession of flowering dogwood received at the PMC. This accession was planted in Field #11 May 1993. Five of the ten plants are surviving in good vigor. Height ranges from 4.0 feet to 4.5 feet; spread ranges from three to three and a half feet. Vigor is excellent along with its resistance to insects and diseases. There have been no indications of the anthranose disease affecting these plants.

## 2000

The five remaining plants of accession 9083225 are surviving in good vigor. Height ranges from 4.8 to 5.0 feet and spread ranges from 3.9 feet to 4.1 feet. There have been no signs of insects or diseases associated with this accession.

## 2001

Only three plants of a total of ten plants are surviving in good vigor. The reason for the decline in the number of plants surviving was due to severe mechanical damage resulting in death of the plants. Height ranges from 5.0 feet to 5.6 feet and spread ranges from 4.1 feet to 4.7 feet. No apparent signs of insects or diseases have been associated with this accession (9083225).

## 2002

Two plants are surviving out of a total of ten initially planted. Mechanical damage has caused the decline in the number of surviving plants. Evaluations of the surviving plants were made on June 10 and October 22. The height of these plants (accession 9083225) is 6.3 feet with a spread of 5.0 feet. There were no apparent signs of insect, disease, or fruit production.

### **Study: 29I1320**

**Study Title:** Miscellaneous Wetland Plant Evaluation

**Study Leader:** Henry, J.

### **Introduction:**

Wetlands are areas, periodically saturated or inundated by surface or ground water, that support vegetation adapted for saturated soil conditions. Wetlands in the Environmental Protection Agency (EPA) Region Seven states of Iowa, Kansas, Missouri and Nebraska are generally found along rivers and streams and their associated floodplains or at the margins of lakes and ponds. Wetlands can also occur in upland depressions, such as the prairie “potholes” of Iowa, or in seepage areas along slopes. Because of their location between land and water, wetlands function to improve water quality. They control erosion and trap the runoff from land carrying nutrients, waste, pollution, and sediment and filter the material from flooding waters. Thus ponds, lakes, rivers, streams and our drinking water remain clear and healthy.

Wetland ecosystems support a great diversity of vegetation, which provides food, water, cover, nesting, and wintering ground for many forms of wildlife that use them for all or parts of their life cycles. In fact, wetlands are some of the most biologically unique and productive areas on earth.

**Problem:**

Naturally occurring wetlands and constructed wetlands, for water quality improvement and wildlife habitat enhancement, require plants that respond to different water regimes and pollutant loads. Facets of these plants' establishment, management and benefits must be explored. This information can then be used and recommended.

**Objective:**

Identify, establish, and evaluate for possible increase selected plant materials needed for wetland enhancement, restoration, and creation to meet resource conservation and related water quality program requirements.

**Discussion:****1992-1999**

Initially, seven wetland cells, 16 feet long by four feet wide and 18 inches deep were constructed using landscape ties, tarp and a double layer of plastic (8 mil). Eighteen inches of good topsoil was placed in each cell. Water was then added to saturate the soil before the planting operation. The following plant species were assembled at the PMC and transplanted in the cells during

July 1992: *Scirpus validus*, softstem bulrush; *Sagittaria latifolia*, smooth-cone sedge; *Typha latifolia*, cattails; *Asclepias incarnata*, swamp milkweed and *Ludwigia peploides*, water primrose.

Each species was watered according to its need identified in a literature search. It became evident that each species required different quantities of water. When water was not provided to the smooth cone sedge in the suggested amount, the stand began to deteriorate. The other species reacted less dramatic than the smooth cone sedge to the reduction in water.

Plans are to release the *Carex laericonica*, smooth cone sedge in year 2003.

Table #1 contains information regarding sources for the different collections included in this study.

Table #2 reflects the plants' performance from 1992 – 1999.

**2000**

Evaluations were conducted during year 2000 along with the seed harvest of *Carex laericonica*, smooth cone sedge. Plant performance data can be found in Table #2 for years 1992 – 2000.

**2001**

This study was again evaluated for percent survival, flower date, seed production, spread, vigor, and insect and disease resistance. Seed was not harvested from the *Carex laericonica*, smooth

cone sedge planting this year because the stand is getting thin and thoughts were that this year's seed production might thicken up the stand. Plant performance data can be found in Table #2 for years 1999 – 2001.

**2002**

There was little to no seed production recorded in 2002 for the *Carex laericonica*, smooth cone sedge. Smooth cone sedge was the selection made out of this study. Seed harvests have been attempted in order to accumulate enough seed to make an increase planting. The smooth cone sedge will also be included in the Filter Strip Study (MOPMC-BU-0106). The stand is still quite weak and has a reed canarygrass infestation. This plant's performance along with other species included in this study can be found in Table #2. Table #1 reflects accession information.

**Study 29I1320 - Miscellaneous Wetland Plant Evaluation**

**Table #1**

<b>Genus/species</b>	<b>Accession Number</b>	<b>Source</b>	<b>City/State</b>
<i>Scirpus validus</i> Softstem bulrush	9083201	Kester's Nurseries, Inc.	Omro, Wisconsin
<i>Sagittaria latifolia</i> Arrowhead	9083202	Kester's Nurseries, Inc.	Omro, Wisconsin
<i>Juncus offusus</i> Soft rush	9083203	Kester's Nurseries, Inc.	Omro, Wisconsin
<i>Carex laericonica</i> Smoothcone sedge	9083204	Field #7, PMC	Elsberry, Missouri
<i>Typha latifolia</i> Cattail	9083205	County Route #79	Elsberry, Missouri
<i>Ludwigia peplaides</i> Water primrose	9083206	BK Leach Wildlife Area	Elsberry, Missouri
<i>Ascepias incarnata</i> Swamp milkweed	9083207	BK Leach Wildlife Area	Elsberry, Missouri

Study 29I1320 Miscellaneous Wetland Plant Evaluation									Table #2
	Year	Percent	Flower	Seed	End of			Insect	Disease
Genus/Species	Eval.	Survival	Date	Prod.	Season Ht	Spread	Vigor	Resist.	Resist
				√1			√1	√1	√1
<b>Scirpus validus</b>	1992	100	5/19/92	5	50 inches	solid	1	1	1
softstem bulrush	1993	100	5/21/93	5	53 inches	solid	1	1	1
9083201	1994	100	5/17/94	3	55 inches	solid	1	1	1
	1995	100	5/24/95	3	55 inches	solid	1	1	1
	1996	100	5/20/96	2	55 inches	solid	1	1	1
	1997	95	5/23/97	3	55 inches	solid	1	1	1
	1998	90	5/18/98	5	55 inches	solid	1	1	1
	1999	85	5/10/99	5	50 inches	solid	1	1	1
	2000	75	5/17/00	4	50 inches	solid	2	1	1
	2001	60	5/21/01	6	47 inches	solid	4	1	1
	2002	50	5/17/02	7	45 inches	solid	5	1	1
<b>Sagittaria latifolia</b>	1992	100	5/27/92	6	65 inches	solid	1	1	1
arrowhead	1993	100	5/25/93	6	68 inches	solid	1	1	1
9083202	1994	100	5/23/94	6	75 inches	solid	1	1	1
	1995	100	5/24/95	6	75 inches	solid	1	1	1
	1996	95	5/27/96	6	75 inches	solid	1	1	1
	1997	95	5/23/97	6	75 inches	solid	1	1	1
	1998	90	5/26/98	6	75 inches	solid	1	1	1
	1999	90	5/21/99	7	72 inches	solid	1	1	1
	2000	85	5/23/00	6	70 inches	solid	2	2	2
	2001	80	5/28/01	6	65 inches	solid	3	2	2
	2002	75	5/24/02	7	60 inches	solid	3	2	2
<b>Juncus offusus</b>	1992	100	5/19/92	5	38 inches	solid	1	1	1
soft rush	1993	100	5/25/93	5	45 inches	solid	1	1	1
9083203	1994	100	5/23/94	5	52 inches	solid	1	1	1
	1995	100	5/26/95	5	52 inches	solid	1	1	1
	1996	95	5/21/96	5	52 inches	solid	1	1	1
	1997	95	5/23/97	5	50 inches	solid	1	1	1
	1998	90	5/26/98	5	50 inches	solid	1	1	1
	1999	90	5/21/99	6	48 inches	solid	1	1	1
	2000	90	5/23/00	7	45 inches	solid	2	2	2
	2001	20	5/28/01	8	40 inches	solid	3	2	2
	2002	15	5/24/02	8	37 inches	solid	3	2	2
<b>Carex laericonica</b>	1992	100	6/3/92	6	24 inches	solid	4	1	1
smooth cone sedge	1993	100	6/6/93	5	30 inches	solid	3	1	1
9083204	1994	90	6/1/94	5	32 inches		3	1	1
	1995	85	5/31/95	6	32 inches		2	1	1
	1996	70	6/4/96	7	32 inches		2	1	1
	1997	60	6/6/97	7	32 inches		2	1	1
	1998	50	6/8/98	7	32 inches		2	1	1
	1999	50	6/4/99	7	30 inches		3	1	1
	2000	50	6/9/00	5	32 inches		3	1	1
	2001	45	6/11/01	5	30 inches		5	1	1
	2002	40	6/7/02	5	30 inches	spotty	5	1	1

Study 29I1320 Miscellaneous Wetland Plant Evaluation							Table #2 - continued		
	Year	Percent	Flower	Seed	End of			Insect	Disease
Genus/Species	Eval.	Survival	Date	Prod.	Season Ht	Spread	Vigor	Resist.	Resist
				√1			√1	√1	√1
<b><i>Typha latifolia</i></b>	1992	100	5/5/92	2	60 inches	solid	1	1	1
cattail	1993	100	5/7/93	2	80 inches	solid	1	1	1
9083205	1994	100	5/3/94	2	80 inches	solid	1	1	1
	1995	100	5/1/95	2	80 inches	solid	1	1	1
	1996	100	5/8/96	2	80 inches	solid	1	1	1
	1997	100	5/2/97	2	75 inches	solid	1	1	1
	1998	100	5/4/98	2	70 inches	solid	1	1	1
	1999	100	5/7/99	1	68 inches	solid	1	1	1
	2000	100	5/10/00	2	65 inches	solid	1	1	1
	2001	50	5/14/01	2	60 inches	solid	1	1	1
	2002	45	5/10/02	2	60 inches	solid	1	1	1
<b><i>Ludwigia peploides</i></b>	1992	90	6/21/92	0	3 inches		3	3	3
water primrose	1993	80	6/24/93	0	6 inches		3	2	2
9083206	1994	70	6/21/94	0	6 inches		3	2	2
	1995	70	6/27/95	0	6 inches		3	2	2
	1996	60	6/24/96	0	6 inches		3	2	2
	1997	60	6/30/97	0	6 inches		3	2	2
	1998	60	6/26/98	0	6 inches		3	2	2
	2000	40	6/29/00	0	4 inches		4	3	3
	2001	30	6/23/01	0	4 inches		4	3	3
	2002	25	6/24/02	0	4 inches		4	3	3
<b><i>Asclepias incarnata</i></b>	1992	died 1992							
swamp milkweed									
9083207									
Rating: Vigor, Insect & Disease Resist: 1 = Excellent, 9 = Poor									
Rating: Seed Production: 1 = Excellent, 9 = Poor & 0 = No Seed Produced									

**Study No. 29I134J**

**Study Title:** Assembly and Evaluation of Eastern Redcedar, *Juniper virginiana L.*

**Study Leader:** Henry, J.

**Introduction:**

Eastern redcedar has the most uniform distribution of the four species of conifers native to Missouri. Although it is most common in the Ozark region, it is found throughout the state. Scale-like or awl-shaped leaves are opposite or ternate around a minute four-angled dark green central stem. The flowers are male and female on separate trees with the male flowers being cone-like, with four to six scales. The female flower structure has fleshy scales. Fruits are bluish

in color and about the size of a pea with a white frost-like bloom and contain one to four seeds. The flesh is sweet and resinous and twigs are slender, four-angled and become reddish-brown with inconspicuous buds. Its bark ranges in color from a tan to reddish-brown and shreddy.

Eastern redcedar flowers during March-May with fruit ripening during September-November.

**Problem:**

There is a lack of an available cultivar of eastern redcedar specifically for this area. NRCS and other conservation and wildlife agencies have identified a need for developing a selection and also source identified sources of redcedar for use as a native juniper for windbreaks and secondary benefits for wildlife habitat in the three states being served by the center.

**Objective:**

The objective is to assemble, comparatively evaluate, select and release a selected, tested and/or cultivar of redcedar for the PMC service area. The selection criteria are for a columnar, upright selection with minimal production of seed.

**Discussion:**

**1989 - 1992**

Collections were received from Illinois and Missouri between 1989 and 1991. Forty-six collections were made (16 from Illinois and 30 from Missouri) and the seed was stratified the fall of 1992.

**1993 - 1998**

Thirty-four of the total 46 collections germinated and were grown out in the PMC greenhouse to a height ranging from 1.5 to 3.0 feet. The planting of the redcedar assembly was made in Field #7 on the PMC on May 17 and 18, 1994. The plot design was a randomized complete block with six replications.

Table #1 reflects the different accessions, states, county or city where these collections were made; Tables #2, #3, #4, #5, and #6 reflect the plants' performance.

**1999**

Evaluations were made on November 22, 1999 for the following: height, spread, vigor, insect and disease resistance and form; this information was not added to Tables #2, #3, #4, #5, and #6.

## 2000

The evaluations documented on November 22, 1999 along with the evaluations made on October 10, 2000 were added to Tables #2, #3, #4, #5 and #6 reflecting plants' performance for years 1997, 1998 and 2000.

## 2001

The only evaluations made in 2001 regarding this study were insect and disease resistance. A severe infestation of bagworms completely defoliated more than half of the entire planting (southern half). All accessions were documented as having bagworms, the southern half of the planting was completely defoliated. The northern half was not defoliated by the bagworms but there was a heavy infestation present. No pesticides were applied, as this was one of the criteria being evaluated. Evaluations on height, spread and form are scheduled for every 3-5 years.

## 2002

The only evaluations conducted on this study were for bagworm infestation as the height, spread, form and fruit production are conducted every 3–5 years. The bag worm evaluations revealed none present in the entire assembly.

### Accessions of Eastern Redcedar Collected for this Study.

**Table # 1**

ACCESSION	STATE	COUNTY OR CITY
9057099	Illinois	Tazewell
9057105	Illinois	Tazewell
9057106	Illinois	Mason
9057115	Illinois	Grundy
9057116	Illinois	Jo Daviess
9057117	Illinois	Jo Daviess
9057136	Illinois	Kendall
9057156	Illinois	Mason
9057180	Illinois	Pope
9068488	Illinois	Jo Daviess
9068579	Illinois	Jo Daviess
9057196	Illinois	Henderson
9068498	Illinois	Ogle
9068497	Illinois	Henderson
9068495	Illinois	Carroll
9068531	Illinois	Cole

<b>ACCESSION</b>	<b>STATE</b>	<b>COUNTY OR CITY</b>
		<b>Table #1 cont.</b>
9068487	Missouri	Cooper
9068486	Missouri	Pettis
9057198	Missouri	Bates
9057199	Missouri	Cooper
9058476	Missouri	Pettis
9057187	Missouri	Johnson
9057190	Missouri	St. Clair
9057189	Missouri	Morgan
9068504	Missouri	Hickory
9068503	Missouri	Mercer
9068502	Missouri	Cooper
9068501	Missouri	St. Clair
9068500	Missouri	Mercer
9068499	Missouri	Camden
9068496	Missouri	Mercer
9068495	Missouri	Carroll
9068494	Missouri	Livingston
9068493	Missouri	Mercer
9068492	Missouri	Cooper
9068532	Missouri	Miller
9068530	Missouri	Vernon
9068554	Missouri	Phelps
9068551	Missouri	Lafayette
9068566	Missouri	Plattsburg/Clinton
9068569	Missouri	Lincoln
9068564	Missouri	Cole
9068582	Missouri	Warren
9068584	Missouri	Moniteau
9068583	Missouri	Dent
9068588	Missouri	Clinton
9068486	Missouri	Pettis
9057198	Missouri	Bates
9057199	Missouri	Cooper
9058476	Missouri	Pettis
9057187	Missouri	Johnson
9057190	Missouri	St. Clair
9057189	Missouri	Morgan

<b>ACCESSION</b>	<b>STATE</b>	<b>COUNTY OR CITY</b>
		<b>Table #1 cont.</b>
9068504	Missouri	Hickory
9068503	Missouri	Mercer
9068502	Missouri	Cooper
9068501	Missouri	St. Clair
9068500	Missouri	Mercer
9068499	Missouri	Camden
9068496	Missouri	Mercer
9068495	Missouri	Carroll
9068494	Missouri	Livingston
9068493	Missouri	Mercer
9068492	Missouri	Cooper
9068532	Missouri	Miller
9068530	Missouri	Vernon
9068554	Missouri	Phelps
9068551	Missouri	Lafayette
9068566	Missouri	Plattsburg/Clinton
9068569	Missouri	Lincoln
9068564	Missouri	Cole
9068582	Missouri	Warren
9068584	Missouri	Moniteau
9068583	Missouri	Dent
9068588	Missouri	Clinton

Study 291134J - Assembly and Evaluation of Eastern Redcedar, <i>Juniper virginia</i> L.																				
											Height in Feet									
1997											1998									
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Ave.	Best	Location		Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Ave.	Best	Location
9068493	7.60	8.00	6.60	8.00	10.00	5.80	7.67	10.20	R5		9068493	8.00	8.20	7.00	8.40	10.60	6.20	8.07	10.60	R5
9068486	9.70	8.00	9.00	7.00	7.00	9.30	8.33	9.70	R1		9068532	10.60	6.30	8.40	6.60	8.60	8.60	8.18	10.60	R1
9068566	8.60	9.60	7.90	8.00	6.20	6.60	7.82	9.60	R2		9057106	9.00	6.60	10.00	6.60	7.70	8.60	8.08	10.00	R3
9057106	8.60	6.30	9.40	6.00	7.00	8.10	7.57	9.40	R3		9068486	10.00	8.20	9.50	7.80	7.20	10.00	8.78	10.00	R1,6
9057196	9.40	6.00	5.00	7.50	7.20	8.00	7.18	9.40	R1		9057196	10.00	7.50	5.60	8.00	7.60	8.50	7.87	10.00	R1
9057198	8.00	9.30	7.50	6.00	7.00	7.20	7.50	9.30	R2		9057198	8.50	10.00	8.60	7.20	7.30	7.50	8.18	10.00	R2
9057199	9.30	8.70	7.00	7.40	7.00	7.00	6.20	9.30	R1		9068530	10.00	8.70	8.00	8.60	9.00	8.60	8.82	10.00	R1
9068530	9.20	8.20	8.20	8.20	8.00	8.10	8.32	9.20	R1		9068566	9.00	10.00	9.20	8.70	6.60	7.00	8.42	10.00	R2
9068500	9.00	8.80	9.10	8.00	5.10	4.20	7.37	9.10	R3		9068583	5.50	9.00	8.80	8.00	7.80	10.00	8.18	10.00	R6
9068499	8.60	9.10	4.60	5.60	6.50	6.80	6.87	9.10	R2		9057136	9.90	8.20	8.50	8.00	6.20	3.00	7.30	9.90	R1
9057105	8.30	9.00	6.80	8.30	7.40	8.50	8.05	9.00	R2		9057099	9.00	9.60	6.70	7.00	8.00	7.20	7.92	9.60	R2
9057136	9.00	7.60	8.00	7.60	5.40	2.50	6.68	9.00	R1		9068499	9.00	9.60	6.50	5.00	8.00	5.20	7.22	9.60	R2
9068531	9.00	6.80	6.60	8.20	7.10	7.40	7.52	9.00	R1		9068496	8.00	9.60	8.20	6.50	5.70	4.50	7.08	9.60	R2
9057190	8.90	8.50	6.90	7.80	8.20	8.60	8.15	8.90	R1,6		9057105	8.60	9.50	7.00	8.50	9.00	9.00	8.60	9.50	R2
9068532	8.90	5.90	7.30	6.00	8.40	8.10	7.43	8.90	R1		9068500	9.50	9.00	9.20	8.20	5.80	4.60	7.72	9.50	R1
9068496	7.30	8.80	6.70	6.30	5.20	4.10	6.40	8.80	R2		9057190	9.40	8.00	7.20	8.00	Dead	8.80	8.28	9.40	R1
9068501	8.70	8.20	8.40	7.60	6.10	7.00	7.67	8.70	R1		9068504	8.00	8.50	7.70	9.40	6.80	8.00	8.07	9.40	R4
9068495	6.80	7.40	8.70	5.20	7.00	6.00	6.85	8.70	R3		9068531	9.40	7.10	7.00	7.80	6.70	6.80	7.47	9.40	R1
9057099	8.00	8.60	6.70	6.40	6.80	6.30	7.13	8.60	R2		9057117	8.00	9.00	6.50	7.30	8.00	7.00	7.63	9.00	R2
9057189	7.80	7.80	7.00	8.60	7.10	8.20	7.75	8.60	R4		9057193	8.50	9.00	8.00	8.00	9.00	7.60	8.35	9.00	R2,5
9068583	5.00	8.60	8.00	7.20	7.00	7.80	7.27	8.60	R2		9057199	9.00	8.00	7.30	6.40	7.60	6.40	7.45	9.00	R1
9068588	8.60	8.50	5.90	5.70	6.70	7.80	7.20	8.60	R1		9068503	8.80	9.00	7.80	6.60	7.50	8.00	7.95	9.00	R2
9057117	7.20	8.50	6.00	7.00	7.90	6.20	7.13	8.50	R2		9068501	9.00	9.00	8.60	8.00	6.50	7.80	8.15	9.00	R1,2
9057193	8.00	8.50	7.40	7.40	8.10	7.80	7.87	8.50	R2		9068588	9.00	9.00	6.20	6.00	7.00	8.20	7.57	9.00	R1,2
9068503	8.30	8.50	7.50	7.60	6.80	6.90	7.60	8.50	R2		9057180	8.00	8.60	8.80	8.00	8.80	8.20	8.40	8.80	R3,5
9068504	7.80	8.20	7.20	6.30	6.20	7.30	7.17	8.20	R2		9057116	8.60	6.30	7.00	6.00	7.70	8.20	7.30	8.60	R1
9068492	8.20	8.20	5.30	7.20	8.10	7.10	7.35	8.20	R1,2		9057189	8.00	8.40	8.00	8.60	8.00	7.50	8.08	8.60	R2
9068502	7.70	8.10	6.20	5.50	6.20	5.80	6.58	8.10	R2		9068502	8.00	8.60	6.50	5.50	6.50	7.60	7.12	8.60	R2
9068554	7.80	7.00	8.10	7.70	8.00	7.00	7.60	8.10	R3		9068492	8.60	8.60	6.20	8.40	8.00	7.20	7.83	8.60	R1,2
9068584	7.00	8.10	5.60	7.30	8.00	6.80	7.13	8.10	R2		9068495	8.00	8.00	8.60	5.60	7.40	6.20	7.30	8.60	R3
9057116	8.00	6.00	6.50	5.50	6.50	8.00	6.75	8.00	R1,6		9068554	7.00	7.30	8.50	8.00	8.40	7.40	7.77	8.50	R3
9068476	6.60	7.80	6.70	7.30	7.60	8.00	7.33	8.00	R6		9068476	7.00	8.00	7.10	8.20	8.00	8.40	7.78	8.40	R6
9057180	7.60	6.90	7.80	7.50	6.70	7.20	7.28	7.80	R3		9068584	7.50	8.60	6.00	8.30	8.40	7.20	7.67	8.40	R5
9057115	6.30	4.50	4.70	7.50	4.50	5.60	5.52	7.50	R4		9057115	6.80	5.00	5.40	7.80	5.70	6.00	6.12	7.80	R4
Height measured in feet																				



Study 291134J - Assembly and Evaluation of Eastern Redcedar, <i>Juniper virginia</i> L.										Table #3									
										Spread in Feet									
1997										1998									
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Ave.	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Ave.	Best	Location
9057115	4.70	4.60	5.00	5.60	4.00	4.20	4.68	<b>5.60</b>	R4	9057115	5.10	5.00	5.80	6.10	4.70	4.80	5.25	<b>6.10</b>	R4
9068503	6.00	5.50	5.10	5.00	5.60	5.00	5.37	<b>6.00</b>	R1	9068503	6.40	6.00	5.60	5.60	6.10	5.90	5.93	<b>6.40</b>	R1
9068493	6.00	5.00	5.60	5.20	4.90	5.30	5.33	<b>6.00</b>	R1	9068493	6.60	5.40	6.00	5.80	5.30	5.90	5.83	<b>6.60</b>	R1
9068476	4.60	6.50	5.70	6.70	5.10	5.30	5.65	<b>6.50</b>	R2	9068530	6.80	6.90	6.00	6.50	6.00	5.50	6.28	<b>6.70</b>	R2
9068495	6.60	6.10	4.90	4.00	5.70	4.30	5.27	<b>6.60</b>	R1	9068495	6.80	6.70	5.20	4.40	6.20	4.90	5.70	<b>6.80</b>	R1
9068530	6.20	6.60	5.70	6.00	5.60	5.00	5.85	<b>6.60</b>	R2	9057106	7.00	6.00	7.60	5.40	6.60	5.80	6.40	<b>7.00</b>	R1
9068584	5.70	5.50	5.40	5.70	4.20	6.60	5.52	<b>6.60</b>	R6	9068476	5.00	7.00	6.30	7.00	5.70	5.90	6.15	<b>7.00</b>	R2
9068492	6.30	5.80	5.90	6.70	5.80	6.00	6.08	<b>6.70</b>	R4	9068492	7.00	6.20	6.30	6.90	6.20	6.50	6.52	<b>7.00</b>	R1
9068499	5.50	6.90	5.10	5.80	5.30	4.00	5.43	<b>6.90</b>	R2	9068554	7.00	7.00	7.00	6.90	6.90	5.30	6.68	<b>7.00</b>	R1,2,3
9068554	6.70	6.80	6.90	6.70	6.60	5.00	6.45	<b>6.90</b>	R3	9068584	7.00	6.00	4.90	5.40	5.90	5.20	5.73	<b>7.00</b>	R1
9057106	6.70	5.50	7.00	5.00	6.10	5.20	5.92	<b>7.00</b>	R3	9068588	7.90	7.30	7.40	5.90	6.60	4.60	6.62	<b>7.00</b>	R1,2
9057193	7.00	6.10	5.80	5.80	6.70	5.40	6.13	<b>7.00</b>	R1	9068499	6.00	7.20	5.60	6.20	5.90	4.80	5.95	<b>7.20</b>	R2
9057116	7.10	5.70	4.80	5.80	6.20	5.20	5.80	<b>7.10</b>	R1	9057193	7.50	6.60	6.30	6.50	6.00	5.30	6.37	<b>7.50</b>	R1
9057199	5.30	7.10	5.80	6.00	5.90	5.00	5.85	<b>7.10</b>	R2	9068504	7.50	7.50	6.80	6.40	5.60	5.70	6.58	<b>7.50</b>	R1,2
9068504	7.00	7.10	6.30	6.00	5.00	5.10	6.08	<b>7.10</b>	R2	9057116	7.60	6.00	5.30	6.00	6.80	6.00	6.28	<b>7.60</b>	R1
9068502	6.60	7.20	5.20	3.10	5.20	5.00	5.38	<b>7.20</b>	R2	9057199	6.00	7.60	6.20	6.50	6.30	5.40	6.33	<b>7.60</b>	R2
9068500	6.10	7.20	6.20	5.10	4.20	2.20	5.17	<b>7.20</b>	R2	9068500	6.50	7.60	6.60	5.90	4.90	2.80	5.72	<b>7.60</b>	R2
9068501	5.70	7.30	5.10	5.70	7.10	4.60	5.92	<b>7.30</b>	R2	9068501	6.00	7.70	5.60	6.00	7.60	5.20	6.35	<b>7.70</b>	R2
9057099	7.60	6.50	3.90	6.80	3.80	6.90	5.92	<b>7.60</b>	R1	9057180	7.00	6.00	6.20	7.80	7.00	5.90	6.65	<b>7.80</b>	R4
9057180	6.60	5.70	5.80	7.60	6.50	5.30	6.25	<b>7.60</b>	R4	9068502	7.00	7.80	5.80	3.60	5.70	5.40	5.88	<b>7.80</b>	R2
9057189	5.70	7.60	7.40	6.30	5.00	5.90	6.32	<b>7.60</b>	R2	9057099	8.00	7.00	4.30	7.10	4.10	7.30	6.30	<b>8.00</b>	R1
9068532	7.60	6.20	6.50	6.70	6.50	6.30	6.63	<b>7.60</b>	R1	9057189	6.00	7.00	8.00	7.20	5.40	6.20	6.63	<b>8.00</b>	R3
9068566	6.80	7.70	7.20	5.90	6.00	4.90	6.42	<b>7.70</b>	R2	9068566	7.40	7.90	8.00	6.30	6.40	5.20	6.87	<b>8.00</b>	R3
9068588	7.70	7.00	7.10	5.30	6.20	4.10	6.23	<b>7.70</b>	R1	9068532	8.20	6.60	6.80	7.20	6.80	6.80	7.07	<b>8.20</b>	R1
9057196	8.00	7.10	5.20	6.10	6.80	4.70	6.32	<b>8.00</b>	R1	9068496	7.50	8.40	6.80	4.40	5.30	4.40	6.13	<b>8.40</b>	R2
9068496	7.00	8.00	6.30	4.00	4.80	4.00	5.68	<b>8.00</b>	R2	9057196	8.50	7.60	6.00	6.60	7.30	5.80	6.97	<b>8.50</b>	R1
9068531	8.30	6.50	5.60	6.00	5.50	6.60	6.42	<b>8.30</b>	R1	9068583	6.00	8.60	7.20	7.00	5.20	6.00	6.67	<b>8.60</b>	R2
9057105	8.50	7.30	5.50	6.60	5.70	6.60	6.70	<b>8.50</b>	R1	9068531	8.90	7.00	6.00	6.40	6.00	7.00	6.88	<b>8.90</b>	R1
9068486	8.50	5.70	6.70	7.00	5.00	5.40	6.38	<b>8.50</b>	R1	9057105	9.00	8.00	6.70	7.00	6.00	7.00	7.28	<b>9.00</b>	R1
9057190	8.50	5.70	4.60	5.80	0.00	5.20	5.96	<b>8.50</b>	R1	9057117	6.60	9.00	5.80	6.90	5.30	5.70	6.55	<b>9.00</b>	R2
9068583	6.80	8.50	7.60	7.00	5.10	6.40	6.90	<b>8.50</b>	R2	9068486	9.00	6.20	7.00	7.50	5.70	6.00	6.90	<b>9.00</b>	R1
9057136	8.60	5.90	6.50	7.20	6.60	4.80	6.60	<b>8.60</b>	R1	9057190	9.00	6.00	5.20	6.00	0.00	5.70	6.38	<b>9.00</b>	R1
9057117	6.10	8.80	5.20	6.30	4.80	5.10	6.05	<b>8.80</b>	R2	9057136	9.30	7.00	6.50	6.60	5.40	9.00	7.30	<b>9.30</b>	R1
9057198	5.40	9.80	5.50	4.60	3.80	5.80	5.82	<b>9.80</b>	R2	9057198	6.00	10.60	5.90	5.00	4.20	6.20	6.32	<b>10.60</b>	R2
<b>Spread measured in feet</b>																			

Study 291134J - Assembly and Evaluation of Eastern Redcedar, <i>Juniper virginia</i> L.										Table #3 - continued		
										Spread in Feet		
										2000		
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Ave.	Best	Location			
9068500	7.00	8.40	7.30	6.40	5.40	3.50	6.33	3.50	R6			
9068502	7.70	8.30	6.20	4.30	6.20	6.40	6.52	4.30	R4			
9057099	8.70	7.40	5.00	7.70	4.70	8.20	6.95	4.70	R5			
9057198	6.40	11.20	6.60	5.40	4.90	6.90	6.90	4.90	R5			
9068496	8.00	9.20	7.50	5.00	6.00	5.00	6.78	5.00	R4,6			
9068499	6.40	8.00	6.10	6.80	6.40	5.30	6.50	5.30	R6			
9057115	5.90	5.70	6.20	6.90	5.40	5.60	5.95	5.40	R5			
9068476	5.40	7.60	6.80	7.60	6.30	6.60	6.72	5.40	R1			
9068495	7.50	7.20	5.80	5.00	6.90	5.40	6.30	5.40	R6			
9068584	7.50	6.40	5.50	6.00	6.40	6.50	6.38	5.50	R3			
9057190	9.70	6.50	5.70	6.40	Dead	6.20	5.75	5.70	R3			
9068583	6.30	9.00	7.80	7.50	5.70	6.60	7.15	5.70	R5			
9057189	6.60	7.50	8.50	7.90	5.80	7.20	7.25	5.80	R5			
9068501	6.40	8.40	6.20	6.50	8.30	5.80	6.93	5.80	R6			
9068554	7.50	7.50	7.50	7.30	7.40	5.80	7.17	5.80	R6			
9057116	8.40	6.60	5.90	6.80	7.40	6.70	6.97	5.90	R3			
9057117	7.30	9.70	6.30	7.50	5.90	6.40	7.18	5.90	R5			
9068493	7.30	6.00	6.40	6.20	5.90	6.30	6.35	5.90	R5			
9057106	7.70	6.50	8.00	6.00	7.20	6.30	6.95	6.00	R4			
9057136	10.00	7.80	7.00	7.10	6.00	9.60	7.92	6.00	R5			
9057193	8.00	7.20	6.80	7.00	6.00	7.10	7.02	6.00	R5			
9068504	8.20	8.00	7.20	7.10	6.00	7.30	7.30	6.00	R5			
9068566	8.20	8.60	8.80	7.00	7.10	6.00	7.62	6.00	R6			
9068503	7.10	6.50	6.10	6.10	6.80	6.50	6.52	6.10	R3,4			
9068486	9.70	6.80	7.50	8.00	6.20	6.50	7.45	6.20	R5			
9068588	8.50	8.10	8.20	6.30	7.40	7.70	7.70	6.30	R4			
9068531	9.50	7.40	6.50	7.00	6.40	7.70	7.42	6.40	R5			
9057105	9.70	8.60	7.30	7.60	6.50	7.70	7.90	6.50	R5			
9057180	7.60	6.50	6.90	8.30	7.50	7.30	7.35	6.50	R2			
9057196	9.00	8.30	6.50	7.10	8.00	7.50	7.73	6.50	R3			
9068530	7.30	7.40	6.50	7.20	6.70	6.90	7.00	6.50	R3			
9057199	6.60	8.40	6.80	7.00	6.90	6.80	7.08	6.60	R1			
9068492	7.60	6.80	6.90	7.40	7.00	7.10	7.13	6.80	R2			
9068532	9.00	7.20	7.40	8.00	7.50	7.50	7.77	7.20	R2			
Spread measured in feet												





Study 291134J - Assembly and Evaluation of Eastern Redcedar, <i>Juniper virginia</i> L.										Table #5									
										Insect/Disease									
1997										1998									
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Ave.	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Ave.	Best	Location
9068588								0.00		9068531								0.00	
9068486	1.00	2.00	1.00	1.00	2.00	2.00	1.50	1.00	R1,3,4	9057099	2.00	1.00	4.00	2.00	3.00	3.00	2.50	1.00	R2
9057198	2.00	1.00	2.00	5.00	1.00	2.00	2.17	1.00	R2,5	9057106	1.00	3.00	2.00	2.00	3.00	3.00	2.33	1.00	R1
9057199	2.00	1.00	3.00	2.00	3.00	1.00	2.00	1.00	R2,6	9057193	2.00	2.00	2.00	3.00	1.00	2.00	2.00	1.00	R5
9068504	1.00	2.00	2.00	1.00	2.00	2.00	1.67	1.00	R1,4	9057196	1.00	2.00	3.00	2.00	3.00	3.00	2.33	1.00	R1
9068503	2.00	2.00	3.00	3.00	1.00	3.00	2.33	1.00	R5	9057198	2.00	2.00	1.00	7.00	1.00	3.00	2.67	1.00	R3,5
9068502	2.00	2.00	1.00	2.00	3.00	3.00	2.17	1.00	R3	9057199	1.00	1.00	2.00	2.00	1.00	1.00	1.33	1.00	R1,2,5,6
9068501	2.00	2.00	2.00	1.00	2.00	2.00	1.83	1.00	R4	9068504	1.00	2.00	2.00	1.00	3.00	1.00	1.67	1.00	R1,4
9068500	2.00	2.00	2.00	2.00	1.00	3.00	2.00	1.00	R5	9068503	3.00	1.00	3.00	2.00	2.00	3.00	2.33	1.00	R2
9068499	1.00	2.00	3.00	2.00	6.00	3.00	2.83	1.00	R1	9068502	2.00	2.00	1.00	2.00	3.00	2.00	2.00	1.00	R3
9068496	1.00	2.00	4.00	2.00	2.00	2.00	2.17	1.00	R1	9068500	1.00	3.00	2.00	2.00	1.00	3.00	2.00	1.00	R1,5
9068495	1.00	1.00	1.00	-	2.00	2.00	1.40	1.00	R1-3	9068499	1.00	1.00	3.00	2.00	3.00	4.00	2.33	1.00	R1,2
9068493	1.00	2.00	3.00	2.00	1.00	2.00	1.83	1.00	R1,5	9068496	1.00	3.00	2.00	1.00	1.00	3.00	1.83	1.00	R1,4,5
9068554	2.00	4.00	2.00	4.00	1.00	1.00	2.33	1.00	R5,6	9068493	1.00	2.00	2.00	2.00	1.00	2.00	1.67	1.00	R1,5
9068566	2.00	2.00	1.00	1.00	3.00	3.00	2.00	1.00	R3,4	9068554	1.00	3.00	2.00	2.00	1.00	1.00	1.67	1.00	R1,5,6
9068584	2.00	2.00	4.00	2.00	1.00	1.00	2.00	1.00	R5,6	9068584	2.00	1.00	3.00	2.00	1.00	3.00	2.00	1.00	R2,5
9057099	2.00	2.00	4.00	3.00	2.00	2.00	2.50	2.00	R1,2,5,6	9068583	5.00	2.00	1.00	1.00	3.00	1.00	2.17	1.00	R3,4,6
9057105	2.00	2.00	2.00	3.00	3.00	4.00	2.67	2.00	R1-3	9057105	2.00	3.00	2.00	3.00	2.00	2.00	2.33	2.00	R1,3,5,6
9057115	2.00	2.00	2.00	3.00	2.00	2.00	2.17	2.00	R1-3,5,6	9057115	3.00	3.00	3.00	2.00	2.00	2.00	2.50	2.00	R4-6
9057116	2.00	2.00	2.00	2.00	2.00	3.00	2.17	2.00	R1-5	9057116	3.00	2.00	2.00	2.00	3.00	2.00	2.33	2.00	R2-4,6
9057117	2.00	2.00	3.00	3.00	3.00	3.00	2.67	2.00	R1,2	9057117	3.00	2.00	3.00	3.00	2.00	3.00	2.67	2.00	R2,5
9057136	2.00	2.00	2.00	3.00	5.00	2.00	2.67	2.00	R1-3,6	9057136	2.00	3.00	2.00	2.00	3.00	3.00	2.50	2.00	R2,5
9057180	3.00	2.00	3.00	2.00	3.00	3.00	2.67	2.00	R2,4	9068486	2.00	3.00	2.00	2.00	3.00	3.00	2.50	2.00	R1,3,4
9057193	4.00	2.00	2.00	3.00	2.00	2.00	2.50	2.00	R2,3,5,6	9057180	2.00	3.00	2.00	2.00	3.00	3.00	2.50	2.00	R1,3,4
9057196	2.00	2.00	2.00	3.00	4.00	3.00	2.67	2.00	R1-3	9068476	6.00	3.00	2.00	3.00	2.00	3.00	3.17	2.00	R3,5
9068476	4.00	3.00	2.00	2.00	2.00	2.00	2.50	2.00	R3-6	9057190	2.00	2.00	2.00	3.00	0.00	3.00	2.00	2.00	R1-3
9057190	3.00	2.00	2.00	2.00	2.00	3.00	2.33	2.00	R2,5	9057189	2.00	2.00	3.00	2.00	3.00	3.00	2.50	2.00	R1,2,4
9057189	3.00	3.00	3.00	2.00	4.00	3.00	3.00	2.00	R4	9068501	2.00	3.00	2.00	2.00	2.00	2.00	2.17	2.00	R1,3-6
9068492	2.00	2.00	3.00	3.00	2.00	3.00	2.50	2.00	R1,2,5	9068492	2.00	2.00	3.00	3.00	2.00	3.00	2.50	2.00	R1,2,5
9068532	3.00	2.00	3.00	5.00	2.00	3.00	3.00	2.00	R2,5	9068495	2.00	2.00	2.00	2.00	2.00	3.00	2.17	2.00	R1-5
9068531	3.00	2.00	2.00	2.00	2.00	4.00	2.50	2.00	R2-5	9068532	3.00	2.00	2.00	7.00	2.00	3.00	3.17	2.00	R2,3,5
9068530	3.00	4.00	3.00	3.00	2.00	2.00	2.83	2.00	R5,6	9068530	3.00	5.00	2.00	3.00	2.00	2.00	2.83	2.00	R3,5,6
9068583	3.00	2.00	2.00	3.00	3.00	3.00	2.67	2.00	R2,3	9068566	2.00	3.00	2.00	2.00	4.00	4.00	2.83	2.00	R1,3,4
9057106	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	R1-6	9068588	3.00	2.00	3.00	2.00	2.00	2.00	2.33	2.00	R2,4-6
Insect/Disease Ratings: 1=None, 9=Severe																			



Study 291134J - Assembly and Evaluation of Eastern Redcedar, <i>Juniper virginia</i> L.																			Table #6
Seed Production																			
1998										2000									
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Ave.	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Ave.	Best	Location
9057099	9.00	5.00	9.00	9.00	9.00	5.00	7.67	9.00	R1,3,-5	9057099	9.00	4.00	9.00	9.00	9.00	5.00	7.50	9.00	R1,3,4,5
9057105	9.00	6.00	9.00	9.00	9.00	3.00	7.50	9.00	R1,3,-5	9057105	9.00	7.00	9.00	9.00	9.00	3.00	7.67	9.00	R1,3,4,5
9057106	6.00	9.00	5.00	9.00	7.00	7.00	7.17	9.00	R2,4	9057106	7.00	9.00	5.00	9.00	8.00	8.00	7.67	9.00	R2,4
9057115	9.00	8.00	9.00	5.00	5.00	4.00	6.67	9.00	R1,3	9057115	9.00	9.00	9.00	4.00	4.00	4.00	6.50	9.00	R1,2,3
9057116	9.00	9.00	9.00	6.00	9.00	7.00	8.17	9.00	R1-3,5	9057116	9.00	9.00	9.00	7.00	9.00	7.00	8.33	9.00	R1-3,5
9057117	9.00	8.00	9.00	8.00	9.00	9.00	8.67	9.00	R1,3,5,6	9057117	9.00	8.00	9.00	8.00	9.00	9.00	8.67	9.00	R1,3,5,6
9057136	9.00	9.00	4.00	6.00	9.00	9.00	7.67	9.00	R1,2,5,6	9057136	9.00	9.00	3.00	5.00	9.00	9.00	7.33	9.00	R1,2,5,6
9068486	9.00	3.00	9.00	6.00	9.00	6.00	7.00	9.00	R1,3,5	9068486	9.00	2.00	9.00	5.00	9.00	9.00	7.17	9.00	R1,3,5,6
9057180	9.00	5.00	9.00	9.00	9.00	8.00	8.17	9.00	R1,3-5	9057180	9.00	4.00	9.00	9.00	9.00	8.00	8.00	9.00	R1,3-5
9057193	9.00	9.00	9.00	4.00	6.00	9.00	7.67	9.00	R1,3,6	9057193	9.00	9.00	9.00	3.00	6.00	9.00	7.50	9.00	R1-3,6
9057196	9.00	8.00	9.00	9.00	9.00	4.00	8.00	9.00	R1,3-5,6	9057196	9.00	8.00	9.00	9.00	9.00	3.00	7.83	9.00	R1,3-5
9057198	5.00	9.00	9.00	6.00	9.00	4.00	7.00	9.00	R2,3,5	9057198	4.00	9.00	9.00	6.00	9.00	3.00	6.67	9.00	R2,3,5
9057199	9.00	6.00	9.00	4.00	1.00	9.00	6.33	9.00	R1,3,6	9057199	9.00	5.00	9.00	4.00	1.00	9.00	6.17	9.00	R1,3,6
9068476	9.00	9.00	4.00	1.00	4.00	8.00	5.83	9.00	R1,2	9068476	9.00	9.00	3.00	1.00	3.00	8.00	5.50	9.00	R1,2
9057190	9.00	6.00	2.00	8.00	9.00	9.00	7.17	9.00	R1,5,6	9057190	9.00	5.00	2.00	8.00	9.00	9.00	7.00	9.00	R1,5,6
9057189	9.00	9.00	8.00	9.00	9.00	9.00	8.83	9.00	R1,2,4-6	9057189	9.00	9.00	7.00	9.00	9.00	9.00	8.67	9.00	R1,2,4-6
9068504	9.00	9.00	9.00	9.00	9.00	3.00	8.00	9.00	R1-5	9068504	9.00	9.00	9.00	9.00	9.00	2.00	7.83	9.00	R1-5
9068503	9.00	9.00	1.00	4.00	9.00	4.00	6.00	9.00	R1,2,5	9068503	9.00	9.00	1.00	4.00	9.00	3.00	5.83	9.00	R1,2,5
9068502	9.00	9.00	9.00	9.00	9.00	9.00	9.00	9.00	R1-6	9068502	9.00	9.00	9.00	9.00	9.00	9.00	9.00	9.00	R1-6
9068501	9.00	7.00	4.00	1.00	9.00	9.00	6.50	9.00	R1,5,6	9068501	9.00	8.00	3.00	1.00	9.00	9.00	6.50	9.00	R1,5,6
9068500	6.00	9.00	1.00	4.00	8.00	9.00	6.17	9.00	R2,6	9068500	7.00	9.00	1.00	3.00	8.00	9.00	6.17	9.00	R2,6
9068492	9.00	6.00	9.00	4.00	9.00	9.00	7.67	9.00	R1,3,5,6	9068492	9.00	5.00	9.00	3.00	9.00	9.00	7.33	9.00	R1,3,5,6
9068499	6.00	8.00	9.00	9.00	9.00	1.00	7.00	9.00	R3,4,5	9068499	7.00	9.00	9.00	9.00	1.00	7.00	7.00	9.00	R2,-4
9068496	9.00	6.00	9.00	3.00	6.00	9.00	7.00	9.00	R1,3,9	9068496	9.00	5.00	9.00	3.00	5.00	9.00	6.67	9.00	R1,3,6
9068495	9.00	9.00	9.00	9.00	9.00	8.00	8.83	9.00	R1-5	9068495	9.00	9.00	9.00	9.00	9.00	8.00	8.83	9.00	R1-6
9068493	9.00	8.00	9.00	9.00	-	9.00		9.00	R1,3,4,6	9068493	9.00	7.00	9.00	9.00	-	8.00	7.00	9.00	R1,3,4
9068532	9.00	9.00	6.00	9.00	9.00	3.00	7.50	9.00	R1,2,4,5	9068532	9.00	9.00	7.00	9.00	9.00	4.00	7.83	9.00	R1,2,4,5
9068531	6.00	9.00	1.00	1.00	3.00	9.00	4.83	9.00	R2,6	9068531	7.00	9.00	1.00	1.00	2.00	9.00	4.83	9.00	R2,6
9068530	9.00	9.00	3.00	9.00	6.00	9.00	7.50	9.00	R1,2,4,6	9068530	9.00	9.00	2.00	9.00	5.00	9.00	7.17	9.00	R1,2,4,6
9068554	6.00	9.00	3.00	1.00	4.00	9.00	5.33	9.00	R2,6	9068554	7.00	9.00	3.00	9.00	7.00	9.00	7.33	9.00	R2,4,6
9068566	6.00	6.00	6.00	3.00	4.00	4.00	4.83	6.00	R1-3	9068566	7.00	7.00	7.00	2.00	4.00	4.00	5.17	7.00	R1-3
9068584	9.00	8.00	8.00	9.00	6.00	9.00	8.17	9.00	R1,6	9068584	9.00	9.00	9.00	9.00	7.00	9.00	8.67	9.00	R1-4,6
9068583	9.00	3.00	8.00	3.00	1.00	9.00	5.50	9.00	R1,R6	9068583	9.00	5.00	9.00	4.00	1.00	9.00	6.17	9.00	R1,3,6
9068588	9.00	9.00	3.00	3.00	8.00	9.00	6.83	9.00	R1,2,6	9068588	9.00	9.00	4.00	4.00	9.00	9.00	7.33	9.00	R1,2,5,6

**Study:** 29I135J

**Study Title:** Assembly and Evaluation of Hazelnut, *Corylus americana* Walt.

**Study Leader:** Henry, J.

**Introduction:**

American hazelnut is a shrub or very small tree probably native to every county in Missouri. It commonly occurs in dry or moist thickets, woodland, and borders of woodland, in valleys and upland. It ranges from Maine to Saskatchewan, south to Georgia, Arkansas, and Oklahoma. Leaves are borne simply on bristly stalks, the bristles somewhat glandular. Flowers are separate with male and female flowers on the same tree. Male catkins droop and form the season before opening. Female flowers are enclosed in a scaly bud. They have red stigmas that protrude at the tip of the bud. The fruit is a globe-shape nut enclosed in a large, leaf-like covering. This species flowers March-May with fruit ripening July-September.

**Problem:**

There is a lack of an available cultivar of American hazelnut specifically for this area. A need for developing a selection, source identified, and sources of hazelnut for use as wildlife habitat and for agroforestry in the three states being served by the Center has been identified by NRCS and other conservation and wildlife agencies.

**Objective:**

The objective is to assemble, comparatively evaluate, select and release an adapted cultivar of source identified or selected hazelnut.

**Discussion**

**1989 - 1994**

Collections of hazelnut were assembled at the PMC between 1989 and 1992. Thirty-six accessions from Illinois and Missouri were stratified and placed in the greenhouse in 1993. Twenty-one accessions germinated and were grown out in two-gallon containers. These accessions were placed in a randomized complete block with eight replications. The planting was established May 3 and 4, 1989, in Field #11E on the PMC.

The summer of 1994 had several significant dry spells and considerable time was spent irrigating. Many plants were stressed, lost leaves, and resprouted. Four plants in the evaluation block failed to survive in 1994.

**1995-1998**

The assembly was evaluated in 1995, 1996, 1997 and 1998. Of the original 138 plants being evaluated a total of 11 died. The survival was good but the rate of growth seems to be slow, which seems characteristic of hazelnuts.

The following accessions were selected in 1997 for field plantings: 9057168 and 9057169 (Iroquois County, Illinois), 9057188 and 9068528 (Coles County, Illinois), 9068562 (Adams County, Illinois), and 9068573 and 9068574 both from Chariton County, Missouri. The selection criteria for these accessions are as follows: form, growth, height, width and fruit production and resistance to insect and disease.

## 1999

The selected accessions continue to be utilized in the plant materials field-planting program throughout the PMC service area. The plants' performance data for 1999 was recorded only for nut production. This information can be found in the following tables.

Nut production for the selected accessions for 1998:

9057168	=	1.75 pounds	9057169	=	1.00 pound
9057188	=	1.90 pounds	9068528	=	1.00 pound
9068562	=	1.67 pounds	9068573	=	1.50 pounds
9068574	=	1.30 pounds			

Nut production for the selected accession for 1999.

9057169	=	1.4 pounds	9068528	=	2.2 pounds
9057188	=	0.5 pound	9068573	=	1.9 pounds
9068562	=	2.7 pounds	9057168	=	1.8 pounds
9068574	=	4.3 pounds			

## 2000

Nut production is being harvested from those accessions selected for field plantings in the service area of the PMC (Iowa, Illinois and Missouri). One-tenth of the nut production for each of the selected accessions was left on the shrubs to determine the dates the fruits would fall to the ground. The following chart reflects the selected accessions, fruit production and dates nuts fell to the ground. There were no plant evaluations on the assembly of plants this year.

Accession Numbers	Nut Production With Husks	Date Nut Dropped
9057188	1.4 Pounds	11/27/00
9068562	10.3 Pounds	11/27/00
9068574	4.6 Pounds	11/27/00
9068528	12.2 Pounds	11/27/00
9068573	3.7 Pounds	11/27/00
9057168	3.2 Pounds	11/16/00

## 2001 - 2002

The following table reflects the performance of those accessions initially selected and placed in the plant materials field planting program in the PMC service area for years 1997 through 2002. As reflected in the PMC Business Plan, releases from these accessions will be made in 2004 – 2005. One plant out of the following accessions will be moved to a crossing block in Field #6 on the PMC in the February 2003. The progeny from this crossing block will be assigned a separate accession number and only one release (Tested Class) will be made for the PMC service area. These accessions are: 9057168, 9068562, 9068573, 9068574, 9057188, and 9068528. The remaining plants in the assembly will be left until the area is needed for a new study.

**Table #1 reflects performance data 1997 – 2002.**

**Table # 1**

Acc. Number	1997	1998	1999	2000	2001	2002	Averages
<b>9057168</b>							
Height (Ft.)	4.3	5.4	5.7	6	6.5	6.8	5.8
Spread (Ft.)	4.2	7	7.3	7.5	8	8.3	7.05
Ins/Disease	2	2	2	2	2	2	2
Form	3	3	3	3	3	3	3
Nut Prod.		1.8 lbs.	1.3 lbs.	2.0 lbs.	2.3 lbs.	1.5 lbs	1.8 lbs.
<b>9068562</b>							
Height	5.2	7	7.4	8	8.2	8.5	7.4
Spread	6.5	7.4	7.6	8	8.5	8.8	7.8
Ins/Disease	2	3	2	2	2	2	2.9
Form	2	2	2	1	2	2	1.8
Nut Prod.		1.67 lbs.	1.60 lbs.	1.7 lbs.	1.9 lbs.	6.2	2.6 lbs.
<b>9068573</b>							
Height	4.6	6.3	6.5	6.7	7.1	7.3	6.4
Spread	5	6	6.3	6.5	7	7.5	6.4
Ins/Disease	2	2	2	2	2	2	2
Form	3	3	3	3	3	3	3
Nut Prod.		1.5 lbs.	1.9 lbs.	2.6 lbs.	4.3 lbs.	2.6 lbs	2.6 lbs.
<b>9068574</b>							
Height	6.8	6.9	7	7.3	7.5	7.9	7.2
Spread	4.5	5.8	6	6.3	6.5	6.8	6.0
Ins/Disease	2	3	2	2	2	2	2.2
Form	3	4	3	3	3	4	3.3
Nut Prod.		1.3 lbs.	1.8 lbs.	1.3 lbs.	2.1 lbs.	3.5	2.0 lbs.
<b>9057188</b>							
Height	5.1	6.4	6.7	6.8	7	7.3	6.6
Spread	3.7	7	7.5	7.8	8	8.2	7.03
Ins/Disease	2	2	2	2	2	2	2
Form	3	3	3	3	3	3	3
Nut Prod.		1.0 lbs.	0.5 lb.	1.4 lbs.	1.9 lbs.	8.2 lbs.	2.6 lbs.
<b>9068528</b>							
Height	3.5	4.3	5.0	6.3	6.7	7.0	5.4
Spread	3.0	4.1	5.5	5.8	6.3	6.8	5.3
Ins/Disease	2	2	2	2	2	2	2
Form	5	4	3	3	3	3	3.5
Nut Prod.	0	1.0 lbs.	2.2 lbs.	12.2 lbs.	4.2 lbs.	6.3 lbs.	4.3 lbs.

**Table #2 reflects accession information****Table #2**

<b>Accession Number</b>	<b>State or Origin</b>	<b>City or County</b>
9057081	Illinois	Coles
9057082	Illinois	Coles
9057087	Illinois	Coles
9057119	Illinois	Whiteside
9057120	Illinois	Carroll
9057167	Illinois	Will
9057168	Illinois	Iroquois
9057169	Illinois	Iroquois
9057184	Illinois	Clark
9057186	Illinois	Coles
9057188	Illinois	Coles
9057192	Illinois	Montgomery
9057195	Illinois	Morgan
9068505	Illinois	Coles
9068507	Illinois	Cumberland
9068508	Illinois	Mercer
9068509	Illinois	Ogle
9068510	Illinois	Iroquois
9068511	Illinois	Effingham
9068512	Illinois	Clay
9068513	Illinois	Pike
9068525	Illinois	Cumberland
9068526	Illinois	Coles
9068527	Illinois	Maultrie
9068528	Illinois	Coles
9068529	Illinois	Vermilion
9068562	Illinois	Adams
9068565	Illinois	Jo Daviess
9068585	Illinois	DeWitt
9068586	Illinois	Vermilion
9068570	Missouri	Lincoln
9068573	Missouri	Chariton
9068574	Missouri	Chariton
9068575	Illinois	Johnson

**Tables # 3 – 6 reflect the performance data for all accessions included in this study for 1995 – 1999.**

Study 291135J - Assembly and Evaluation of Hazelnut, <i>Corylus americana</i> , Walt.													Table #3										
													Height in Feet										
1995													1997										
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Tallest	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Tallest	Location
9068562	1.2	2.5	1.4	1.3	1.5	1.7	2.9	4.0	2.1	4.0	R8	9068574	4.9	4.3	3.8	3.9	6.8	3.8	3.2	2.2	4.1	6.8	R5
9057188	2.6	4.0	1.6	3.1	2.6	2.0	2.3	2.2	2.6	4.0	R2	9068562	3.3	5.2	2.7	2.7	3.4	4.6	4.2	4.5	3.8	5.2	R2
9068573	3.6	2.7	3.2	1.5	3.0	2.2	2.5	3.2	2.7	3.6	R1	9057188	4.0	5.0	2.9	4.2	5.1	3.7	4.7	4.0	4.2	5.1	R5
9068508	2.0	3.0	2.2	2.3	1.3	1.0	1.6	1.5	1.9	3.0	R2	9057169	5.0	4.1	3.4	3.5	2.3	3.6	3.2	2.8	3.5	5.0	R1
9068574	1.7	2.0	1.7	3.0	2.3	2.2	1.3	2.0	2.0	3.0	R4	9057168	3.8	1.2	4.6	2.4	4.3	4.1	3.0	2.0	3.2	4.6	R3
9057169	2.9	1.6	1.4	1.7	0.8	1.0	1.4	1.6	1.6	2.9	R1	9068573	4.2	4.5	4.0	3.4	4.6	3.1	2.5	3.4	3.7	4.6	R4
9068507	1.7	1.0	2.6	Dead	Dead	2.0	1.3	1.8	1.7	2.6	R 3	9068528	4.5	4.2	Dead	4.0	3.1	3.2	3.0	2.8	3.5	4.5	R1
9068565	2.3	2.6	2.5	2.0	2.4	2.2	1.6	Dead	2.2	2.6	R2	9068510	3.1	2.0	3.0	4.5	4.3	2.8	2.0	4.0	3.2	4.5	R4
9068558	1.5	2.2	1.7	1.3	2.0	1.5	2.5	Dead	1.8	2.5	R7	9068558	3.6	Dead	2.4	3.5	2.8	4.3	3.9	Dead	3.4	4.3	R6
9057168	1.3	1.3	2.1	1.0	1.9	2.2	1.4	0.9	1.5	2.2	R6	9068507	2.3	Dead	3.5	Dead	Dead	4.0	2.0	2.3	2.8	4.0	R6
9068510	0.6	1.3	2.1	1.7	1.5	1.4	0.6	2.2	1.4	2.2	R8	9068565	2.7	3.3	2.3	3.0	4.0	2.8	1.6	Dead	2.8	4.0	R5
9068528	1.3	1.2	Dead	2.1	Dead	1.7	2.0	1.4	1.6	2.1	R4	9068525	3.3	2.3	4.0	3.6	Dead	3.1	Dead	3.2	2.8	4.0	R3
9068586	Dead	Dead	1.2	1.7	2.0	2.0	1.0	1.3	1.5	2.0	R5,6	9068508	3.2	3.6	3.9	3.3	3.4	2.8	3.5	3.3	3.4	3.9	R3
9068525	1.3	1.2	1.0	1.0	1.0	1.5	Dead	1.7	1.2	1.7	R8	9068586	Dead	Dead	2.9	2.6	3.7	3.0	2.0	3.1	2.9	3.1	R8
1996													1998										
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Tallest	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Tallest	Location
9057188	3.3	4.1	2.6	3.2	4.1	3.2	3.4	2.9	3.4	4.1	R2, 5	9068562	4.7	7.0	4.0	4.6	5.1	4.1	4.6	5.4	4.9	7.0	R2
9068562	2.0	3.8	1.7	1.0	2.7	2.8	3.2	4.1	2.7	3.8	R2	9068558	4.6	Dead	5.0	4.3	4.1	5.0	6.4	Dead	4.9	6.4	R7
9068586	Dead	Dead	2.9	2.6	3.7	3.0	2.0	2.0	2.7	3.7	R5	9057188	4.0	5.8	6.0	5.0	6.4	5.8	5.0	5.7	5.5	6.4	R5
9068573	2.6	3.7	3.4	2.1	3.6	3.0	2.8	3.3	3.1	3.7	R2	9068573	6.3	4.9	5.2	5.0	6.3	5.0	6.0	4.0	5.3	6.3	R5
9068574	3.2	2.3	2.4	3.7	3.5	2.6	2.7	2.0	2.8	3.5	R5	9068574	5.2	5.3	5.0	4.0	6.3	3.2	3.6	3.0	4.5	6.3	R5
9068508	2.3	3.4	3.3	2.5	1.7	1.4	2.5	2.3	2.4	3.4	R2	9057169	5.9	5.2	5.0	5.0	3.2	4.4	3.2	3.3	4.4	5.9	R1
9057168	2.3	1.3	3.3	1.8	3.3	3.0	1.8	1.3	2.3	3.3	R3, 5	9057168	5.0	1.8	5.4	3.8	5.4	5.1	4.2	3.0	4.2	5.4	R5
9068528	3.0	3.2	Dead	3.3	Dead	2.5	2.5	2.1	2.8	3.3	R4	9068528	5.4	4.4	Dead	4.2	4.0	4.0	4.8	3.2	4.3	5.4	R1
9068507	2.1	1.3	3.2	Dead	Dead	2.9	2.0	1.5	2.2	3.2	R3	9068510	3.9	4.8	4.0	4.6	5.4	3.0	4.0	4.6	4.3	5.4	R5
9068558	2.0	Dead	2.1	2.1	2.4	3.2	2.7	Dead	2.4	3.2	R6	9068507	2.3	Dead	4.3	Dead	Dead	5.2	2.8	4.0	3.7	5.2	R6
9057169	2.9	3.1	2.3	2.7	1.6	2.2	2.1	1.9	2.4	3.1	R2	9068525	4.2	3.5	5.2	4.9	Dead	3.4	Dead	4.6	3.7	5.2	R3
9068565	2.3	2.9	2.3	2.3	2.6	2.3	1.4	Dead	2.3	2.9	R2	9068586	Dead	Dead	4.2	4.0	5.0	4.6	3.5	4.1	4.2	5.0	R5
9068510	1.8	2.2	1.7	2.2	2.7	2.3	1.3	2.7	2.1	2.7	R5,8	9068508	3.5	3.8	3.2	4.8	4.7	3.8	4.2	4.0	4.0	4.8	R4
9068525	2.2	1.6	1.7	2.5	1.6	1.9	Dead	2.5	2.0	2.5	R4,8	9068565	2.9	4.8	3.2	Dead	4.4	4.0	3.4	Dead	3.8	4.8	R2
Height Measured in Feet																							

Study 291135J - Assembly and Evaluation of Hazelnut, <i>Corylus americana</i> , Walt.												Table #4											
1995												1997											
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location
9057188	1.0	0.7	0.6	1.2	1.4	0.9	0.9	2.0	1.1	<b>2.0</b>	R8	9068562	3.3	6.5	2.3	2.3	3.8	3.7	3.5	4.2	3.7	<b>6.5</b>	R2
9068562	0.4	1.4	0.3	0.4	0.4	0.6	0.4	1.5	0.7	<b>1.5</b>	R8	9068573	4.1	3.5	4.3	5.1	5.0	3.6	2.5	2.9	3.9	<b>5.1</b>	R4
9068573	1.5	0.6	0.8	0.8	1.0	0.7	0.9	0.3	0.8	<b>1.5</b>	R1	9057188	3.6	5.0	4.2	4.7	3.7	4.5	4.0	4.4	4.3	<b>5.0</b>	R2
9068574	1.5	0.8	1.0	1.0	0.9	0.9	0.6	0.4	0.9	<b>1.5</b>	R1	9057169	3.6	5.0	4.2	4.7	3.7	4.5	4.0	4.4	4.3	<b>5.0</b>	R2
9068507	0.6	0.3	1.2	Dead	Dead	1.0	0.3	0.3	0.6	<b>1.2</b>	R3	9068574	4.9	4.4	4.6	3.7	4.5	3.2	3.0	2.0	3.8	<b>4.9</b>	R1
9068510	0.2	1.2	0.6	0.4	0.9	0.6	0.2	0.8	0.6	<b>1.2</b>	R2	9057168	4.4	1.5	4.2	2.0	4.2	3.3	2.5	2.0	3.0	<b>4.4</b>	R1
9057168	0.7	0.4	1.1	0.4	1.1	0.8	0.7	0.5	0.7	<b>1.1</b>	R3, 5	9068528	3.0	4.4	Dead	3.3	2.9	2.0	3.4	2.3	3.0	<b>4.4</b>	R2
9068558	0.3	0.3	0.5	0.7	0.9	1.1	0.7	Dead	0.6	<b>1.1</b>	R6	9068508	4.0	Dead	3.2	3.7	3.9	3.0	3.4	3.4	3.5	<b>4.0</b>	R1
9068586	Dead	Dead	0.4	0.6	1.0	0.9	0.1	0.2	0.5	<b>1.0</b>	R5	9068510	3.0	3.2	3.0	3.3	3.9	2.1	4.0	3.3	3.2	<b>4.0</b>	R7
9057169	1.0	0.8	0.6	0.4	0.2	0.5	0.7	0.4	0.6	<b>1.0</b>	R1	9068525	4.0	3.3	4.0	3.4	Dead	2.0	Dead	4.0	3.0	<b>4.0</b>	R1, 3, 8
9068508	0.5	0.4	0.4	0.8	0.6	0.9	0.8	0.8	0.7	<b>0.9</b>	R6	9068586	Dead	Dead	3.7	2.5	3.1	3.5	1.8	2.8	2.9	<b>3.7</b>	R3
9068565	0.6	0.4	0.9	0.8	0.5	0.7	0.7	Dead	0.7	<b>0.9</b>	R3	9068558	3.2	1.5	3.2	3.0	2.7	3.5	3.3	Dead	2.9	<b>3.5</b>	R6
9068528	0.8	0.6	Dead	0.6	Dead	0.5	0.6	0.3	0.6	<b>0.8</b>	R1	9068565	2.8	3.5	2.2	2.0	3.1	3.0	1.5	Dead	2.6	<b>3.5</b>	R2
9068525	0.4	0.4	0.4	0.3	0.3	0.3	Dead	0.6	0.4	<b>0.6</b>	R8	9068507	2.3	Dead	3.0	Dead	Dead	3.2	1.0	1.8	2.3	<b>3.0</b>	R3
1996												1998											
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location
9057188	2.4	2.8	2.4	2.6	2.9	3.3	2.3	3.7	2.8	<b>3.7</b>	R8	9057188	4.6	7.5	5.4	5.4	7.7	7.0	4.8	6.0	6.1	<b>4.6</b>	R1
9068562	1.8	3.6	1.0	0.9	2.2	2.7	1.8	3.3	2.2	<b>3.6</b>	R2	9068508	4.4	5.8	4.4	5.2	4.8	5.4	4.6	4.9	4.9	<b>4.4</b>	R1, 3, 8
9068574	2.8	3.1	2.8	2.3	2.5	1.9	3.4	1.1	2.5	<b>3.4</b>	R 7	9068573	7.0	5.5	5.4	6.0	6.0	5.4	5.7	4.3	5.7	<b>4.3</b>	R8
9068573	3.1	2.7	2.3	2.4	3.0	2.2	2.4	1.2	2.4	<b>3.1</b>	R1	9068558	4.0	Dead	5.0	4.4	4.0	5.0	5.2	Dead	4.6	<b>4.0</b>	R1, 5
9057169	3.1	2.5	3.0	2.4	0.8	2.4	1.3	1.0	2.1	<b>3.1</b>	R1	9068528	4.3	4.6	Dead	4.0	4.4	3.4	3.8	4.0	4.1	<b>3.4</b>	R6
9057168	2.8	1.0	2.9	1.4	2.8	2.1	2.1	1.2	2.0	<b>2.9</b>	R3	9068525	3.4	4.8	5.7	5.2	Dead	3.4	Dead	4.6	3.9	<b>3.4</b>	R1,6
9068508	2.0	2.5	2.3	2.2	2.4	1.7	2.8	1.8	2.2	<b>2.8</b>	R7	9068562	4.2	7.4	4.0	3.3	5.0	5.5	5.1	5.8	5.0	<b>3.3</b>	R4
9068510	1.6	2.7	2.1	1.8	2.6	1.8	1.0	0.6	1.8	<b>2.7</b>	R2	9068510	3.4	3.2	4.0	4.2	4.8	3.5	3.5	4.0	3.8	<b>3.2</b>	R2
9068586	Dead	Dead	2.6	1.5	1.5	2.0	1.1	1.6	1.7	<b>2.6</b>	R3	9057169	4.8	4.6	5.3	5.2	2.8	4.3	3.5	4.0	4.3	<b>2.8</b>	R5
9068565	1.0	2.4	1.6	2.0	1.7	2.6	1.0	Dead	1.8	<b>2.6</b>	R6	9057168	4.0	2.6	6.0	3.4	7.0	5.0	4.6	3.2	4.5	<b>2.6</b>	R2
9068558	1.7	Dead	2.4	2.5	2.0	2.1	2.5	Dead	2.2	<b>2.5</b>	R 4,7	9068574	2.4	5.3	5.2	2.6	5.8	3.8	4.5	3.3	4.1	<b>2.4</b>	R1
9068528	2.2	2.3	Dead	2.2	1.7	2.4	2.4	1.8	2.1	<b>2.4</b>	R6, 7	9068565	4.0	4.6	3.0	Dead	5.0	4.2	2.3	Dead	3.9	<b>2.3</b>	R7
9068525	1.7	2.2	2.0	2.0	1.4	2.0	Dead	2.3	1.9	<b>2.3</b>	R8	9068586	Dead	Dead	4.9	4.0	3.8	3.5	2.1	4.1	3.7	<b>2.1</b>	R7
9068507	1.4	0.8	2.1	Dead	Dead	2.3	1.4	0.6	1.4	<b>2.1</b>	R3	9068507	2.7	Dead	5.0	Dead	Dead	6.0	1.3	4.6	3.9	<b>1.3</b>	R7
Width Measured in Feet																							

Study 291135J - Assembly and Evaluation of Hazelnut, <i>Corylus americana</i> , Walt.													Table #5												
													Form												
1995													1997												
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location		
9057188	3.0	4.0	5.0	5.0	5.0	5.0	5.0	2.0	4.3	2.0	R8	9068562	5.0	3.0	5.0	7.0	7.0	4.0	8.0	3.0	5.3	3.0	R2,8		
9068562	3.0	3.0	3.0	4.0	3.0	5.0	6.0	3.0	3.8	3.0	R1,2,3,5,8	9057168	5.0	8.0	4.0	8.0	3.0	5.0	6.0	7.0	5.8	3.0	R5		
9057168	5.0	8.0	3.0	6.0	4.0	5.0	6.0	7.0	5.5	3.0	R3	9068558	4.0	Dead	5.0	5.0	6.0	5.0	3.0	Dead	4.7	3.0	R7		
9068558	7.0	8.0	5.0	7.0	3.0	4.0	7.0	Dead	5.9	3.0	R5	9068573	7.0	4.0	5.0	5.0	3.0	5.0	5.0	6.0	5.0	3.0	R5		
9068508	5.0	7.0	8.0	5.0	6.0	3.0	5.0	6.0	5.6	3.0	R6	9057188	3.0	4.0	4.0	4.0	3.0	5.0	3.0	4.0	3.8	3.0	R1,5,7		
9068573	3.0	4.0	5.0	5.0	4.0	5.0	4.0	6.0	4.5	3.0	R1	9068565	7.0	3.0	6.0	8.0	5.0	5.0	7.0	Dead	5.9	3.0	R2,8		
9068507	5.0	7.0	4.0	Dead	Dead	5.0	6.0	6.0	5.5	4.0	R3	9068510	7.0	8.0	6.0	5.0	5.0	4.0	6.0	3.0	5.5	3.0	R8		
9057169	4.0	5.0	5.0	8.0	6.0	6.0	6.0	6.0	5.8	4.0	R1	9068574	7.0	6.0	4.0	6.0	3.0	6.0	6.0	6.0	5.5	3.0	R8		
9068510	8.0	5.0	4.0	5.0	8.0	8.0	5.0	6.0	6.1	4.0	R3,4,6	9068507	5.0	Dead	4.0	5.0	Dead	4.0	8.0	6.0	5.3	4.0	R3,6		
9068574	4.0	6.0	4.0	6.0	6.0	6.0	6.0	6.0	5.5	4.0	R1	9068586	Dead	Dead	6.0	7.0	4.0	5.0	6.0	5.0	5.5	4.0	R4		
9068565	5.0	6.0	7.0	5.0	6.0	5.0	7.0	Dead	5.9	5.0	R1,4,6	9068508	7.0	5.0	5.0	5.0	5.0	7.0	6.0	4.0	5.5	4.0	R8		
9068528	5.0	5.0	Dead	5.0	Dead	6.0	6.0	6.0	5.5	5.0	R1,2,4	9057169	4.0	4.0	6.0	4.0	7.0	5.0	5.0	8.0	5.4	4.0	R1,2,4		
9068525	6.0	6.0	5.0	8.0	6.0	8.0	Dead	6.0	6.4	5.0	R3	9068528	4.0	4.0	Dead	5.0	6.0	4.0	6.0	6.0	5.0	4.0	R1,3,6		
9068586	Dead	Dead	6.0	6.0	7.0	6.0	9.0	8.0	7.0	6.0	R3,4,6	9068525	5.0	6.0	7.0	8.0	Dead	8.0	Dead	5.0	6.4	5.0	R1,8		
1996													1998												
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location		
9068573	3.0	4.0	4.0	6.0	4.0	4.0	4.0	5.0	4.3	3.0	R1	9068586	5.0	Dead	3.0	6.0	5.0	7.0	7.0	2.0	5.0	2.0	R8		
9057188	3.0	5.0	5.0	4.0	4.0	4.0	4.0	5.0	4.3	3.0	R1	9068562	5.0	2.0	2.0	5.0	3.0	5.0	6.0	2.0	3.8	2.0	R2,3,8		
9057169	3.0	5.0	6.0	5.0	4.0	5.0	5.0	5.0	4.8	3.0	R1	9068558	3.0	5.0	3.0	5.0	3.0	2.0	2.0	Dead	3.3	2.0	R6,7		
9068507	4.0	5.0	4.0	Dead	Dead	4.0	4.0	5.0	4.3	4.0	R1,3,6,7	9068574	5.0	2.0	3.0	6.0	5.0	6.0	3.0	5.0	4.4	2.0	R2		
9068586	Dead	Dead	5.0	7.0	4.0	5.0	5.0	4.0	5.0	4.0	R3,8	9057168	5.0	7.0	5.0	5.0	3.0	5.0	5.0	7.0	5.3	3.0	R5		
9068562	5.0	5.0	4.0	7.0	5.0	4.0	5.0	4.0	4.9	4.0	R6,8	9068573	5.0	5.0	5.0	3.0	3.0	3.0	4.0	5.0	4.1	3.0	R4,5,6		
9057168	6.0	6.0	5.0	6.0	4.0	4.0	6.0	6.0	5.4	4.0	R5,6	9057188	6.0	5.0	3.0	6.0	3.0	3.0	5.0	3.0	4.3	3.0	R3,5,6,8		
9068558	4.0	Dead	6.0	5.0	6.0	5.0	5.0	Dead	5.2	4.0	R1	9068528	3.0	5.0	3.0	5.0	3.0	7.0	5.0	6.0	4.6	3.0	R1,3,5		
9068565	5.0	4.0	6.0	7.0	5.0	6.0	5.0	Dead	5.4	4.0	R2	9068510	5.0	7.0	5.0	3.0	3.0	7.0	7.0	5.0	5.3	3.0	R4,5		
9068528	5.0	4.0	Dead	5.0	5.0	5.0	6.0	5.0	5.0	4.0	R2	9068565	5.0	5.0	7.0	Dead	5.0	5.0	4.0	Dead	5.2	4.0	R7		
9068510	5.0	7.0	6.0	4.0	5.0	4.0	4.0	5.0	5.0	4.0	R4,6,7	9068507	7.0	Dead	5.0	Dead	Dead	5.0	7.0	7.0	6.2	5.0	R3,6		
9068574	5.0	7.0	4.0	5.0	4.0	5.0	5.0	5.0	5.0	4.0	R3,5	9068508	Dead	5.0	7.0	5.0	7.0	5.0	6.0	5.0	5.7	5.0	R2,4,6,8		
9068508	7.0	5.0	5.0	5.0	5.0	7.0	5.0	5.0	5.5	5.0	R2,3,4,5,7,8	9057169	7.0	5.0	7.0	5.0	7.0	5.0	6.0	5.0	5.9	5.0	R2,4,6,8		
9068525	5.0	5.0	5.0	6.0	6.0	6.0	Dead	6.0	5.6	5.0	R1,2,3,	9068525	5.0	7.0	5.0	7.0	Dead	7.0	Dead	6.0	6.0	5.0	R1,3,5		

Rating: 1-Excellent, 9=Poor

Study 291135J - Assembly and Evaluation of Hazelnut, <i>Corylus americana</i> , Walt.												Table #6											
												Fruit Production											
1997												1998											
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location
9057169	2.0	3.0	9.0	9.0	0.0	0.0	0.0	0.0	5.8	2.0	R1	9068507	5.0	Dead	5.0	Dead	Dead	2.0	0.0	0.0	4.0	2.0	R6
9068562	0.0	7.0	0.0	0.0	0.0	3.0	9.0	7.0	6.5	3.0	R6	9068586	Dead	Dead	7.0	7.0	7.0	7.0	5.0	2.0	5.8	2.0	R8
9057168	9.0	9.0	3.0	0.0	7.0	9.0	0.0	0.0	7.4	3.0	R3	9068562	2.0	2.0	7.0	0.0	7.0	5.0	2.0	2.0	3.9	2.0	R1,2,7,8
9057188	3.0	7.0	Dead	9.0	9.0	9.0	7.0	7.0	7.3	3.0	R1, R7	9057168	7.0	5.0	2.0	0.0	2.0	5.0	7.0	0.0	4.7	2.0	R3,5
9068574	6.0	0.0	0.0	8.0	3.0	0.0	0.0	0.0	5.7	3.0	R5	9068558	2.0	Dead	5.0	2.0	0.0	5.0	5.0	Dead	3.8	2.0	R2,4
9068573	3.0	6.0	9.0	0.0	6.0	0.0	0.0	0.0	6.0	6.0	R2, R5	9068508	5.0	5.0	2.0	5.0	2.0	5.0	2.0	2.0	3.5	2.0	R1,2,3,5,7,8
9068528	9.0	6.0	0.0	9.0	0.0	6.0	8.0	0.0	7.6	6.0	R2,6	9068573	7.0	2.0	2.0	5.0	2.0	7.0	5.0	7.0	4.6	2.0	R2,3,5
9068510	0.0	7.0	0.0	0.0	6.0	0.0	0.0	0.0	6.5	6.0	R5	9068565	7.0	7.0	2.0	7.0	0.0	2.0	5.0	0.0	5.0	2.0	R3,6
9068507	0.0	Dead	7.0	Dead	Dead	0.0	0.0	0.0	7.0	7.0	R3	9057169	7.0	7.0	2.0	7.0	0.0	2.0	5.0	0.0	5.0	2.0	R3,6
9068565	8.0	0.0	9.0	7.0	9.0	9.0	0.0	Dead	8.4	7.0	R4	9068528	2.0	2.0	Dead	5.0	2.0	5.0	5.0	2.0	3.3	2.0	R1,2,5,8
9068508	9.0	Dead	9.0	0.0	9.0	0.0	9.0	8.0	8.8	8.0	R8	9068510	7.0	2.0	7.0	7.0	7.0	5.0	0.0	5.0	5.7	2.0	R2
9068558	9.0	Dead	0.0	0.0	0.0	0.0	9.0	Dead	9.0	9.0	R1, R7	9068574	5.0	7.0	7.0	2.0	2.0	5.0	5.0	0.0	4.7	2.0	R4,5
9068525	0.0	0.0	0.0	0.0	0.0	0.0	Dead	9.0	9.0	9.0	R8	9068525	5.0	5.0	7.0	7.0	2.0	7.0	Dead	2.0	5.0	2.0	R5,8
9068586	Dead	Dead	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-	-	9057188	7.0	7.0	5.0	7.0	5.0	0.0	0.0	Dead	6.2	5.0	R3,5
1=Heavy Fruit Production; 9=Poor Fruit Production												1=Heavy Fruit Production; 9=Poor Fruit Production											
												Insect/Disease											
1997												1998											
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location
9068586	Dead	Dead	4.0	3.0	4.0	4.0	5.0	4.0	4.0	2.0	R2	9068507	4.0	Dead	6.0	Dead	Dead	2.0	9.0	3.0	4.8	2.0	R6
9068562	3.0	2.0	4.0	5.0	4.0	2.0	3.0	2.0	3.1	2.0	R2	9068586	Dead	Dead	4.0	3.0	3.0	3.0	4.0	2.0	3.2	2.0	R8
9057168	3.0	4.0	3.0	3.0	2.0	3.0	4.0	4.0	3.3	2.0	R5	9057168	2.0	4.0	3.0	4.0	2.0	2.0	6.0	3.0	3.3	2.0	R1,5,6
9068558	2.0	Dead	3.0	5.0	7.0	3.0	3.0	Dead	3.8	2.0	R1,3,6	9068558	3.0	Dead	4.0	3.0	3.0	2.0	2.0	Dead	2.8	2.0	R6,7
9068508	3.0	3.0	3.0	3.0	2.0	3.0	4.0	5.0	3.3	2.0	R5	9068573	5.0	3.0	2.0	3.0	2.0	3.0	3.0	3.0	3.0	2.0	R5
9068573	8.0	3.0	3.0	2.0	2.0	3.0	3.0	3.0	3.4	2.0	R4, 5	9057188	7.0	3.0	3.0	2.0	3.0	2.0	2.0	4.0	3.3	2.0	R4,6,7
9057188	2.0	2.0	2.0	4.0	2.0	6.0	2.0	2.0	2.8	2.0	R1,2,3,5,7,8	9057169	2.0	4.0	4.0	3.0	2.0	3.0	2.0	3.0	2.9	2.0	R1,5,7
9068565	3.0	2.0	7.0	6.0	3.0	5.0	4.0	5.0	4.4	2.0	R2	9068528	3.0	4.0	Dead	3.0	3.0	3.0	2.0	2.0	2.9	2.0	R7,8
9057169	2.0	2.0	6.0	2.0	3.0	2.0	3.0	7.0	3.4	2.0	R1,2,4,6	9068510	6.0	4.0	3.0	3.0	5.0	3.0	3.0	2.0	3.6	2.0	R8
9068510	4.0	4.0	4.0	3.0	4.0	5.0	2.0	2.0	3.5	2.0	R7,8	9068574	3.0	6.0	4.0	4.0	3.0	2.0	3.0	3.0	3.5	2.0	R6
9068574	3.0	3.0	5.0	4.0	2.0	5.0	3.0	3.0	3.5	2.0	R5	9068562	3.0	3.0	5.0	4.0	3.0	4.0	3.0	3.0	3.5	3.0	R1,2,5,7,8
9068525	2.0	3.0	2.0	7.0	Dead	3.0	Dead	2.0	3.2	2.0	R 1,3,8	9068508	4.0	4.0	3.0	3.0	3.0	4.0	3.0	4.0	3.5	3.0	R3,4,5,7
9068507	3.0	Dead	3.0	Dead	Dead	3.0	4.0	4.0	3.4	3.0	R1,3,6	9068565	7.0	3.0	4.0	6.0	3.0	4.0	3.0	Dead	4.3	3.0	R2,5,7
9068528	3.0	3.0	Dead	3.0	3.0	4.0	4.0	4.0	3.4	3.0	R1,2,4,5	9068525	3.0	4.0	3.0	3.0	Dead	3.0	Dead	3.0	3.2	3.0	R1,3,4,6,8
1=No Insect/Disease; 9=Severe Insect/Disease												1=No Insect/Disease; 9=Severe Insect/Disease											

**Study: 29I136J**

**Study Title** - Assembly and Evaluation of Wild Plum, *Prunus americana* Marsh.

**Study Leader:** Henry, J.

**Introduction:**

Wild plum is recognized as an excellent wildlife plant that also has some aesthetic value. It is a shrub or small tree with shaggy bark. Leaves are narrow to wedge-shaped, hairless or nearly so, somewhat long-pointed, sharply and often doubly toothed. Usually no glands are found on leaf-stalks. Twigs are typically hairless. Buds are red-brown, mostly about 1/8 inch in length. Leaf/scars are not abnormally enlarged. Leaves are one to five inches long. Wild plum reaches a height of 15 – 30 feet; with a diameter of five to ten inches. Flowers are white, three - five inch clusters, appearing March - May. Fruits are red and yellow, usually 7/8 - 1 1/4 inches, seed are somewhat flattened and ripen June - October. This species occurs from Massachusetts to Manitoba, New Mexico, Central Texas and southwest Florida.

**Problem:**

There is a lack of an available cultivar of wild plum specifically for this area. A need for developing a local selection or source identified selected sources of wild plum for use as wildlife food and habitat in the three states being served by the center has been identified by NRCS and other conservation and wildlife agencies.

**Objective:**

The objective is to assemble, comparatively evaluate, select and release an adapted cultivar selection of wild plum.

**Discussion**

**1990-1993**

Seed was collected from native stands during 1990, 1991, and 1992. A total of 27 collections were made in Missouri, Iowa, and Illinois. The seed was stratified, germinated in the greenhouse and grown out in open bottom milk-carton type containers. Eighteen of the 27 collections germinated.

**1994-1998**

The plants were transplanted into a randomized complete block with seven replications and one nonrandomized block. The planting was established May 16, 1994 in Field #11e at the PMC. There were several significant dry periods throughout the summer and the plants were under stress several times. The plants were hand watered several times and only four out of 120 plants under evaluation were lost.

The planting was evaluated in 1995, 1996, 1997 and 1998 with very good survival considering the tough establishment year and a very droughty 1998.

The following accessions were selected in 1998 for field plantings: 9062309 (South Dakota), 9057088 (Moultrie County, Illinois), 9068546 (Dallas County, Missouri), 9068545 (Phelps County, Missouri), and 9068580 from Livingston County, Missouri.

## 1999

The 1999 evaluations of this study took place at different times of the year to capture the purposes for the evaluations: height, spread, fruit production, and form.

Table #2 lists the different accessions included in this assembly along with the locations and collectors' names.

Tables # 2, #3, #4, #5 and #6 reflect the plants' performance from 1995 to 1999. These tables can be found in the 1999 Elsberry Technical Report.

## 2000

There were no plant performance evaluations done on this study in year 2000 other than fruit production and insect and disease resistance evaluations. On April 28, 2000 an infestation of the caterpillar tent worm, *Malacosoma americanum* was noted in the planting. A closer observation revealed a severe infestation of the caterpillar tent worm affecting every plant in the assembly. The pesticide Malathion 57 EC liquid was used following the label recommendations. The control was very effective.

Tables #2, #3, #4, #5, and #6 reflect the plants' locations, collectors and performance for years 1995 to 1999.

The following information (Table #1) pertains to the fruit production harvested from selected accessions in year 2000.

**Table #1**

<b>Accession Number</b>	<b>Amount of Clean Seed Produced</b>
<b>9062309</b>	<b>4.9 ounces</b>
<b>9068580</b>	<b>11.5 ounces</b>
<b>9068485</b>	<b>5.5 ounces</b>
<b>9057088</b>	<b>1.7 ounces</b>
<b>9068546</b>	<b>11.0 ounces</b>

**Table #2 – Accessions, Locations and Collector’s Name**

<u>Accession Number</u>	<u>Locations Collected</u>	<u>Collector’s Name</u>
9062309	PMC, Bismarck, North Dakota	Dwight Tober
9057096	Kendall Co., Illinois	William D. Glass
9057085	Coles Co., Illinois	Robert E. Szafoni
9057088	Moultrie Co., Illinois	Robert E. Szafoni
9057130	Grundy Co., Illinois	William D. Glass
9057139	Iroquois Co., Illinois	William D. Glass
9057146	Will Co., Illinois	William D. Glass
9057163	Ogle Co., Illinois	Jim R. Heim
9057164	Woodbury Co., Iowa	Harry A. Minor
9057165	Kankakee Co., Illinois	William D. Glass
9957166	Woodbury Co., Iowa	Harry A. Minor
9068480	Livingston Co, Illinois	William D. Glass
9068485	Ogle Co., Illinois	Jim R. Heim
9057185	Cooper Co., Missouri	David M. Skaer
9867516	Livingston Co., Illinois	Mark Baron
9068515	Moniteau Co., Missouri	Henry E. Knipker
9068514	Grundy Co., Illinois	William D. Glas
9068546	Dallas Co., Missouri	David L. Wright
9068545	Phelps Co., Missouri	Melodie marshall
9068544	Cooper Co., Missouri	Linda Young
9068543	Kendall Co., Illinois	Dayle Saar
9068580	Livingston Co., Missouri	Mac Ellis
9068581	Lincoln Co., Missouri	Bruce Schuette

**2001**

A similar infestation of the caterpillar tent worm, *Malacosoma americanum*, occurred this year as it did last year. The infestation affected all accessions to some degree and was noted encroaching into the planting during the last week of April 2001. The pesticide Malathion 57 EC liquid was used following the label recommendation. The control was again very effective.

Evaluations made this year included insect and disease resistance and fruit production.

The following Table #3 is the summary of plant performance of the selected accessions of wild plum out of the initial assembly.

**2002**

Plant performance evaluations were documented on those accessions selected for field plantings in the PMC service. A light infestation of the caterpillar tent worm, *Malacosoma americanum*, was noted in this assembly in early spring, however no pesticide was applied. No plants were seriously affected as a result of not applying the pesticide Malathoin 57 EC. Table #3 reflects those evaluations. The accessions listed in the following table will be allowed to remain and all the rest of the plants will be removed. The remaining accessions will be allowed to cross and the progeny will be assigned a separate accession number and only one release (Tested Class) will be made for the PMC service area.

**Table #3**

<b>Acc. Number</b>	<b>1995</b>	<b>1996</b>	<b>1997</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>	<b>2001</b>	<b>2002</b>	<b>Average</b>
<b>9062309</b>									
Height (ft)	2.7	4.7	7.1	8.2	9.4				6.4 feet
Spread (ft)	0.8	3.2	6.9	7.7	10.6				
Ins/Dis	4	4	4	4	4	4	4	4	4.6
Form	4	4	4	4	4	4	4	4	4.0
Fruiting			4	4	5	7	6	9	4.3
<b>9068580</b>									
Height (ft)	3.1	6.1	9.0	9.8	10.4				7.7
Spread (ft)	0.93	4.6	9.3	10.0	11.3				7.3
Ins/Dis	2	2	2	3	3	4	4	4	2.9
Form	4	3	3	3	3	3	3	3	3.1
Fruiting			4	3	4	6	4	6	4.5
<b>9068485</b>									
Height (ft)	2.2	3.9	5.5	6.3	9.1				5.4
Spread (ft)	0.93	4.6	9.2	10.5	11.3				
Ins/Dis	2	2	2	2	2	4	4	4	2.6
Form	4	3	3	2	2	2	2	2	2.6
Fruiting			4	3	4				
<b>9068545</b>									
Height (ft)	2.2	3.9	5.5	6.3	7.8				5.4
Spread (ft)	0.3	3.0	5.6	6.8	8.5				4.8
Ins/Dis	3	3	3	3	3	5	4	4	3.4
Form	5	3	3	3	3	3	3	3	3.3
Fruiting			3	4	4	8	1	1	3.5

**Table 3 continued**

<b>Acc. Number</b>	<b>1995</b>	<b>1996</b>	<b>1997</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>	<b>2001</b>	<b>2002</b>	<b>Average</b>
<b>9068546</b>									
Height (ft)	2.9	5.2	7.9	16.6	17.3				10.0
Spread (ft)	0.8	4.2	8.1	8.5	10.9				6.5
Ins/Dis	3	2	2	3	3	4	5	5	3.1
Form	4	4	3	3	3	3	3	3	3.3
Fruiting			3	2	2	6	2	3	2.7

Rating for Insect/Disease: 1 = Exc Resistance, 9 = Poor Resistance

Rating for Fruiting: 1 = Heavy Fruit Production, 9 = Poor Fruit Production

Rating for Form: 1 = Excellent, 9 = Poor

#### **Fruit Production Chart for 2000 - 2002**

**Table #4**

<b>Acc. Number</b>	<b>2000 Pound(s)</b>	<b>2001 Pound(s)</b>	<b>2002 Pound(s)</b>	<b>Average</b>
9062309	0.31	0.60	12 Seeds	0.30
9068580	0.72	1.50	0.70	0.97
9068485	0.11	3.45	3.00	2.19
9068546	0.69	3.57	1.90	2.05
9057088	0.00	5.86	0.40	2.09

Study 291136J Assembly and Evaluation of <i>Prunus Americana</i> , Wild Plum												Table #4											
												Height in Feet											
1995												1996											
Accssion	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Ave.	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Ave.	Best	Location
434240	4.50	5.30	3.80	4.30	2.60	Dead	4.10	-	4.10	<b>5.30</b>	R2	9068545	7.70	6.40	6.80	6.20	5.70	Dead	5.40	-	6.37	<b>7.70</b>	R1
9068580	3.60	5.00	2.60	4.30	1.50	2.00	2.60	3.00	3.08	<b>5.00</b>	R2	434240	7.10	7.30	6.30	6.00	5.10	Dead	6.00	-	6.30	<b>7.30</b>	R2
9057088	4.30	3.10	3.10	4.80	2.50	2.50	2.60	3.50	3.30	<b>4.80</b>	R4	9057096	5.20	7.00	6.20	Dead	Dead	1.30	Dead	-	4.93	<b>7.00</b>	R2
9068545	4.50	3.00	3.00	3.20	2.30	Dead	2.00	-	3.00	<b>4.50</b>	R1	9068514	7.00	6.50	5.10	4.50	Dead	4.10	4.50	-	5.28	<b>7.00</b>	R1
9068546	3.70	4.30	3.60	2.30	2.60	1.80	2.10	2.40	2.85	<b>4.30</b>	R2	9068580	6.90	7.00	6.60	6.80	4.40	4.60	6.00	6.30	6.08	<b>7.00</b>	R2
9068516	2.50	2.00	Dead	4.00	2.00	Dead	Dead	-	2.63	<b>4.00</b>	R4	9068480	4.70	3.10	5.10	6.80	2.90	Dead	Dead	Dead	4.52	<b>6.80</b>	R4
9068515	2.50	0.60	3.80	2.70	1.50	2.50	2.30	2.30	2.28	<b>3.80</b>	R3	9057088	6.50	5.70	5.20	4.60	5.60	5.10	5.40	-	5.44	<b>6.50</b>	R1
9057096	3.60	2.30	1.40	Dead	Dead	1.10	Dead	-	2.10	<b>3.60</b>	R1	9068546	5.50	6.20	6.50	5.60	4.70	3.60	4.40	5.20	5.21	<b>6.50</b>	R3
9068485	3.30	2.00	2.30	2.70	1.50	Dead	1.20	-	2.17	<b>3.30</b>	R1	9062309	6.30	Dead	3.60	4.80	3.80	4.80	Dead	-	4.66	<b>6.30</b>	R1
9068514	3.10	1.90	2.60	2.00	Dead	1.80	2.10	-	2.25	<b>3.10</b>	R1	9057165	5.30	5.00	6.20	6.00	5.10	-	-	-	5.52	<b>6.20</b>	R3
9068480	2.60	3.10	2.40	3.00	1.60	Dead	Dead	Dead	2.54	<b>3.10</b>	R2	9068516	4.90	5.00	Dead	5.10	6.10	Dead	Dead	-	5.28	<b>6.10</b>	R5
9068478	2.60	2.40	3.00	2.80	1.60	2.60	1.40	-	2.34	<b>3.00</b>	R3	9068543	4.20	6.00	5.30	4.70	Dead	Dead	Dead	-	5.05	<b>6.00</b>	R2
9062309	2.80	Dead	2.00	3.00	2.60	2.90	Dead	-	2.66	<b>3.00</b>	R4	9068515	5.10	2.40	5.90	5.30	4.30	4.20	4.10	4.80	4.51	<b>5.90</b>	R3
9057165	1.90	1.80	2.80	2.00	1.40	-	-	-	1.98	<b>2.80</b>	R3	9062308	4.40	5.00	3.10	4.80	Dead	Dead	2.60	Dead	3.98	<b>5.00</b>	R2
9068543	2.40	2.70	2.50	2.00	Dead	Dead	Dead	-	2.40	<b>2.70</b>	R2	9068478	3.10	4.50	3.40	4.50	4.30	4.30	3.40	-	3.93	<b>4.50</b>	R2,4
9062308	2.00	2.20	2.30	1.60	Dead	Dead	1.75	Dead	1.97	<b>2.30</b>	R3	9068485	4.10	4.10	4.00	4.50	4.00	Dead	2.60	-	3.88	<b>4.50</b>	R4
9057146								1.60	1.60	<b>1.60</b>	R8	9057146								4.50	4.50	<b>4.50</b>	R8
ND-286								Dead		<b>0.00</b>		ND-286								Dead		<b>0.00</b>	
1997												1998											
Accssion	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Ave.	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Ave.	Best	Location
9057088	9.50	6.40	7.40	7.30	8.60	7.00	9.00	10.00	8.15	<b>10.00</b>	R8	9068545	12.10	10.90	7.70	10.40	9.60	Dead	7.90	-	9.77	<b>12.10</b>	R1
9068545	11.00	9.80	6.60	9.10	8.00	Dead	7.00	-	8.58	<b>10.00</b>	R1	9068580	11.30	11.00	10.90	11.80	9.00	8.80	9.40	-	10.31	<b>11.30</b>	R1
9068580	10.00	10.00	9.60	10.80	7.20	7.00	8.20		8.97	<b>10.00</b>	R1,2	9057088	10.20	7.70	8.30	8.20	9.60	8.00	7.30	11.20	8.81	<b>11.20</b>	R8
9068546	7.20	9.70	9.00	8.40	7.00	6.00	7.60	8.00	7.86	<b>9.70</b>	R2	434240	10.20	10.00	10.70	8.90	8.60	Dead	8.60	-	9.50	<b>10.70</b>	R3
434240	9.50	9.00	9.50	7.60	7.30	Dead	8.20	-	8.52	<b>9.50</b>	R1,3	9068515	8.90	5.80	10.30	8.10	6.00	7.00	9.90	7.10	7.89	<b>10.30</b>	R3
9068515	8.20	4.20	9.10	7.40	5.00	6.00	8.20	6.20	6.79	<b>9.10</b>	R3	9068480	8.80	6.80	10.20	7.70	7.00	Dead	Dead	6.90	7.90	<b>10.20</b>	R3
9057096	7.30	7.20	8.00	Dead	Dead	2.50	Dead	-	6.25	<b>8.00</b>	R3	9068546	8.70	10.20	10.00	9.90	8.20	67.90	8.20	9.80	16.61	<b>10.20</b>	R2
9062309	8.00	Dead	7.00	7.20	6.40	7.00	Dead	-	7.12	<b>8.00</b>	R1	9057146								8.90	8.90	<b>8.90</b>	R8
9068516	7.80	7.20	Dead	6.00	7.20	Dead	Dead	-	7.05	<b>7.80</b>	R1	9062309	8.90	Dead	8.10	8.40	7.10	8.30	Dead	-	8.16	<b>8.90</b>	R1
9062308	6.40	2.50	5.10	7.60	Dead	Dead	4.00	Dead	5.12	<b>7.60</b>	R4	9068514	8.80	7.30	8.10	7.40	Dead	8.10	7.40	-	7.85	<b>8.80</b>	R1
9068514	7.60	6.40	7.40	6.30	Dead	7.00	6.60	-	6.88	<b>7.60</b>	R1	9057096	7.90	7.70	8.60	Dead	Dead	4.50	Dead	-	7.18	<b>8.60</b>	R3
9068543	6.00	5.00	7.20	7.00	Dead	Dead	Dead	-	6.30	<b>7.20</b>	R3	9068516	8.10	8.60	Dead	7.20	8.30	Dead	Dead	-	8.05	<b>8.60</b>	R2
9057146								7.20	7.20	<b>7.20</b>	R8	9068543	7.00	6.00	8.30	8.10	Dead	Dead	Dead	-	7.35	<b>8.30</b>	R3
9068480	7.00	5.40	9.00	6.30	6.00	Dead	Dead	6.00	6.62	<b>7.00</b>	R1	9062308	7.30	4.90	6.60	8.00	Dead	Dead	5.00	Dead	6.36	<b>8.00</b>	R4
9057165	5.30	5.10	6.10	7.00	5.70	-	-	-	5.84	<b>7.00</b>	R4	9057165	6.60	6.80	7.40	8.00	6.80	-	-	-	7.12	<b>8.00</b>	R4
9068478	3.20	6.50	4.40	6.40	Dead	Dead	4.60	-	5.02	<b>6.80</b>	R6	9068478	4.00	6.90	5.40	7.20	Dead	Dead	5.20	-	5.74	<b>7.20</b>	R4
9068485	5.70	6.30	5.00	6.80	6.30	Dead	2.70	-	5.47	<b>6.80</b>	R4	9068485	6.10	7.20	6.00	7.40	7.10	Dead	3.80	-	6.27	<b>7.20</b>	R2
ND-286								Dead		<b>0.00</b>		ND-286										<b>0.00</b>	
Height measured in feet																							





Study 29I136J Assembly and Evaluation of Prunus Americana, Wild Plum											
Table #5 - continued											
1999											
Spread in Feet											
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Average	Best	Location
9068480	9.00	8.50	10.50	5.00	8.50	0.00	0.00	9.00	8.42	<b>10.50</b>	R3
9057096	8.50	9.50	10.50	0.00	0.00	0.00	0.00	0.00	9.50	<b>10.50</b>	R3
9068478	5.00	8.00	8.00	0.00	8.00	11.50	11.00	0.00	8.58	<b>11.50</b>	R6
9068515	10.00	7.50	7.00	12.00	9.00	11.00	9.00	0.00	9.36	<b>12.00</b>	R4
9062308	9.00	0.00	7.00	10.00	0.00	0.00	10.00	0.00	9.00	<b>10.00</b>	R4
9068485	8.30	8.50	7.00	12.00	10.00	0.00	5.00	0.00	8.47	<b>12.00</b>	R4
9057088	12.50	10.00	12.50	11.00	14.00	9.00	13.00	13.00	11.88	<b>14.00</b>	R5
9068545	14.50	12.00	0.00	11.00	14.00	0.00	0.00	0.00	12.88	<b>14.50</b>	R1
9068543	8.50	10.00	12.00	10.00	0.00	0.00	0.00	0.00	10.13	<b>12.00</b>	R3
9068516	10.00	11.00	0.00	11.00	11.00	0.00	9.00	0.00	10.40	<b>11.00</b>	R2,4,5
9068514	9.50	9.00	9.00	9.00	0.00	10.00	8.00	0.00	9.08	<b>10.00</b>	R6
9068580	11.00	10.00	0.00	13.00	11.00	11.00	10.00	13.00	11.29	<b>13.00</b>	R4,8
9057146								10.00	10.00	<b>10.00</b>	R8
9068546	10.00	5.00	13.00	11.50	11.00	12.00	14.00	11.00	10.94	<b>14.00</b>	R7
434240	0.00	11.00	11.00	11.00	0.00	0.00	0.00	0.00	0.00	<b>11.00</b>	R2,3,4
ND-286								0.00	0.00	<b>0.00</b>	
9062309	11.50	0.00	8.50	11.00	11.00	11.00	0.00	0.00	10.60	<b>11.50</b>	R1
9057165	8.00	9.00	11.00	10.00	8.50	0.00	0.00	0.00	9.30	<b>11.00</b>	R3
Spread Measured in Feet											
0 = Dead plant											

Study 291136J Assembly and Evaluation of <i>Prunus Americana</i> , Wild Plum																								Table #6
												Form												
1995												1996												
Accssion	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Ave.	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Ave.	Best	Location	
ND-286								Dead	-	0.00		ND-286								Dead	-	0.00		
434240	1.00	2.00	2.00	2.00	4.00	Dead	5.00	-	2.67	1.00	R1	9068478	3.00	4.00	2.00	6.00	6.00	4.00	3.00	-	4.00	2.00	R3	
9057088	2.00	4.00	5.00	4.00	4.00	4.00	2.00	4.00	3.63	2.00	R1, 7	9068515	2.00	5.00	3.00	3.00	4.00	6.00	3.00	6.00	4.00	2.00	R1	
9068545	2.00	3.00	3.00	5.00	2.00	Dead	6.00	-	3.50	2.00	R1, 5	9068514	2.00	5.00	4.00	5.00	Dead	5.00	8.00	-	4.83	2.00	R1	
9068516	2.00	8.00	Dead	7.00	2.00	Dead	Dead	-	4.75	2.00	R1, 5	9068546	2.00	6.00	2.00	3.00	4.00	5.00	7.00	3.00	4.00	2.00	R1, 3	
9068478	4.00	7.00	4.00	3.00	5.00	4.00	4.00	-	4.43	3.00	R3	9068480	8.00	4.00	5.00	3.00	6.00	Dead	Dead	Dead	5.20	3.00	R4	
9068515	4.00	6.00	5.00	5.00	7.00	3.00	5.00	6.00	5.13	3.00	R6	9057096	4.00	3.00	3.00	Dead	Dead	6.00	Dead	-	4.00	3.00	R2, 3	
9062308	5.00	3.00	6.00	6.00	Dead	Dead	6.00	Dead	5.20	3.00	R2	9062308	3.00	5.00	3.00	5.00	Dead	Dead	6.00	Dead	4.40	3.00	R1, 3	
9068580	5.00	3.00	5.00	3.00	5.00	5.00	5.00	5.00	4.50	3.00	R2, 4	9068485	5.00	3.00	3.00	3.00	4.00	Dead	3.00	-	3.50	3.00	R2,3,4,7	
9068546	4.00	5.00	3.00	5.00	7.00	5.00	5.00	5.00	4.88	3.00	R3	9057088	3.00	6.00	4.00	6.00	4.00	4.00	3.00	4.00	4.25	3.00	R1, 7	
9068480	4.00	8.00	5.00	7.00	6.00	Dead	Dead	Dead	6.00	4.00	R1	9068545	5.00	4.00	3.00	5.00	4.00	Dead	7.00	-	4.67	3.00	R3	
9068514	4.00	7.00	7.00	8.00	Dead	4.00	5.00	-	5.83	4.00	R1, 6	9068516	4.00	4.00	Dead	5.00	3.00	Dead	5.00	-	4.20	3.00	R5	
9057165	4.00	5.00	8.00	8.00	8.00	-	-	-	6.60	4.00	R1	9068580	5.00	5.00	3.00	3.00	3.00	3.00	4.00	3.00	3.63	3.00	R3,4,5,6,8	
9068485	7.00	7.00	8.00	7.00	5.00	Dead	8.00	-	7.00	5.00	R5	9057146								3.00	3.00	3.00	R8	
9068543	5.00	8.00	5.00	8.00	Dead	Dead	Dead	-	6.50	5.00	R1, 3	434240	3.00	3.00	4.00	7.00	4.00	Dead	3.00	-	4.00	3.00	R1,2, 7	
9062309	5.00	Dead	6.00	6.00	6.00	7.00	Dead	-	6.00	5.00	R1	9062309	3.00	Dead	5.00	3.00	4.00	4.00	Dead	-	3.80	3.00	R1, 4	
9057096	6.00	7.00	6.00	Dead	Dead	8.00	Dead	-	6.75	6.00	R1, 3	9068543	5.00	4.00	5.00	4.00	Dead	Dead	Dead	-	4.50	4.00	R2, 4	
9057146								7.00	7.00	7.00	R8	9057165	5.00	4.00	5.00	5.00	6.00	-	-	-	5.00	4.00	R2	
1997												Form	1998											
Accssion	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Ave.	Best	Location	Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Ave.	Best	Location	
ND-286								Dead		0.00		ND-286								Dead	-	0.00		
9068545	1.00	3.00	8.00	7.00	7.00	Dead	5.00	-	5.17	1.00	R1	9057088	1.00	6.00	5.00	7.00	5.00	Dead	4.00	-	4.67	1.00	R1	
9068580	1.00	3.00	7.00	2.00	5.00	6.00	2.00	2.00	3.50	1.00	R1	9068580	1.00	3.00	5.00	2.00	5.00	5.00	2.00	2.00	3.13	1.00	R1	
434240	1.00	5.00	6.00	8.00	5.00	Dead	3.00	-	4.67	1.00	R1	434240	1.00	4.00	6.00	8.00	4.00	Dead	3.00	-	4.33	1.00	R1	
9057088	1.00	7.00	6.00	8.00	5.00	4.00	3.00	2.00	4.50	2.00	R8	9068545	1.00	2.00	6.00	5.00	6.00	3.00	3.00	-	3.71	2.00	R8	
9068546	5.00	3.00	2.00	2.00	5.00	5.00	3.00	5.00	3.75	2.00	R3,4	9068514	6.00	8.00	5.00	Dead	6.00	7.00	2.00	-	5.67	2.00	R7	
9068515	3.00	6.00	5.00	5.00	7.00	5.00	3.00	5.00	4.88	3.00	R1, 7	9068546	5.00	3.00	2.00	2.00	4.00	4.00	3.00	4.00	3.38	2.00	R3,4,8	
9068516	3.00	7.00	Dead	8.00	5.00	Dead	4.00	-	5.40	3.00	R1	9068515	3.00	5.00	4.00	4.00	7.00	5.00	3.00	4.00	4.38	3.00	R1,7	
9068514	6.00	8.00	5.00	Dead	6.00	8.00	3.00	-	6.00	3.00	R1	9068516	3.00	6.00	Dead	8.00	5.00	Dead	4.00	-	5.20	3.00	R1	
9068480	4.00	5.00	8.00	5.00	6.00	Dead	3.00	6.00	5.29	4.00	R1	9068480	4.00	6.00	7.00	4.00	6.00	Dead	3.00	6.00	5.14	4.00	R1,4	
9062308	4.00	9.00	7.00	8.00	Dead	Dead	7.00	-	5.83	4.00	R4	9068478	8.00	6.00	7.00	6.00	Dead	4.00	6.00	-	6.17	4.00	R6	
9057096	6.00	7.00	7.00	5.00	Dead	8.00	Dead	-	6.60	5.00	R4	9062308	4.00	8.00	7.00	8.00	Dead	Dead	7.00	-	6.80	4.00	R1	
9068478	8.00	6.00	7.00	7.00	Dead	5.00	6.00	-	6.50	5.00	R6	9057096	5.00	6.00	6.00	5.00	Dead	8.00	Dead	-	6.00	5.00	R1,4	
9068485	6.00	6.00	6.00	7.00	5.00	Dead	6.00	-	6.00	5.00	R5	9068485	6.00	6.00	5.00	6.00	5.00	Dead	6.00	-	5.67	5.00	R3,5	
9068543	6.00	7.00	5.00	5.00	Dead	Dead	Dead	-	5.75	5.00	R3,4	9068543	6.00	6.00	5.00	5.00	Dead	Dead	Dead	-	5.50	5.00	R3,4	
9057146								5.00	5.00	5.00	R8	9057146								5.00	5.00	5.00	R8	
9062309	5.00	Dead	6.00	5.00	8.00	6.00	Dead	-	6.00	5.00	R1,4	9062309	5.00	Dead	5.00	5.00	7.00	6.00	Dead	-	5.60	5.00	R1,3,4	
9057165	7.00	7.00	6.00	6.00	6.00	-	-	-	6.40	6.00	R4,5,6	9057165	7.00	6.00	6.00	5.00	6.00	-	-	-	6.00	5.00	R4	

Rating: 1= Excellent, 9=Poor 0=Dead Plant





Study 291136J Assembly and Evaluation of <i>Prunus Americana</i> , Wild Plum											Table #8
Insect/Disease Resistance											
1999											
Accession	Rep 1	Rep 2	Rep 3	Rep 4	Rep 5	Rep 6	Rep 7	Rep 8	Ave.	Best	Location
ND-286	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	<b>0.00</b>	
9068480	4.00	4.50	4.50	1.00	4.50	0.00	0.00	1.50	3.33	<b>1.00</b>	R4, 8
9068478	4.00	2.50	1.50	0.00	1.00	1.00	1.00	0.00	1.69	<b>1.00</b>	R3,5,6,7
9068515	2.50	3.50	5.50	1.00	4.50	3.00	3.00	0.00	3.29	<b>1.00</b>	R4
9057088	1.50	2.00	1.50	1.00	1.50	1.00	1.00	1.00	1.31	<b>1.00</b>	R1,3,4,5,6,7,8
9068545	2.00	1.00	0.00	2.00	1.50	0.00	0.00	0.00	1.63	<b>1.00</b>	R1,2,5
9068580	2.50	1.00	0.00	1.00	3.00	1.00	1.50	1.50	1.64	<b>1.00</b>	R2,4,6,7,8
9068546	6.00	1.00	1.00	1.00	1.00	1.00	1.00	1.50	1.69	<b>1.00</b>	R2,7,8
9057096	5.50	3.00	1.50	0.00	0.00	0.00	0.00	0.00	3.33	<b>1.50</b>	R3
9062308	6.00	0.00	3.50	1.50	0.00	0.00	2.00	1.50	2.90	<b>1.50</b>	R4,7,8
9068485	5.50	5.50	4.00	1.50	1.50	0.00	3.00	0.00	3.50	<b>1.50</b>	R4,5
9068516	2.50	4.00	0.00	2.50	3.50	0.00	2.00	0.50	2.73	<b>1.50</b>	R4,7,8
9057146	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.50	1.50	<b>1.50</b>	R8
9062309	3.50	0.00	3.50	1.50	3.50	2.50	0.00	0.00	2.90	<b>1.50</b>	R4,6,7,8
9068543	2.50	4.50	2.00	2.50	0.00	0.00	0.00	0.00	2.88	<b>2.00</b>	R1,3,4
9068514	3.00	3.00	2.00	2.00	0.00	2.50	2.00	0.00	2.42	<b>2.00</b>	R3,4,6,7
9057165	6.00	4.50	2.50	3.00	3.00	0.00	0.00	0.00	3.80	<b>2.00</b>	R3,5
434240	0.00	5.50	5.00	0.00	0.00	0.00	0.00	0.00	5.25	<b>3.00</b>	R2,5
Disease Resistance Rating:				1-Excellent resistance							
				9-Poor resistance							
				0=Dead plant							

**Study: 29A1370**

**Study Title:** Wetland/Riparian Propagation, Establishment, and Demonstration

**Study Leader:** Henry, J.

**Introduction:**

There is a growing interest in wetland restoration throughout the conservation community. Government programs, such as USDA-Wetland Reserve Program, the USFWS Partners for Wildlife, Wetland Restoration Program, the Missouri Department of Conservation (MDC) Private Lands Wetland Program, and private programs sponsored by Ducks Unlimited and Waterfowl USA have all focused on the need for a suitable supply of plants in wetland restoration efforts.

The increasing use of wetlands as filters in agricultural waste management and the control of non-point source pollution also indicates the need for a greater knowledge base for proper plant selection.

Understanding wetland ecosystems will require improved and increased quality of information on wetland plants and ecosystems. Innovative approaches to field management and additional training of personnel in wetland conservation and management will also be needed. Intra- and interagency coordination and information exchange among state and federal agencies will help standardize monitoring and management strategies.

**Problem:**

Information is largely unavailable related to the propagation, adaptation, and potential use of many of the wetland species found in the Midwest. Wetland plants of interest often have multi-use potential providing wildlife benefits, shoreline stabilization, water quality improvement, and/or aesthetic benefits. They are also needed to fulfill conservation needs resulting from increased demands in wetland development and water treatment. The ability to document this information or to observe the interaction of selected species is restricted by the availability of plants and plant communities especially under controlled conditions. Proper use of species to address conservation problems is limited by specific knowledge and technology for using these plants.

**Objectives:**

The objectives of the Elsberry PMC wetland study are:

1. Provide a demonstration of various plant materials for wetland conservation and aesthetic values.
2. Provide an area for interagency research on the biology of selected wetland plants.

**Discussion:**

## 1994 – 1999

A large wetland was constructed in Field #4 on the Plant Materials Center in July 1994. Selected plant materials were planted with the intent of evaluating these plants for flood tolerance. The PMC has been working with a flood tolerant switchgrass since 1991. As a result, it was placed in this wetland for further testing along with six accessions of eastern gamagrass which were found growing in wet conditions: accessions 9078842, 9078844 and 9078843 were collected in Atchison County, Missouri, 9078845 collected in Holt County, Missouri, 9078840 collected in Chariton County, Missouri and 9078846 was collected in Clinton County, Missouri. Local collections of bermudagrass and swamp milkweed were planted in the spring of 1998. Two collections of prairie cordgrass (Cuivre Island and Lost Creek) were also planted in this wetland. The switchgrass, eastern gamagrass and the prairie cordgrass were planted in 1997. All plants in this wetland were given time to establish prior to the beginning of the flooding operation which took place in October 1999. The wetland was flooded to a depth of 40 inches. This water remained in the wetland until early spring of 2000. Once the water is drained out of the wetland and enough time elapses for plant regrowth, evaluations on survival will take place.

The following Tables #1, #2, and #3 reflect the plants' performance.

## 2000

Water was drained out of the wetland in segments because the drainpipe was not functioning properly. This operation started on March 21, 2000 and ended on March 30, 2000. The prairie cordgrass were the first plants to begin green up (March 30) followed by the bermudagrass planting. 'Cave-In-Rock' switchgrass sod (23 plugs) was planted on the west side of the flood tolerant switchgrass (sod) for comparison with other plant species in the wetland. On June 1, 2000, flood tolerant switchgrass was seeded in a plot 50 feet long and three feet wide. On August 9 an evaluation of the seeded flood tolerant switchgrass revealed no germination had taken place in the plot seeded on June 1. Poor germination has been experienced with this selection since 1998. There was no flooding of the wetland this fall to allow the Cave-In-Rock to get fully established. The following is a listing of percent survival of plants included in this study. The best performing plants in this study are Cuivre Island and Lost Creek collection of *Spartina pectinata*, *Tripsacum dactyloides* accessions 9078843, 9078845, and 'Pete'; and *Cynodon dactylon*. The following tables reflect the different plants' performance before and after a flooding event.

## 2001

Began pumping water into wetland on April 24, 2001. The objective of the pumping was to flood the wetland to a depth of approximately 32 inches of water, which was achieved by April 27, 2001. The water was allowed to remain in the wetland for seven days. Water was then allowed to drained out of the wetland starting on April 30, 2001. All the water was drained out of the wetland by May 1, 2001. On May 8 evaluations were conducted to document re-growth after flooding. Again on June 11 a quick flooding scenario was conducted in the wetland to simulate a flash flooding event. Thirty-four inches of water was pumped into the wetland. The PMC began draining the

water out of the wetland on June 15. The process of draining the water out of the wetland was completed on June 19.

The following is a listing of plant vigor ratings for each accession/variety included in this study. Plant vigor evaluations were taken on June 21 and 26, 2001.

## 2002

Plant performance evaluations were performed on April 24, 2002. The wetland was not burned this year as in previous years. Flooding of the wetland began on April 29, 2002. A total of 45 inches of water were pumped into the wetland before the de-watering process began on May 15, 2002. All water was drained out of the wetland by May 17, 2002. The plants were under water for 17 days. Once all the water was drained out of the wetland, follow-up evaluations took place on June 17, 2002. The following table (#1A) reflects the plant performances during 2002 before and after the flood event. Previous years plant performances can be found on Tables #1-#3.

Table #1A

Genus/Species	Common Name	Accession No.	Vigor Rating		Date of Rating	
			BFE	AFE	BFE	AFE
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9098840	Good	Exc.	4/24/02	6/17/02
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078844	Fair	Exc.	4/24/02	6/17/02
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078842	Fair	Exc.	4/24/02	6/17/02
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078846	Good	Good	4/24/02	6/17/02
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078843	Fair	Good	4/24/02	6/17/02
<i>Tripsacum dactyloides</i>	Eastern gamagrass	9078845	Good	Exc.	4/24/02	6/17/02
<i>Tripsacum dactyloides</i>	Pete EGG		Exc.	Exc.	4/24/02	6/17/02
<i>Panicum virgatum</i>	Switchgrass	9062193	Good	Exc.	4/24/02	6/17/02
<i>Panicum virgatum</i>	Switchgrass	9062235	Good	Good	4/24/02	6/17/02
<i>Panicum virgatum</i>	Switchgrass	9062213	Good	Exc.	4/24/02	6/17/02
<i>Panicum virgatum</i>	Switchgrass	C-I-R	Good	Good	4/24/02	6/17/02
<i>Panicum virgatum</i>	Switchgrass	Flood-Tolerant	Good	Good	4/24/02	6/17/02
<i>Spartina pectinata</i>	Prairie cordgrass	Cuivre Island	Exc.	Exc.	4/24/02	6/17/02
<i>Spartina pectinata</i>	Prairie cordgrass	Lost Creek	Exc.	Exc.	4/24/02	6/17/02
<i>Cynodon dactylon</i>	Bermuda grass	Elsberry	Good	Good	4/24/02	6/17/02
<i>Asclepias incarnata</i>	Swamp milkweed	Iowa		Good	4/24/02	6/17/02
<i>Lobelia cardinalis</i>	Cardinal flower	Forrest Keeling	Fair	Good	4/24/02	6/17/02
<i>Carex scoparia</i>	Broomsedge	MDC		Poor	4/14/02	6/17/02
<i>Elymus virginicus</i>	Cuivre River	Forrest Keeling	Exc.	Good	4/14/02	6/17/02

**BFE = Before Flood Event**

**AFE = After Flood Event**

Study 29A1370 - Wetland Species in Wetland at Elsberry PMC							Table #1
<b>Plugs Planted 5-2-97 (Eastern Gamagrass)</b>							
<b>2002 Data</b>	<b>Began Flooding on 4/24/02</b>						
	<b>Total #</b>	<b>Active</b>	<b>Weed</b>	<b>Disease/</b>	<b>Developed</b>		
	<b>Planted</b>	<b>Growing</b>	<b>Comp.</b>	<b>Insect</b>	<b>Seed Head</b>	<b>Vigor</b>	<b>Ave. Ht.</b>
<b>Eastern Gamagrass 9078840 Chariton, Missouri. 5' spacing, planted 5/2/97.</b>							
						<b>25 plants planted</b>	
<b>Dates Evaluated</b>							
7/9/98	20	20	severe	moderate	yes	good	2'5"
9/29/99	20	20	moderate	light rust	yes	good/exc	3'5"
5/11/00	19	17	moderate	moderate	none	poor	6"
9/19/00	13	13	mod/sev	light rust	none	good	2' 5"
6/26/01	20	20	light	none	yes	good	40"
<b>4/24/02 (BFE)</b>	<b>18</b>	<b>18</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>8"</b>
<b>6/17/02 (AFE)</b>	<b>15</b>	<b>15</b>	<b>light</b>	<b>none</b>	<b>yes</b>	<b>exc</b>	<b>24"</b>
<b>Percent surviving as of 6/17/02 was 75%</b>							
<b>Eastern Gamagrass 9078844 Atchison, Missouri. 7' spacing, planted 5/2/97.</b>							
						<b>18 plants planted</b>	
<b>Dates Evaluated</b>							
7/9/98	12	12	severe	moderate rust	yes	poor	2'5"
9/29/99	12	12	moderate	moderate rust	yes	fair	2'5"
5/11/00	12	10	moderate	moderate	none	poor	6"
9/19/00	12	13	severe	light rust	Yes	fair	2' 0"
6/26/01	12	9	light	light rust	yes	fair	34"
<b>4/24/02 (BFE)</b>	<b>9</b>	<b>9</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>fair</b>	<b>7"</b>
<b>6/17/02 (AFE)</b>	<b>9</b>	<b>9</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>exc.</b>	<b>24"</b>
<b>Percent surviving as of 6/17/02 was 75%</b>							
<b>Eastern Gamagrass 9078842 Atchison, Missouri. 15' spacing, planted 5/2/97.</b>							
						<b>9 plants planted</b>	
<b>Dates Evaluated</b>							
7/9/98	5	5	severe	none	yes	fair	2'0"
9/29/99	5	5	severe	none	yes	fair	2'5"
5/11/00	5	3		none	0	poor	6"
9/19/00	5	4	severe	none	none	fair	20"
6/26/01	3	3	light	none	yes	fair	26"
<b>4/24/02 (BFE)</b>	<b>4</b>	<b>4</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>fair</b>	<b>7"</b>
<b>6/17/02 (AFE)</b>	<b>4</b>	<b>4</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>exc.</b>	<b>24"</b>
<b>Percent surviving as of 6/17/02 was 44%</b>							
Rating for Vigor: 1=Excellent; 9=Poor							
Rating for Weed Competition and Dis/Insect: 1=Excellent; 9=Severe							

Study 29A1370 - Wetland Species in Wetland at Elsberry PMC						Table #1-continued	
	<b>Total #</b>	<b>Active</b>	<b>Weed</b>	<b>Disease/</b>	<b>Developed</b>		
	<b>Planted</b>	<b>Growing</b>	<b>Comp.</b>	<b>Insect</b>	<b>Seed Head</b>	<b>Vigor</b>	<b>Ave. Ht.</b>
<b>Eastern Gamagrass 9078846 Clinton, Missouri. 8' spacing, total planted 5/2/97.</b>							
						<b>16 plants planted</b>	
<b>Dates Evaluated</b>							
7/9/98	11	11	severe	none	yes	good	2'0"
9/29/99	11	11	moderate	none	yes	good	2'5"
5/11/00	8	8	moderate	none	none	poor	7"
9/19/00	10	10	severe	light rust	none	fair	2' 0"
6/26/01	8	8	light	light rust	yes	good	38"
<b>4/24/02 (BFE)</b>	<b>10</b>	<b>10</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>8"</b>
<b>6/17/02 (AFE)</b>	<b>10</b>	<b>10</b>	<b>light</b>	<b>none</b>	<b>yes</b>	<b>exc.</b>	<b>30"</b>
<b>Percent surviving as of 6/17/02 was 63%</b>							
<b>Eastern Gamagrass 9078843 Atchison, Missouri. 15' spacing, planted 5/2/97.</b>							
						<b>9 plants planted</b>	
<b>Dates Evaluated</b>							
7/9/98	13	13	severe	none	yes	poor	2'5"
9/29/99	13	13	moderate	none	yes	moderate	3'0"
5/11/00	5	5		none	none	poor	7"
9/19/00	10	10	severe	slight rust	none	fair	2' 0"
6/26/01	4	4	light	light	none	fair	30"
<b>4/24/02 (BFE)</b>	<b>4</b>	<b>4</b>	<b>light</b>	<b>light</b>	<b>none</b>	<b>fair</b>	<b>8"</b>
<b>6/17/02 (AFE)</b>	<b>4</b>	<b>4</b>	<b>light</b>	<b>light</b>	<b>none</b>	<b>good</b>	<b>24"</b>
<b>Percent surviving as of 6/17/02 was 44%</b>							
<b>Eastern Gamagrass 9078845 Holt, Missouri. 8' spacing, planted 5/2/97.</b>							
						<b>16 plants planted</b>	
<b>Dates Evaluated</b>							
7/9/98	12	12	severe	none	yes	good	3'5"
9/29/99	12	12	severe	none	yes	good	3'0"
5/22/00	12	9	severe	none	none		8"
9/19/00	16	16	severe	slight rust	yes	good	2' 5"
6/26/01	10	10	light	none	yes	good	38"
<b>4/24/02 (BFE)</b>	<b>10</b>	<b>10</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>8"</b>
<b>6/17/02 (AFE)</b>	<b>10</b>	<b>10</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>exc.</b>	<b>30"</b>
<b>Percent surviving as of 6/17/02 was 63%</b>							
Rating for Vigor: 1=Excellent; 9=Poor							
Rating for Weed Competition and Dis/Insect: 1=Excellent; 9=Severe							
BFE - Before Flooding Event							
AFE - After Flooding Event							



Study 29A1370 - Wetland Species in Wetland at Elsberry PMC							Table #2
Plugs Planted 6-24-97 (Flood Tolerant Switchgrass)							
2002 Data: Flood Event from 4/29/02 to 5/17/02							
	% Cover/ Plant #	Active Growing	Weed Comp.	Disease/ Insect	Developed Seed Head	Vigor	Ave. Ht.
<b>Switchgrass 9062213 3' spacing, 41 total planted (plugs) 6/24/97.</b>							
<b>Dates Evaluated</b>							
7/9/98		35 plants	moderate	none	all plants	poor/fair	2'.0"
9/29/99		35 plants	moderate	none	all plants	fair	2'.5"
4/26/00		35 plants	moderate	none	none	exc.	5" regrowth
9/19/00	85% row	35 plants	moderate	none	all plants	exc.	4'.5"
6/26/01		33 plants	light	none	none	exc.	40"
<b>4/24/02 (BFE)</b>		<b>31 plants</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>6"</b>
<b>6/17/02 (AFE)</b>		<b>31 plants</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>exc.</b>	<b>30"</b>
<b>Percent surviving as of 6/17/02 was 76%</b>							
<b>Switchgrass 9062235 4' spacing, 31 total planted (plugs) 6/24/97.</b>							
<b>Dates Evaluated</b>							
7/9/98		22 plants	moderate	none	all plants	poor/fair	5'.5"
9/29/99		22 plants	moderate	none	all plants	fair	5'.0"
4/26/00		26 plants	moderate	none	none	exc.	6'.5"
9/19/00		26 plants	moderate	none	All plants	exc.	4' 5"
6/26/01		24 plants	light	none	none	exc.	35"
<b>4/24/02 (BFE)</b>		<b>20 plants</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>6"</b>
<b>6/17/02 (AFE)</b>		<b>20 plants</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>24"</b>
<b>Percent surviving as of 6/17/02 was 65%</b>							
<b>Switchgrass 9062193 5' spacing; 25 total planted (plugs) 6/24/97.</b>							
<b>Dates Evaluated</b>							
7/9/98		17 plants	moderate	none	all plants	fair	3'.5"
9/29/99		17 plants	moderate	none	all plants	good	4'.5"
4/26/00		21 plants	moderate	none	all plants	exc.	6'.5"
9/19/00		21 plants	moderate	none	all plants	exc.	5'.0"
6/26/01		20 plants	light	none	none	exc.	42"
<b>4/24/02 (BFE)</b>		<b>16 plants</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>5"</b>
<b>6/17/02 (AFE)</b>		<b>14 plants</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>exc.</b>	<b>30"</b>
<b>Percent surviving as of 6/17/02 was 56%</b>							
<b>BFE - Before Flooding Event</b>							
<b>AFE - After Flooding Event</b>							

Study 29A1370 - Wetland Species in Wetland at Elsberry PMC						Table #2 - continued		
	% Cover/ Plant #	Active Growing	Weed Comp.	Disease/ Insect	Developed Seed Head	Vigor	Ave. Ht.	
Evaluation Dates:		4/24/02 & 6/17/02						
<b>Cave-In-Rock Switchgrass 23 plants planted.</b>								
<b>Dates Evaluation</b>								
4/18/00	23	23	severe	none	none	good	5"	
9/19/00	9	growing weak	severe	none	yes	poor	2'. 0"	
6/21/01	21	21	light	light	none	good	30"	
<b>4/24/02 (BFE)</b>	<b>10</b>	<b>10</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>8"</b>	
<b>6/17/02 (AFE)</b>	<b>12</b>	<b>12</b>	<b>light</b>	<b>light</b>	<b>none</b>	<b>good</b>	<b>30"</b>	
Percent surviving as of 6/17/02 was 52%								
<b>Flood Tolerant Switchgrass, seeded 50' row plus 3' wide.</b>								
<b>Dates Evaluated</b>								
Seeded 6/1/00 50' x 40" plot - .0038 ac. Rate 6# PLS/ac.								
9/19/00	15%- 20% of 50' row	fair	moderate	none	6/5 5%	good	8"	
6/21/01	22	22	light	none	none	exc.	36"	
<b>4/24/02 (BFE)</b>	<b>16</b>	<b>16</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>5"</b>	
<b>6/17/02 (AFE)</b>	<b>33</b>	<b>33</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>18"</b>	
<b>Flood tolerant switchgrass plugs block, 63 plants planted 5/25/99.</b>								
<b>Dates Evaluated</b>								
4/26/00	92%	58 plants	none	none	6/5 100%	exc.	6'5"	
9/19/00	95%	95%	none	none	6/5 100%	exc.	4' 5"	
6/21/01	80%	66 plants	light	none	none	exc.	36"	
<b>4/24/02 (BFE)</b>	<b>85%</b>	<b>66 plants</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>6"</b>	
<b>6/17/02 (AFE)</b>	<b>85%</b>	<b>66 plants</b>	<b>light</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>24"</b>	
<b>Bermudagrass block plugs, planted 5/25/99.</b>								
<b>Dates Evaluated</b>								
9/28/99	35%	100%	light	none	50%	exc.	3"	
4/26/00		100%	light	none	none	exc.	3-5"	
9/19/00	100%	100%	light	none	100%	exc.	9"	
6/21/01	100%	100%	none	none	none	exc.	6"	
<b>4/24/02 (BFE)</b>	<b>100%</b>	<b>50%</b>	<b>none</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>2"</b>	
<b>6/17/02 (AFE)</b>	<b>90%</b>	<b>90%</b>	<b>none</b>	<b>none</b>	<b>none</b>	<b>good</b>	<b>3"</b>	



Study 29A1370 - Wetland Species in Wetland at Elsberry PMC							Table #3		
Prairie Cordgrass									
2002 Data: Flood Event from 4/29/02 to 5/17/02									
		Active					Ave. Ht.	Average	
	Total #	Growing	Weed	Disease/	Developed		Seed	Forage	
	Planted	Spreading	Comp.	Insect	Seed Head	Vigor	Head	Height	
Prairie Cordgrass Collection, planted 9/29/97							10' x 10'		
East →							3   2   1		
							6   5   4		
							9   8   7		
7/9/98	9	6" average	severe	none	NA	exc.	-	-	
8/1/99	9	30" average	moderate	none	9/9	good	-	-	
9/19/00	9	4.5" ave.	none	none	9/9	exc.	6'.5"	5'.0 forage	
6/21/01	9	6'	light	none	none	exc.	6'	45"	
4/24/02 (BFE)	9	7.5'	light	none	none	exc.	none	17"	
6/17/02 (AFE)	9	8'	light	none	none	exc.	none	36"	
Percent surviving as of 6/17/02 was 100%									
Cuivre Island Prairie Cordgrass Collection, planted 5/15/98							3' x 3'		
					North		4   3   2   1		
							8   7   6   5		
7/9/98	8	5.5"	severe	none	6 plants	good/exc.	4.0"	4.0"	
5/25/99	8	1.5" each	moderate	none	none	exc.	none		
		direction							
Lost Creek Prairie Cordgrass Collection, planted 5/15/98							3'x3'		
							12 11 10 9		
							16 15 14 13		
7/9/98	8	6"	severe	none	4 plants	good/exc.	4.0"	4.0"	
5/25/99	8	1.5" each	moderate	none	none	exc.	none		
		direction							
9/19/00									
Total block for both collections			none	none	35%	exc.	6' 0"	5' 0"	
									More lodging Cuivre Island collection
9/19/00									
14' x 13'5" total spread of blocks			none	none	35%	exc.	6'.0"	More lodging Cuivre Island collection	
9/19/00									
3' x 3' block is filled in total									
prairie cordgrass			none	none	35%	exc.	6'.0"	More lodging Cuivre Island collection	
6/26/01		solid	none	none	none	exc.	6',0"	50"	
4/24/02 (BFE)		80%	none	none	*	exc,	15"		
6/17/02 (AFE)		solid block	none	none	none	exc.	48"		
BFE - Before Flooding Event									
AFE - After Flooding Event									

**Study: 29I141G**

**Study Title:** Assembly and Evaluation of Little Bluestem, *Schizachyrium scoparium*, Nichx.

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

Little bluestem is a native warm season prairie grass. It was a major component making up as much as 50 percent of the tall grass prairie that was native to much of the Elsberry PMC service area. It can also be a major component of glade areas and mixed grass prairies. Little bluestem can be found in prairies, open woods, dry hills, and fields, from Quebec and Maine to Alberta and Idaho, south to Florida and Arizona.

**Problem:**

There are no current varieties of little bluestem on the market that have an origin within the three-state service area. Available varieties do not always perform as well as expected. There is a need for an adapted and improved variety of little bluestem for pasture and range seedings, surface mine reclamation, critical area planting, wildlife plantings, recreational area development and other conservation uses in Missouri, Iowa, and Illinois.

**Objective:**

The objective is to assemble, evaluate, develop and cooperatively release an adapted variety and/or varieties of tested class of little bluestem for conservation use in Missouri, Iowa, and Illinois.

**Procedure:**

Vegetative material from native ecotypes was collected throughout the states of Missouri, Iowa, and Illinois. A minimum of three collections per Major Land Resource Area/state was requested. (Approximately 60 collections total.) Field selection of collected plant material was based on forage quantity and plant vigor.

Each collection (accession) was one individual plant. A collection was made up of more than one plant if they were in the same immediate area (within five feet) and appeared to be clones of each other.

**Discussion:****1996**

The study was approved in July 1996. Collection instructions were sent out and plants were dug in October and November. The samples were picked up shortly after collection and stored in the packing shed at the Plant Materials Center. At this time we received 113 collections from the three-state area. There are a few additional collections expected.

## **1997 - 1998**

The collections were vegetatively propagated in containers in January and grown out in the greenhouse until April. These plants were then transplanted in Field #1 on the PMC from April 22-24, 1997 in a randomized complete block with four replications (see Table #2 for map of plot layout). Thirteen additional collections were made in the summer of 1997 and planted into the replications August 14-15, 1997. This brought the total accessions represented to 130: 79 from Missouri, 20 from Illinois, 27 from Iowa, and four standards of comparison. A list of collectors can be seen in Table #1. First year evaluation consisted of survival. The second year evaluations consisted of survival, height, late dormancy, and form.

## **1999**

The assembly was evaluated in 1999 for forage amount and vigor (Tables #3 and #4). The higher rated plants will have forage quality samples taken in 2000.

## **2000**

The assembly was evaluated for mid season forage production, quality and vigor on June 27, 2000. The entire planting was then clipped to a height of six inches on June 28, 2000. The assembly was evaluated for amount of regrowth and vigor on July 25, 2000 and forage quality samples were taken on August 1, 2000. The assembly was clipped the second time on August 2, 2000 and evaluations for regrowth amount and vigor were taken October 24, 2000.

## **2001**

Evaluations from previous years were correlated and the best plants from the top 10-20 percent of the total accessions were propagated in the greenhouse from clonal material from each individual plant. Plants were then isolated in two locations. A northern region was established containing plants from Iowa, northern Missouri, and northern Illinois. A southern region was established containing plants from southern Missouri and central and southern Illinois. These isolation blocks will receive additional evaluation to remove unwanted plants and the remaining plants will be allowed to produce seed. Plants from this seed will be selected for the next evaluation nursery. After further evaluation, plants from the nursery planted in 2003 will be used as a breeder's block for improved selections. Plants selected for each region can be found in Table #5.

## **2002**

The south region crossing block did very well in 2002. Very few plants were rouged out and seed was harvested from each accession in the block. This seed will be used to establish the next evaluation nursery scheduled for 2003.

The north region crossing block did not do well in 2002. Weed control became a problem and many of the plants were reestablished and did not make seed. Filling in additional plants is scheduled for 2002 and also seed production from this crossing block.

<b>Study 29I141G - Assembly and Evaluation of Little Bluestem, <i>Schizachyrium scoparium</i>, Nichx.</b>					
<b>Little Bluestem</b>					<b>Table #1</b>
	<b>REFERENCE</b>				
<b>ACCESSION</b>	<b>NUMBER</b>	<b>COLLECTOR</b>	<b>MLRA</b>	<b>COUNTY</b>	<b>STATE</b>
9078894	MO-1	Robert S. Crowder	M115	Chariton	Missouri
9078951	MO-2	Robert J. Crowder/ George L. Pollard	109	Chariton	Missouri
9078895	MO-3	Joe Tousignant	N116B	Cape Girardeau	Missouri
9078896	MO-4	Douglas Rainey	M115	Clark	Missouri
9078897	MO-5	David S. Mackey	113	Knox	Missouri
9078898	MO-6	Larry R. Brewer	M109	Putnam	Missouri
9078899	MO-7	Tommy Robins/ Jim Hoefer	116	Ripley	Missouri
9078900	MO-8	Grant P. Butler	N116B	Jefferson	Missouri
9078901	MO-9			Iron	Missouri
9078902	MO-10	Tommy Robins/ Jim Hoefer	116	Carter	Missouri
9078903	MO-11	Arch J. Mueller	M115	Ste. Genevieve	Missouri
9078904	MO-12			St. Francois	Missouri
9078905	MO-13	J. Mark Mitchell		Butler	Missouri
9078906	MO-14	Randy C. Miller	N116A	Shannon	Missouri
9078907	MO-15	Tom Johnson	N116B	Bollinger	Missouri
9078908	MO-16	Tom Johnson	N116A	Bollinger	Missouri
9078909	MO-17	Randy C. Miller	N116B	Reynolds	Missouri
9078910	MO-18			Franklin	Missouri
9078911	MO-19	Tom Johnson	N116A	Wayne	Missouri
9078912	MO-20	Mark E. Nussbaum	N116B	Cape Girardeau	Missouri
9078913	MO-21	Frank Oberle	115	Adair	Missouri
9078914	MO-22	David S. Mackey	113	Knox	Missouri
9078915	MO-23	Claude F. Peifer	116B	Perry	Missouri
9078916	MO-24	Grant P. Butler/ Bryan L. Westfall	N116A	Washington	Missouri
9078917	MO-25	John E. Turner	113/115	Monroe	Missouri
9078918	MO-26	David S. Mackey	113	Knox	Missouri
9078919	MO-27	Douglas Rainey	M115	Clark	Missouri
9078920	MO-28	Frank Oberle	115	Adair	Missouri
9078921	MO-29		M115	Montgomery	Missouri
9078922	MO-30	David S. Mackey	113	Knox	Missouri
9078923	MO-31	Curtis W. Walker	109	Clinton	Missouri
9078924	MO-32	James A. Mayberry	109	Carroll	Missouri
9078925	MO-33	Gary J. Barker	M109	Gentry	Missouri
9078926	MO-34			Vernon	Missouri
9078927	MO-35	Louis Byford		Atchison	Missouri
9078928	MO-36	Todd E. Mason	M109	Worth	Missouri
9078929	MO-37	Louis Byford		Atchison	Missouri
9078930	MO-38	Louis Byford		Atchison	Missouri
9078931	MO-39	Ronald L. Musick	M109	Harrison	Missouri

Study 29I141G - Little Bluestem			Table #1 - continued		
ACCESSION	REFERENCE NUMBER	COLLECTOR	MLRA	COUNTY	STATE
9078932	MO-40	Gary J. Barker	M109	Gentry	Missouri
9078933	MO-41	Curtis Walker	109	Gentry	Missouri
9078934	MO-42	Curtis Walker	107	Buchanan	Missouri
9078935	MO-43	Louis Byford		Atchison	Missouri
9078936	MO-44	Ronald L. Musick	M109	Harrison	Missouri
9078937	MO-45	Louis Byford		Atchison	Missouri
9078938	MO-46	Louis Byford		Atchison	Missouri
9078939	MO-47	Bob Sipec		Holt	Missouri
9078940	MO-48	Bib Sipec		Holt	Missouri
9078941	MO-49	Bob Sipec		Holt	Missouri
9078942	MO-50	Ian S. Kurtz	116A	Taney	Missouri
9078943	MO-52	Dennis Shirk/ Ed Gillmore	115	Gasconade	Missouri
9078944	MO-53	Dennis Shirk/ Ed Gillmore	116	Osage	Missouri
9078945	MO-54	Raleigh Redman	112	Henry	Missouri
9078946	MO-55	Dennis Shirk/ Ed Gillmore	116	Maries	Missouri
9078947	MO-56	Jerry Cloyed	M112	Barton	Missouri
9078948	MO-57	Ian S. Kurtz	116A	Taney	Missouri
9078949	MO-58	Ben A. Reed	M112	Barton	Missouri
9078950	MO-59	Jerry Cloyed	M112	Barton	Missouri
9078952	MO-60	M. Denise Brown	N116A	Miller	Missouri
9078953	MO-61	M. Denise Brown	N116B	Miller	Missouri
9078954	MO-62	Howard L. Coambes	N116B	Cedar	Missouri
9078955	MO-63	Howard L. Coambes	N116B	Cedar	Missouri
9078956	MO-64	Douglas G. Newman		Shannon	Missouri
9078957	MO-65	Tom E. Toney		Wayne	Missouri
9078958	MO-66	Rod Doolen		Wayne	Missouri
9078959	MO-67	Rod Doolen		Wayne	Missouri
9078960	MO-68	Kenneth L. Dalrymple		Pike	Missouri
9078963	MO-69	Maurice Davis/ Steve Clubine		Pettis	Missouri
	MO-70	Maurice Davis/ Steve Clubine		Benton	Missouri
	MO-71	Maurice Davis/ Steve Clubine		St. Clair	Missouri
	MO-72	Maurice Davis/ Steve Clubine		Benton	Missouri
9078964	MO-73	Maurice Davis/ Steve Clubine		Pettis	Missouri
9078965	MO-74	Maurice Davis/ Steve Clubine		Pettis	Missouri

Study 29I141G - Little Bluestem			Table #1 - continued		
ACCESSION	REFERENCE NUMBER	COLLECTOR	MLRA	COUNTY	STATE
	MO-75	Maurice Davis/ Steve Clubine		Pettis	Missouri
	MO-76	Maurice Davis/ Steve Clubine		Benton	Missouri
9078966	MO-77	Maurice Davis/ Steve Clubine		Maries	Missouri
9078967	MO-78	Dennis Shirk		Maries	Missouri
9078968	MO-79	Steve Clubine		Maries	Missouri
9078969	MO-80	Maurice Davis		Maries	Missouri
9078970	MO-81			Lawrence	Missouri
9078961	IA-27	Robert R. Bryant/ Shawn Dettman	108	Scott	Iowa
9078847	IA-1	Curt Donohue	109	Clarke	Iowa
9078848	IA-2	Curt Donohue	109	Clarke	Iowa
9078849	IA-3	Janet M. Thomas/ John P. Vogel	107	Cherokee	Iowa
9078850	IA-4	John P. Vogel	107	Woodbury	Iowa
9078851	IA-5	Henry D. Tordoff	107	West Pottawattamie	Iowa Iowa
9078852	IA-6	Henry D. Tordoff/ Galen Barrett	107	West Pottawattamie	Iowa Iowa
9078853	IA-7	John P. Vogel	107	Woodbury	Iowa
9078854	IA-8	Henry D. Tordoff	107	West Pottawattamie	Iowa Iowa
9078855	IA-9	John P. Vogel	107	Plymouth	Iowa
9078856	IA-10	Henry D. Tordoff	107	West Pottawattamie	Iowa Iowa
9078857	IA-11	Julie K. Watkins/ Charlie E. Kiepe	108	Franklin	Iowa
9078858	IA-12	Brad Harrison	103	Dallas	Iowa
9078859	IA-13	Shawn A. Dettman	108	Muscatine	Iowa
9078860	IA-14	Jim Ranum	105	Allamakee	Iowa
9078861	IA-15	Rick Cordes	104	Howard	Iowa
9078862	IA-16	James Ranum	105	Allamakee	Iowa
9078863	IA-17	Jay E. Ford	107	Crawford	Iowa
9078864	IA-18	Steve Maternack	103	Polk	Iowa
9078865	IA-19	Jay E. Ford	107	Crawford	Iowa
9078866	IA-20	Jay E. Ford	107	Crawford	Iowa
9078867	IA-21	Al Ehley	104	Cerro Gordo	Iowa
9078868	IA-22	Al Ehley	104	Cerro Gordo	Iowa
9078869	IA-23	John P. Vogel	102	Lyon	Iowa
9078870	IA-24	Jay E. Ford	107	Crawford	Iowa







Study 29I141G				Rep #2				Table #2 - continued			
Little Bluestem											
				North	↑						
PLT #	32 33 34	35 36 37	38 39 40	41 42 43	44 45 46	47 48 49	50 51 52	53 54 55	56 57 58	59 60 61	
TIER #											
I	? ? ?	W W W	W W W	W W b	b b b	b b b	b b b	b b b	b R R	R R R	I
II	MO-34	IL-18	IA-7	MO-31	MO-6	MO-53	MO-2	IA-18	MO-22	MO-48	II
III	MO-71	MO-24	MO-35	IA-14	IA-23	IA-2	MO-74	MO-28	CAMPER	MO-57	III
IV	MO-8	MO-42	MO-67	IL-1	MO-60	MO-33	MO-37	MO-26	IL-21	IL-7	III
V	IA-13	IA-3	MO-9	MO-39	IL-16	IA-8	MO-15	MO-69	MO-14	MO-25	IV
VI	MO-50	CIMMERON	IL-4	MO-59	MO-52	MO-40	MO-51	IA-27	MO-81	IA-16	VI
VII	IA-17	MO-63	MO-66	IL-20	MO-72	IL-19	MO-19	MO-23	IL-11	IL-10	VII
VIII	MO-32	IA-6	MO-4	IA-11	IL-2	MO-54	IA-26	IL-8	MO-41	IA-4	VIII
IX	IA-10	MO-77	IL-5	MO-46	MO-56	MO-64	MO-1	MO-21	MO-65	MO-10	IX
X	IL-14	MO-38	MO-49	MO-27	IL-12	MO-79	IA-19	MO-68	IA-1	ALDOUS	X
XI	MO-61	IA-9	MO-55	IL-15	IA-25	MO-17	MO-7	IA-5	IL-9	IL-3	XI
XII	IA-24	MO-47	MO-78	MO-43	PASTURA	MO-20	MO-73	MO-12	IA-20	MO-13	XII
XIII	MO-30	MO-18	MO-11	IL-6	MO-3	IL-13	IA-12	IA-22	MO-29	MO-44	XIII
XIV	MO-45	MO-62	MO-58	MO-5	IL-17	IA-21	MO-80	MO-16	MO-36	IA-15	XIV
XV	Y Y S	i S S	S S S	S S S	S S S	S S S	h S S	Y h Y	Y Y Y	h J J	XV



Study 29I141G												Rep #4		Table #2 - continued	
Little Bluestem															
PLT #	94 95 96	97 98 99	100 101 102	103 104 105	106 107 108	109 110 111	112 113 114	115 116 117	118 119 120	121 122 123	124				
TIER #															
I	R i R	a a a	X X X	X X U	i U U	U U U	U U U	W W W	W W W	W W W	d	I			
II	IA-9	IL-18	MO-8	MO-74	MO-40	IA-25	MO-5	MO-42	IA-4	IA-20	d	II			
III	MO-58	IA-19	MO-28	IL-17	MO-53	IL-8	PASTURA	MO-37	IL-10	MO-77	d	III			
IV	ALDOUS	MO-80	IA-21	MO-2	IA-8	MO-26	IA-26	MO-68	MO-14	MO-52	d	IV			
V	MO-51	IA-18	MO-20	MO-46	IL-1	MO-1	MO-62	MO-44	MO-9	MO-34	d	V			
VI	IA-17	IA-10	MO-33	IA-24	MO-43	IL-12	IA-5	MO-81	CIMMERON	MO-19	d	VI			
VII	MO-64	IA-10	CAMPER	MO-3	MO-69	MO-61	IA-16	IL-4	MO-35	MO-21	d	VII			
VIII	IA-27	MO-39	IL-19	MO-57	IL-6	MO-38	MO-67	MO-25	MO-48	IL-14	e	VIII			
IX	MO-60	MO-15	MO-63	IA-7	MO-36	IL-15	MO-49	IA-13	MO-29	MO-30	e	IX			
X	MO-12	MO-41	MO-32	MO-55	IA-12	MO-47	IA-26	IL-21	MO-65	IL-9	e	X			
XI	IL-20	IA-23	IA-11	MO-46	MO-17	IL-2	IL-13	MO-45	IL-11	IA-22	f	XI			
XII	MO-50	MO-6	MO-59	IA-14	MO-31	MO-54	MO-79	IA-3	MO-16	IL-7	f	XII			
XIII	MO-71	MO-78	MO-27	MO-73	MO-18	IA-15	MO-66	MO-72	MO-22	MO-10	f	XIII			
XIV	MO-7	MO-11	IL-16	MO-23	IA-1	IL-5	IA-6	MO-13	IL-3	MO-56	f	XIV			
XV	c R R	MO-24	R h R	R S h	h S S	S S T	h h h	T V V	V h g	g g g	g	XV			
.															

Study 29I141G			Forage Rating: 8/9/99										Table #3			
Little Bluestem																
			1 = High					9 = Low								
													Ave.			
Local	Rep 1			Rep 2			Rep 3			Rep 4			Percent	Living	Best	
Number	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	Survival	Plants	Plant	Location/s
MO-7	2	3	1	3	3	2	2	1	2	5	1	3	100	2.33	1	P 1, 8, 11
MO-12	1	2	1	3	2	2	3	2	2	1	1	1	100	1.75	1	P 1, 3, 12, 11, 12
MO-21	1	2	2	6	2	3	4	3	3	4	4	5	100	3.25	1	P 1
MO-74	3	3	5	4	4	4	5	5	4	1	2	1	100	3.42	1	P 10, 12
MO-80	3	3	x	4	5	5	4	4	2	1	4	3	92	3.45	1	P 10
MO-4	x	5	5	4	8	2	3	4	4	6	x	x	83	4.10	2	P 6
MO-9	4	4	4	3	4	4	3	4	3	2	3	3	100	3.42	2	P 10
MO-14	4	4	3	4	4	4	5	2	2	4	4	3	100	3.58	2	P 8, 9
MO-15	3	2	3	5	4	3	6	4	5	4	3	5	100	3.92	2	P 2
MO-22	4	5	5	3	4	2	5	5	6	x	8	x	83	4.70	2	P 6
MO-23	3	5	6	2	6	8	5	4	5	8	8	3	100	5.73	2	P 4
MO-24	3	x	2	x	4	4	3	4	3	3	4	5	83	3.18	2	P 3
MO-32	4	x	8	6	7	3	3	4	5	2	5	6	92	4.82	2	P 10
MO-34	4	4	4	3	4	3	x	x	4	2	x	5	75	3.00	2	P 10
MO-37	2	4	3	7	5	4	x	5	4	3	4	3	92	3.67	2	P 1
MO-42	5	5	6	4	5	2	4	4	4	5	5	7	100	4.67	2	P 6
MO-50	3	3	4	2	2	2	3	4	6	2	3	4	100	3.17	2	P 4, 5, 6, 10
MO-51	3	3	3	3	4	4	4	6	3	4	3	2	100	3.50	2	P 12
MO-53	4	4	5	5	5	5	2	4	5	5	6	7	100	4.75	2	P 7
MO-56	3	3	2	2	5	4	5	3	3	3	3	3	100	3.25	2	P 3, 4
MO-58	3	3	3	5	4	5	5	5	5	2	2	4	100	3.83	2	P 10, 11
MO-59	2	3	4	4	4	5	3	3	3	3	4	4	100	3.50	2	P 1
MO-66	3	3	x	3	3	3	3	2	4	4	5	5	92	3.45	2	P 8
MO-73	7	4	4	3	3	2	4	5	5	7	8	6	100	4.83	2	P 6
MO-79	2	3	2	5	3	5	3	8	5	4	4	3	100	3.92	2	P 1, 3
MO-2	4	5	3	5	5	5	5	3	3	3	4	3	100	4.00	3	P 3, 8, 9, 10, 12
MO-5	7	3	3	5	5	5	6	8	4	4	5	4	100	4.92	3	P 2, 3
MO-8	6	x	5	5	4	5	7	4	8	3	3	4	92	4.91	3	P 10, 11
MO-10	4	5	5	3	3	5	5	5	5	7	5	4	100	4.67	3	P 4, 12
MO-11	x	7	x	4	5	6	6	6	5	3	3	6	83	4.25	3	P 10, 11
MO-13	5	8	5	5	x	5	4	4	3	6	4	6	100	4.58	3	P 9
MO-16	4	3	8	6	6	5	4	6	4	4	5	100	75	3.00	3	P 2
MO-17	4	4	3	4	3	7	8	6	5	4	5	5	100	4.83	3	P 3, 5
MO-18	3	4	3	7	7	8	x	x	x	5	5	5	75	3.92	3	P 1, 3
MO-19	3	5	5	3	4	3	4	6	5	3	5	4	100	4.17	3	P 1, 4, 6, 10
MO-20	8	7	6	7	6	5	3	4	5	4	8	3	100	6.60	3	P 7, 12
MO-25	3	3	x	5	5	5	5	4	6	5	5	6	92	4.33	3	P 1, 2
MO-26	3	4	4	5	x	4	3	4	4	3	4	5	92	4.30	3	P 1, 7, 10
MO-27	5	6	3	4	5	4	6	5	4	5	5	7	100	5.36	3	P 3
MO-29	4	3	x	4	5	4	4	6	3	3	5	8	92	4.45	3	P 2, 9, 10
MO-30	3	4	5	7	7	x	4	4	7	4	3	4	92	4.73	3	P 1, 11
MO-31	7	3	4	4	4	6	7	8	x	5	5	5	92	5.27	3	P 2

Study 29I141G			Forage Rating: 8/9/99												Table #3 - continued		
Little Bluestem																	
			1 = High				9 = Low										
															Ave.		
Local	Rep 1			Rep 2			Rep 3			Rep 4			Percent	Living	Best		
Number	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	Survival	Plants	Plant	Location/s	
MO-33	3	x	3	5	5	3	4	5	5	8	8	4	92	5.89	3	P 1, 3, 6	
MO-35	4	7	8	5	6	7	5	3	6	5	4	x	92	5.45	3	P 8	
MO-38	6	6	5	3	3	4	4	6	7	3	3	4	100	5.40	3	P 4, 5, 10, 11	
MO-41	5	6	5	4	4	7	6	x	4	3	x	5	83	4.90	3	P 10	
MO-43	4	4	x	5	5	5	5	6	5	4	3	4	92	4.55	3	P 11	
MO-46	4	x	4	4	3	3	3	5	5	4	4	4	92	3.91	3	P 5, 6, 7	
MO-47	5	6	6	6	5	4	3	4	5	5	8	4	100	5.08	3	P 7	
MO-48	3	7	8	5	5	6	4	4	6	4	5	5	100	5.17	3	P 1	
MO-52	3	3	3	4	3	3	4	5	4	4	3	4	100	3.58	3	P 1, 2, 3, 5, 6, 11	
MO-54	x	x	x	5	5	5	4	5	5	6	4	3	75	4.67	3	P 12	
MO-57	4	4	x	3	5	x	4	4	x	5	4	3	92	3.27	3	P 4, 12	
MO-60	7	4	6	4	6	3	6	4	6	5	5	4	100	5.00	3	P 6	
MO-61	5	8	6	x	4	5	x	8	8	3	7	5	83	5.90	3	P 10	
MO-65	4	5	6	7	x	x	4	5	3	4	6	6	83	5.00	3	P 9	
MO-67	3	3	3	3	3	3	6	5	x	3	3	3	92	3.45	3	P 1, 2, 3, 4, 5, 6, 10, 11, 12	
MO-69	4	5	4	3	3	5	4	5	4	7	4	5	100	4.42	3	P 3, 4	
MO-71	x	5	5	4	3	5	4	4	5	4	5	3	92	4.27	3	P 5, 12	
MO-77	6	x	6	4	6	4	3	4	5	6	6	5	92	5.00	3	P 7	
MO-78	5	6	5	5	3	5	3	5	6	4	3	3	100	4.42	3	P 5, 7, 11, 12	
MO-1	4	5	4	4	4	6	4	7	5	4	5	5	100	4.75	4		
MO-3	4	7	4	5	4	4	4	4	4	5	4	5	100	4.50	4		
MO-6	7	7	7	7	7	5	x	8	7	4	4	4	92	6.09	4		
MO-28	6	5	6	6	7	5	4	7	7	4	x	x	83	4.75	4		
MO-36	4	4	5	6	6	6	x	5	5	5	6	5	92	5.18	4		
MO-39	4	6	7	4	6	4	6	5	x	6	5	x	83	5.89	4		
MO-40	7	6	7	5	4	4	x	6	5	5	5	5	92	5.36	4		
MO-44	7	4	5	5	6	7	7	x	6	5	4	6	92	5.64	4		
MO-45	4	4	4	5	6	6	5	6	5	4	4	4	100	4.75	4		
MO-49	6	5	6	6	5	x	5	5	4	7	5	6	92	5.45	4		
MO-55	x	6	x	4	4	5	4	5	x	8	x	5	67	5.13	4		
MO-62	4	4	5	5	4	5	5	7	6	5	5	6	100	5.08	4		
MO-63	5	6	5	5	4	4	8	4	6	4	5	5	100	5.08	4		
MO-68	7	6	6	6	8	4	5	6	5	4	4	4	100	5.42	4		
MO-72	5	6	5	5	6	5	4	6	6	5	4	4	100	5.08	4		
MO-81	x	4	5	5	4	6	x	x	x	6	x	8	58	5.43	4		
MO-64	x	7	6	7	6	6	6	5	8	x	7	5	92	5.73	5		
MO-70																	
MO-75																	
MO-76																	

Study 29I141G				Forage Rating: 8/9/99												Table #3 - continued			
Little Bluestem																			
				1 = High			9 = Low												
																Ave.			
Local	Rep 1			Rep 2			Rep 3			Rep 4			Percent	Living	Best				
Number	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	Survival	Plants	Plant	Location/s			
IA-16	x	x	4	3	6	5	3	x	1	x	5	5	75	3.56	1	P 9			
IA-27	1	1	3	3	4	5	5	5	4	5	4	2	100	3.50	1	P 1, 2			
IA-6	4	5	6	5	2	4	3	4	3	7	4	5	100	4.33	2	P 5, 6			
IA-8	5	6	3	5	3	5	5	5	5	5	3	2	100	4.33	2	P 12			
IA-12	7	5	7	x	4	5	4	3	2	4	5	5	92	4.64	2	P 9			
IA-15	5	4	5	x	x	x	2	x	5	5	5	6	67	4.63	2	P 7			
IA-23	6	5	5	8	8	6	5	4	x	2	4	6	92	5.36	2	P 10			
IA-1	8	5	5	5	4	4	4	5	x	3	7	3	92	4.82	3	P 10, 12			
IA-2	4	4	4	3	4	4	6	5	5	4	x	6	92	4.45	3	P 4			
IA-3	x	x	8	x	3	3	4	5	4	4	5	4	75	4.44	3	P 5, 6			
IA-4	5	8	4	3	x	3	4	7	5	4	7	5	92	5.00	3	P 4, 6			
IA-5	4	5	4	3	6	8	6	4	4	3	5	x	92	4.73	3	P 4, 10			
IA-7	5	3	3	5	5	5	4	4	6	5	5	5	100	4.58	3	P 2, 3			
IA-9	4	6	7	6	6	6	8	6	6	4	3	4	100	5.50	3	P 11			
IA-11	6	5	6	5	7	3	5	5	6	4	x	5	92	5.18	3	P 6			
IA-13	4	4	6	4	7	x	5	4	x	3	4	3	83	4.40	3	P 10, 12			
IA-17	3	7	4	5	x	4	6	x	6	4	6	5	83	5.00	3	P 1			
IA-19	6	x	x	6	3	3	x	4	4	x	x	x	50	4.33	3	P 5, 6			
IA-20	x	4	x	7	5	5	4	x	4	6	7	3	75	5.00	3	P 12			
IA-24	4	5	3	5	4	4	4	4	5	5	5	4	100	4.33	3	P 3			
IA-25	4	5	6	6	5	6	6	4	5	3	5	3	100	4.83	3	P 10, 12			
IA-26	x	3	4	3	3	6	x	x	4	5	6	x	67	4.25	3	P 2, 4, 5			
IA-10	6	7	7	4	5	5	5	6	7	6	4	x	92	5.64	4				
IA-14	4	6	4	5	5	6	4	5	5	5	7	5	100	5.08	4				
IA-18	5	6	5	6	5	6	5	4	5	4	5	5	100	5.08	4				
IA-21	4	5	4	4	x	6	x	x	6	-	4	5	67	4.75	4				
IA-22	x	x	x	7	x	x	7	6	6	5	8	8	58	6.71	5				
IL-12	8	7	5	3	8	4	5	5	4	4	2	x	92	5.00	2	P 11			
IL-17	3	4	3	2	3	5	3	4	2	2	3	3	100	3.08	2	P 4, 9, 10			
IL-18	5	4	6	3	3	3	5	6	4	3	2	4	100	4.00	2	P 11			
IL-2	6	6	6	4	5	6	5	3	5	4	5	3	100		3	P 8			
IL-5	6	5	7	4	8	3	4	5	5	5	4	5	100	5.08	3	P 6			
IL-7	4	4	3	4	7	6	8	6	8	6	8	8	100	6.00	3	P 3			
IL-8	x	x	5	4	x	8	x	6	4	x	4	3	58	4.86	3	P 12			
IL-11	x	x	3	x	4	x	5	x	6	x	x	x	33	4.50	3	P 3			
IL-14	4	5	x	3	5	x	6	4	7	6	5	6	83	5.10	3	P 4			
IL-16	5	5	4	4	3	3	4	x	3	7	6	4	92	4.36	3	P 5, 6, 9			
IL-19	5	6	7	3	3	3	4	3	4	3	4	3	100	4.00	3	P 4, 5, 6, 8, 12			

Study 29I141G			Forage Rating: 8/9/99										Table #3 - continued			
Little Bluestem																
			1 = High			9 = Low										
													Ave.			
Local	Rep 1			Rep 2			Rep 3			Rep 4			Percent	Living	Best	
Number	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	Survival	Plants	Plant	Location/s
IL-20	5	3	3	x	6	5	4	4	4	3	5	3	92	4.09	3	P 2, 3, 10, 12
IL-21	5	5	4	3	4	4	5	4	4	5	4	4	100	4.25	3	P 4
IL-1	4	x	4	6	7	6	4	7	7	5	6	5	92	5.55	4	
IL-6	7	7	4	6	5	7	x	x	x	6	5	5	75	5.78	4	
IL-9	6	x	6	x	5	7	6	5	4	4	4	7	83	5.40	4	
IL-10	x	x	x	4	6	7	x	x	7	x	5	7	50	6.00	4	
IL-13	x	7	x	5	7	4	6	6	7	x	8	6	83	5.60	4	
IL-15	8	8	x	x	7	6	4	5	5	5	4	5	83	5.70	4	
IL-3	5	4	x	7	x	x	8	7	6	5	x	x	58	6.00	5	
IL-4	6	7	4	4	6	5	6	5	5	5	5	5	100	5.25	5	
IL-22																
Aldous	2	3	3	3	3	3	5	4	5	3	2	2	100	3.17	2	P 1, 11, 12
Cimmeron	2	3	2	4	2	3	3	2	5	3	5	3	100	3.08	2	P 1, 3, 5, 8
Camper	3	4	5	4	5	6	5	4	5	x	3	5	92	4.45	3	P 1, 11, 12
Pastura	x	x	5	6	x	6	6	6	x	3	3	x	58	5.00	3	P 10, 11

Study 29I141G			Vigor Rating: 8/9/99											Table #4		
Little Bluestem																
1 = High 9 = Low																
Local	Rep 1			Rep 2			Rep 3			Rep 4			Percent	Living	Best	
Number	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	Survival	Plants	Plant	Location/s
														Ave.		
MO-4	x	3	4	4	6	2	4	5	5	3	x	x	75	4.00	2	P 6
MO-7	2	3	2	3	3	3	5	2	2	5	2	2	100	2.83	2	P 1, 3, 8, 9, 11, 12
MO-12	3	3	3	4	2	2	4	4	3	3	3	3	100	3.08	2	P 5, 6
MO-16	3	2	6	6	4	3	4	5	6	4	5	3	100	4.25	2	P 2
MO-24	5	x	5	x	5	3	5	5	5	2	4	6	83	4.50	2	P 10
MO-25	2	3	x	5	4	3	5	4	4	5	6	6	92	4.27	2	P 1
MO-32	3	x	6	5	5	3	4	6	4	2	4	6	92	4.36	2	P 10
MO-35	2	6	7	2	4	5	6	6	3	5	4	x	92	4.55	2	P 1, 4
MO-42	5	4	5	3	4	2	4	4	6	4	5	6	100	4.33	2	P 6
MO-47	4	5	6	4	5	4	2	2	3	4	6	3	100	4.00	2	P 7, 8
MO-56	3	4	3	3	3	2	4	4	4	4	3	3	100	3.33	2	P 6
MO-61	5	5	4	x	3	4	x	7	7	2	5	4	83	4.60	2	P 10
MO-67	3	3	3	2	3	3	5	4	x	4	5	5	92	3.64	2	P 4
MO-69	4	5	6	3	3	4	2	3	5	8	4	5	100	4.33	2	P 7
MO-79	2	3	3	3	3	4	5	6	4	5	4	3	100	3.75	2	P 1
MO-1	3	4	3	4	3	5	5	5	5	3	5	4	100	4.08	3	P 1, 3, 5, 10
MO-3	3	4	4	5	4	3	4	5	5	4	3	4	100	4.00	3	P 1, 6, 11
MO-5	5	3	3	5	4	6	5	7	4	5	6	4	100	4.75	3	P 2, 3
MO-6	3	7	6	6	5	5	x	5	5	5	5	3	92	5.00	3	P 1, 12
MO-8	5	x	4	6	3	3	6	6	5	6	5	7	92	5.09	3	P 5, 6
MO-9	5	5	6	3	3	3	4	4	4	5	5	5	100	4.33	3	P 3, 4, 5
MO-11	x	5	x	5	6	6	7	5	3	5	4	6	83	5.20	3	P 9
MO-13	5	7	6	6	x	5	5	6	3	6	5	7	92	5.55	3	P 9
MO-14	4	4	3	5	5	5	4	6	6	4	5	4	100	4.58	3	P 3
MO-15	3	3	3	4	3	3	5	4	4	4	4	4	100	3.67	3	P 1, 2, 3, 5, 6
MO-17	5	5	5	4	4	7	7	5	4	3	4	5	100	4.83	3	P 10
MO-19	3	3	3	4	4	4	4	5	5	4	4	4	100	3.92	3	P 1, 2, 3, 5, 6
MO-21	3	3	3	6	4	4	5	4	4	6	6	6	100	4.50	3	P 1, 2, 3
MO-22	4	3	3	3	3	3	5	5	4	x	7	x	83	4.00	3	P 2, 3, 4, 5, 6
MO-23	5	5	3	4	5	7	5	5	6	5	7	5	100	5.17	3	P 3
MO-26	4	4	4	3	x	3	6	5	5	4	5	5	92	4.36	3	P 4, 6
MO-27	3	5	3	3	4	3	6	6	5	3	4	6	100	4.25	3	P 1, 3, 4, 6, 10
MO-29	4	3	x	6	5	5	5	5	3	5	5	6	92	4.73	3	P 2, 9
MO-31	6	3	3	3	3	4	5	5	x	6	4	5	92	4.27	3	P 2, 3, 4, 5
MO-33	4	x	6	4	4	3	4	4	4	6	6	5	92	4.55	3	P 6
MO-34	4	3	3	3	3	4	x	x	4	4	x	3	75	3.44	3	P 2, 3, 4, 5, 12
MO-36	4	3	3	6	5	5	x	4	5	3	5	5	92	4.36	3	P 2, 3, 10
MO-37	3	3	3	4	3	4	x	5	5	4	4	4	92	3.82	3	P 1, 2, 3, 5
MO-38	4	4	3	5	4	4	3	5	5	3	3	4	100	3.92	3	P 7, 10, 11
MO-39	5	6	7	4	3	3	5	3	x	5	7	x	83	4.80	3	P 5, 6, 8
MO-40	3	8	8	4	5	3	x	5	4	8	8	7	100	5.25	3	P 1, 6, 11

Study 29I141G			Vigor Rating: 8/9/99											Table #4 - continued		
Little Bluestem																
1 = High 9 = Low																
Local	Rep 1			Rep 2			Rep 3			Rep 4			Percent	Living	Best	
Number	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	Survival	Plants	Plant	Location/s
MO-43	6	3	4	4	4	4	5	6	5	4	5	3	100	4.42	3	P 2, 12
MO-45	4	4	3	3	4	3	4	5	3	3	4	4	100	3.67	3	P 3, 4, 6, 9, 10
MO-46	3	x	3	3	3	4	5	5	3	5	3	4	92	3.73	3	P 1, 3, 4, 5, 9, 11
MO-48	4	5	5	3	4	4	5	3	5	4	6	6	100	4.50	3	P 4, 8
MO-51	4	5	4	3	3	3	4	5	4	4	4	4	100	3.92	3	P 3, 4, 5
MO-52	5	4	5	5	3	4	5	6	5	5	4	5	100	4.67	3	P 5
MO-53	5	5	6	4	5	6	3	4	4	5	5	6	100	4.83	3	P 7
MO-54	x	x	x	5	7	3	6	7	7	6	3	4	75	5.33	3	P 11
MO-60	4	4	4	3	4	3	5	3	5	5	6	6	100	4.33	3	P 4, 6, 8
MO-62	4	4	4	3	4	5	4	4	4	5	6	7	100	4.50	3	P 4
MO-63	4	4	4	3	3	3	5	5	4	4	6	4	100	4.08	3	P 4, 5, 6
MO-65	3	4	4	6	x	x	5	6	5	5	7	6	83	5.10	3	P 1
MO-66	5	5	x	4	3	3	6	6	5	6	7	7	92	5.18	3	P 5, 6
MO-71	x	3	5	5	3	4	5	4	5	3	4	4	92	4.09	3	P 2, 5, 10
MO-72	3	3	3	3	5	4	3	4	5	5	4	3	100	3.75	3	P 1, 2, 3, 4, 7, 12
MO-73	6	5	3	3	3	3	5	7	4	6	7	6	100	4.83	3	P 3, 4, 5, 6
MO-77	6	x	6	5	3	5	3	4	5	6	6	6	92	5.00	3	P 5, 7
MO-78	6	4	4	4	6	4	4	5	3	4	4	3	100	4.25	3	P 9, 12
MO-80	4	3	x	3	3	3	6	6	5	3	6	6	92	4.36	3	P 2, 4, 5, 6, 10
MO-81	x	3	5	5	4	4	x	x	x	6	x	5	58	4.57	3	P 2
MO-2	4	5	5	4	5	6	4	4	5	4	4	4	100	4.50	4	
MO-18	4	6	4	4	5	7	x	x	x	6	4	6	75	5.11	4	P 1, 3, 4, 11
MO-20	4	6	6	6	5	5	6	5	5	4	6	4	100	5.17	4	
MO-28	6	4	5	4	6	5	5	6	5	4	x	x	83	5.00	4	
MO-30	4	5	5	4	4	x	5	5	6	5	4	4	92	4.64	4	
MO-41	4	7	4	5	5	4	6	x	5	4	x	4	83	4.80	4	
MO-44	6	4	4	5	5	5	7	x	6	5	4	6	92	5.18	4	
MO-49	8	8	8	8	8	x	7	7	6	6	4	4	92	6.73	4	
MO-50	5	5	5	4	4	4	6	6	4	5	5	5	100	4.83	4	
MO-55	x	5	x	4	6	5	5	4	x	6	x	4	67	4.88	4	
MO-57	4	5	x	5	4	x	6	5	x	5	6	5	75	3.75	4	
MO-58	6	5	4	6	5	6	7	7	7	4	4	5	100	5.50	4	
MO-59	7	6	5	5	4	4	7	6	7	6	6	5	100	5.67	4	
MO-68	5	5	5	4	5	5	5	4	4	6	4	5	100	4.75	4	
MO-74	5	6	6	4	4	5	5	5	5	5	5	4	100	4.92	4	
MO-10	6	7	7	5	5	5	5	6	6	7	6	4	100	5.75	5	
MO-64	x	7	7	5	7	7	6	6	6	x	7	5	83	6.30	5	
MO-70																
MO-75																
MO-76																

Study 29I141G			Vigor Rating: 8/9/99												Table #4 - continued		
Little Bluestem																	
1 = High 9 = Low																	
Local	Rep 1			Rep 2			Rep 3			Rep 4			Percent	Living	Best		
Number	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	Survival	Plants	Plant	Location/s	
IA-3	x	x	5	x	3	2	6	6	7	7	5	5	75	5.11	2	P 6	
IA-4	4	5	3	4	x	3	4	6	4	2	5	5	92	4.09	2	P 10	
IA-5	6	6	6	4	5	6	6	5	6	2	5	x	92	5.18	2	P 10	
IA-9	4	4	4	4	4	5	6	5	5	3	2	5	100	4.25	2	P 11	
IA-10	3	4	5	3	4	4	5	5	5	6	2	x	92	4.18	2	P 11	
IA-13	2	3	4	3	5	x	5	4	x	4	5	3	92	3.45	2	P 1	
IA-15	5	4	4	x	x	x	2	x	6	4	4	5	67	4.25	2	P 7	
IA-27	2	2	2	2	3	3	5	6	5	4	3	3	100	3.33	2	P 1, 2, 3, 4	
IA-1	6	3	3	5	5	4	4	4	x	4	7	4	92	4.45	3	P 2, 3	
IA-2	3	3	3	4	5	5	6	5	5	5	x	6	92	4.55	3	P 1, 2, 3	
IA-6	6	4	4	4	3	3	5	4	4	7	3	5	100	4.33	3	P 5, 6, 11	
IA-7	3	3	4	3	3	3	3	4	6	4	4	4	100	3.67	3	P 1, 2, 4, 5, 6, 7	
IA-8	5	6	3	3	3	4	5	6	5	4	3	4	100	4.25	3	P 3, 4, 5, 11	
IA-12	4	5	6	x	5	4	3	5	4	3	3	3	92	4.09	3	P 7, 10, 11, 12	
IA-14	6	5	5	3	3	3	5	7	7	4	6	5	100	4.92	3	P 4, 5, 6	
IA-16	x	x	4	3	5	4	3	x	5	x	5	6	67	4.38	3	P 4, 7	
IA-17	4	6	5	4	x	4	5	x	4	3	5	3	83	4.30	3	P 10, 12	
IA-18	5	6	5	5	4	5	4	4	5	3	3	4	100	4.42	3	P 10, 11	
IA-23	4	4	4	5	6	6	5	5	x	3	3	4	100	4.08	3	P 10	
IA-25	5	5	5	5	4	4	4	5	5	4	4	3	100	4.42	3	P 12	
IA-26	x	6	4	3	4	5	x	x	4	4	6	x	67	4.50	3	P 4	
IA-11	7	6	7	4	5	4	6	6	7	5	x	5	92	5.64	4		
IA-19	6	x	x	5	4	4	x	4	4	x	x	x	50	4.50	4		
IA-20	x	4	x	7	5	5	5	x	6	5	6	5	75	5.33	4		
IA-21	4	4	5	4	x	5	x	x	4	x	5	4	67	4.38	4		
IA-22	x	x	x	5	x	x	5	4	4	6	8	8	58	5.71	4		
IA-24	5	5	4	6	6	6	7	7	7	6	5	5	100	5.75	5		
IL-8	x	x	6	4	x	5	x	2	3	x	5	3	58	4.00	2	P 8	
IL-12	6	6	2	3	5	3	4	4	3	3	2	x	92	3.73	2	P 3, 11	
IL-1	7	x	3	5	7	6	5	6	8	6	5	5	92	5.73	3	P 3	
IL-2	3	3	4	4	5	3	4	5	5	5	4	4	100	4.08	3	P 1, 2, 6	
IL-3	3	7	3	5	x	x	6	7	6	5	x	x	67	5.25	3	P 1, 3	
IL-5	5	5	6	5	3	4	5	6	5	5	4	5	100	4.83	3	P 5	
IL-6	7	5	4	8	3	5	x	x	x	5	4	7	75	5.33	3	P 5	
IL-9	5	x	3	x	4	5	5	3	3	5	4	6	92	3.91	3	P 3, 8, 9	
IL-10	4	4	5	5	4	3	x	x	8	x	6	6	100	3.75	3	P 6	
IL-11	x	x	3	x	4	x	3	x	5	x	x	x	33	3.75	3	P 3, 7	
IL-13	x	5	x	4	5	5	6	6	7	x	6	3	75	5.22	3	P 12	
IL-14	5	4	x	3	4	x	5	3	5	5	4	5	83	4.30	3	P 4, 8	
IL-15	5	7	x	x	5	4	6	6	5	4	4	3	83	4.90	3	P 12	

Study 29I141G		Vigor Rating: 8/9/99												Table #4 - continued		
Little Bluestem																
1 = High 9 = Low																
Local	Rep 1			Rep 2			Rep 3			Rep 4			Percent	Living	Best	
Number	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	Survival	Plants	Plant	Location/s
IL-16	3	3	4	4	4	5	5	x	6	7	5	4	92	4.55	3	P 1, 2, 6
IL-17	4	4	3	3	3	3	3	3	3	3	3	3	100	3.17	3	P 3, 4, 5, 6, 7, 8, 9, 10,,11, 12
IL-18	4	3	5	4	4	3	5	6	4	4	5	4	100	4.25	3	P 2, 6
IL-19	4	4	6	3	3	3	5	3	4	4	3	4	100	3.83	3	P 4, 5, 6, 8, 11
IL-20	4	5	4	x	3	4	4	3	4	4	6	3	92	4.00	3	P 5, 8, 12
IL-21	7	7	7	3	5	4	6	5	5	5	5	5	100	5.33	3	P 4
IL-4	6	5	5	4	4	4	5	5	5	6	6	5	100	5.00	4	
IL-7	4	4	5	4	5	6	6	5	6	6	7	6	100	5.33	4	
IL-22																
Cimmonon	2	2	2	2	3	2	3	5	4	3	4	3	100	2.92	2	P 1, 2, 3, 4, 6
Aldous	4	3	4	3	3	3	5	5	4	3	3	3	100	3.58	3	P 2, 4, 5, 6, 10, 11, 12
Camper	3	3	3	4	5	5	5	5	6	x	5	5	92	4.45	3	P 1, 2, 3
Pastura	x	x	5	5	x	7	5	7	x	3	4	x	58	5.14	3	P 10

<b>Study 29I141G - Assembly and Evaluation of Little Bluestem, <i>Schizachyrium scoparium</i>, Nichx.</b>					
<b>Selected accessions for each region</b>					<b>Table #5</b>
		<b>Northern Region</b>			
		IA - All			
		MO - North of Missouri River			
		IL - Northern 2/3rds of the state			
	<b>REFERENCE</b>				
<b>ACCESSION</b>	<b>NUMBER</b>	<b>COLLECTOR</b>	<b>MLRA</b>	<b>COUNTY</b>	<b>STATE</b>
9078896	MO-4	Douglas Rainey	115	Clark	Missouri
9078913	MO-21	Frank Oberle	115	Adair	Missouri
9078914	MO-22	David S. Mackey	113	Knox	Missouri
9078924	MO-32	James A. Mayberry	109	Carroll	Missouri
9078934	MO-42	Curtis Walker	107	Buchanan	Missouri
9078849	IA-3	Janet M. Thomas/ John P. Vogel	107	Cherokee	Iowa
9078854	IA-8	Henry D. Tordoff	107	West Pottawattamie	Iowa Iowa
9078861	IA-15	Rick Cordes	104	Howard	Iowa
9078862	IA-16	James Ranum	105	Allamakee	Iowa
9078884	IL-12	Bill Kleiman		Lee	Illinois
9078891	IL-19	Martha E. Sheppard	115	Calhoun	Illinois
		<b>Southern Region</b>			
		MO - South of Missouri River			
		IL - Southern 1/3 of state			
	<b>REFERENCE</b>				
<b>ACCESSION</b>	<b>NUMBER</b>	<b>COLLECTOR</b>	<b>MLRA</b>	<b>COUNTY</b>	<b>STATE</b>
9078895	MO-3	Joe Tousignant	N116B	Cape Girardeau	Missouri
9078899	MO-7	Tommy Robins/ Jim Hoefer	116	Ripley	Missouri
9078915	MO-23	Claude F. Peifer	116B	Perry	Missouri
9078942	MO-51	Ian S. Kurtz	116A	Taney	Missouri
9078950	MO-59	Jerry Cloyed	M112	Barton	Missouri
9078952	MO-60	M. Denise Brown	N116A	Miller	Missouri
9078964	MO-73	Maurice Davis/ Steve Clubine		Pettis	Missouri
9078965	MO-74	Maurice Davis/ Steve Clubine		Pettis	Missouri
9078968	MO-79	Steve Clubine		Marries	Missouri
9078969	MO-80	Maurice Davis/			Missouri
9078893	IL-21	Remington T. Irwin	114	Wayne	Illinois

**Study No. 29I142G**

**Study Title:** Production of Native Missouri Ecotypes of Grasses, Legumes and Forbs for Roadsides, Critical Areas, and All Other Vegetative Plantings Where Native Plants are Now Being Planted.

**Study Leader:** Bruckerhoff, S. B.

**Study Coordinator:** Erickson, R., Missouri Department of Conservation

**Introduction:**

Well-adapted native grass, legume and forb plantings offer many advantages as a low cost sustainable vegetative cover for management of soil and water resources. Native plant communities resist noxious weed invasion, provide excellent erosion control, and generally require relatively low maintenance.

These characteristics make native plants an excellent selection for use in roadside plantings, wildlife habitat enhancement, long-term land retirement programs, public land and all other vegetative plantings where mono-cultures of grasses are presently being planted. This is especially true along public transportation corridors that constitute a major land resource and management problem in the state of Missouri. Based on 1987 National Resource Inventory (NRI) data, over one million acres of Missouri land are devoted to rural transportation. Other federal and state agencies also own a significant land base in Missouri.

Proper vegetation management along these corridors is an important element in controlling soil loss and unwanted weedy plant species. Many of these acres are now seeded to introduced cool-season grass and legume species which are often invaded by noxious weeds requiring extensive mowing or herbicide treatment programs. These management techniques are expensive and can also result in additional water quality problems where herbicides are used extensively.

Managing or reseeding these acres to promote native grasses and forbs offers a low cost environmentally sound approach to roadside vegetation management. Herbicide use, soil erosion, and most mowing can be reduced significantly where a vigorous native grass and forb mixture dominates a roadside right-of-way. In addition, these goals are consistent with on-going NRCS programs designed to improve ground and surface water quality, reduce soil loss and increase wildlife habitat.

**Problem:**

Many adapted forb, legume and grass species of native origin are either currently not commercially available or available only in very limited quantities, which makes them very expensive. Species that are available are often varietal releases that have undergone an evaluation and selection process or a plant-breeding program. Most varieties are designed for high forage production and are highly vigorous plants. They are generally excellent for pasture and hay production but can be too domineering for diversified mixtures. Their origins are often not from within the state in which they are being planted. There is a need for additional native species for use on public lands and other types of conservation plantings with origins close to where they are being planted.

## Objective:

The objective of this study is to accelerate the availability of selected native grass, legume and forb species.


## Cooperators:

The Missouri Department of Conservation (MDC), USDA Natural Resources Conservation Service (NRCS), Plant Materials Center (PMC), the University of Missouri at Columbia, Missouri (UMC), and the National Audubon Society-Audubon Missouri (NAS).

## Procedures:

The state of Missouri was divided into four zones: Northern Glaciated Plains, Zone #1; Western Prairie, Zone #2; Ozarks, Zone #3; and the Bootheel Region, Zone #4 (See Table #1). Plant materials were collected as seed by the study coordinator, selected personnel from USDA-NRCS, MDC, UMC, and other knowledgeable interested persons. Collections were made from prairie remnants throughout each zone striving for a relatively equal and representative sample. Large collections from one site were not allowed to dominate the mixture from throughout the zone. Seed from each collection site was inventoried by location. Seed collected from within each zone was kept separate from the other zones. Increase plots were and will be established, as seed becomes available. Each species will be released as 'Source Identified' germplasm from the zone in which it was collected. Evaluation and selection or plant breeding procedures has not improved 'Source Identified' seed.

Table #1



## **Discussion:**

### **1997**

The Missouri Ecotype Enhancement Program was officially started as a plant materials study with the signing of the study plan in December of 1997. This plan is an agreement between cooperators and funded by a grant from the MDC. Several meetings preceded the document signing that included MDC, NRCS, UMC, Department of Transportation, Missouri Department of Natural Resources, and other interested individuals.

The initial grant from MDC to UMC was received July 1997 and a program coordinator was hired by UMC in September 1997 to work at the Elsberry Plant Materials Center.

### **1998**

A grant was given to UMC once again by MDC that would fund the program through August of 1999. Goals were established for 1998 collections. Some species from 1997 were recollected and new species were added. See Table #2 for listing of species collected and status of collections.

### **1999**

The Missouri Ecotype program continued during 1999 and the species released listed in Table #3. Beginning in September, the Lincoln County Soil and Water Conservation District took over as the administrator for the Missouri Ecotype Program replacing UMC. MDC funded the program for the 1999-2000 fiscal year.

### **2000**

The Missouri Ecotype program continued through August until funding was depleted. The program was continued under direction of Missouri Audubon Society and MDC in cooperation with the NRCS Plant Materials Center.

### **2001**

The Missouri Ecotype program is growing increase plots at Elsberry and also at the Charles Green Conservation area near Ashland, Missouri. A list of species in production at both sites is in Table #2 and plants released through the program in Table #3.

### **2002**

Missouri Department of Conservation took over as administrator of this study and is currently still funding the program with the aid of grants. The Missouri Ecotype program is continuing to increase plots at the Elsberry PMC and Green Conservation Area. New collections are being made of both old and new species.

Study No. 29I142G - Production of Native Missouri Ecotypes of Grasses, Legumes, and Forbs for Roadsides, Critical Areas, and All Other Vegetative Plantings Where Native Plants are Now Being Planted.					
Project Status			Table #2		
Common Name Genus/Species	Zone	Accession Number	Collection Date	Status of Increase Plot Planting (Where/When)	Status of Accession
<b>Big Bluestem</b> <i>Andropogon gerardii</i>	1	9079000	1997	In production PMC/1998	Released in 1999
<b>Butterfly Milkweed</b> <i>Asclepias tuberosa</i>	1	9079024	1997	In production PMC/1998	
	2	9079025	2001	In production Green CA/2002	
<b>Creamy Wild Indigo</b> <i>Baptisia bracteata</i>	1	9079088	2000	In production Green CA/2001	
	2	9079089	2000	In production Green CA/2001	
<b>Foxglove Beardstongue</b> <i>Penstemon digitalis</i>	1	9079064	1999	In production PMC/2000	
	2	9079065	2001	In production Green CA/2002	
<b>Grayhead Coneflower</b> <i>Ratibida pinnata</i>	1	9079060	1999 & 2000	In production PMC/2000, increased in 2001	
	2	9079061	2000	In production Green CA/2001	
<b>Gravelweed</b> <i>Verbesina helianthoides</i>	1	9079080	2001	In production Green CA/2002	
	2	9079081	2001	In production Green CA/2002	
<b>Horsemint</b> <i>Monarda fistulosa</i>	1	9079056	1999	In production PMC/2000	
	1	9079056	1999 & 2000	In production Green CA/2001(increase)	
	2	9079057	2000	In production Green CA/2001	
<b>Indiangrass</b> <i>Sorghastrum nutans</i>	1	9079036	n/a	Not in production on PMC Commercial production only	Released in 1999
	2	9079037	n/a	Not in production on PMC Commercial production only	Released in 1999

<b>Common Name</b>		<b>Accession</b>	<b>Collection</b>	<b>Status of Increase Plot</b>	<b>Status of</b>
<b>Genus/Species</b>	<b>Zone</b>	<b>Number</b>	<b>Date</b>	<b>Planting (Where/When)</b>	<b>Accession</b>
<b>Largeleaf Wild Indigo</b> <i>Baptisia alba</i>	1	9079092	2000	In production Green CA/2001	
	2	9079093	2000	In production Green CA/2001	
<b>Little Bluestem</b> <i>Schizachrium scoparium</i>	1	9079004	1996 & 1997	In production PMC/1997 (increase 1998)	Released in 1999
	2	9079005	1996 & 1997	In production PMC/1997	
	3	9079006	1996 & 1997	In production PMC/1997	
<b>New England Aster</b> <i>Aster oblongifolius</i>	1	9079103	2001	In production Green CA/2002	
<b>Pale Purple Coneflower</b> <i>Echinacea pallida</i>	1	9079032	1997	In production PMC/1998	Released in 2001
	2	9079033	1998	In production PMC/1999	Released in 2001
<b>Prairie Blazing Star</b> <i>Liatris pycnostachya</i>	1	9079020	1998	In production PMC/1999	
	2	9079021	2001	In production Green CA/2002	
<b>Prairie Coreopsis</b> <i>Coreopsis palmata</i>	1	9079028	1997	In production PMC/1998	Released in 2001
	2	9079029	1998	In production PMC/1999	Released in 2001
<b>Purple Milkweed</b> <i>Asclepias purpurascens</i>	1	9079114	2001	In production Green CA/2002	
<b>Purple Prairie Clover</b> <i>Dalea purpurea</i>	1	9079048	2000	In production PMC/2001	
	1	9079048	2000	In production Green CA/2001	
	2	9079049	2000	In production PMC/2001	
<b>Rough Blazing Star</b> <i>Liatris aspera</i>	1	9079068	1999	In production PMC/2000	
	2	9079069	2000	In production Green CA/2001	
<b>Roundhead Bushclover</b> <i>Lespedeza capitata</i>	1	9079008	1999	In production PMC/2000	
	2	9079009	2000	In production Green CA/2001	

<b>Common Name</b>		<b>Accession</b>	<b>Collection</b>	<b>Status of Increase Plot</b>	<b>Status of</b>
<b>Genus/Species</b>	<b>Zone</b>	<b>Number</b>	<b>Date</b>	<b>Planting (Where/When)</b>	<b>Accession</b>
<b>Showy Goldenrod</b> <i>Solidago speciosa</i>	1	9079110	2001	In production Green CA/2002	
<b>Sideoats</b> <i>Bouteloua curtipendula</i>	1	9079072	2001	In production PMC/2002	
	2	9079073	2001	In production PMC/2002	
<b>Stiff Goldenrod</b> <i>Solidago rigida</i>	1	9079107	2001	In production Green CA/2002	
	2	9079107	2001	In production Green CA/2002	
<b>Tall Dropseed</b> <i>Sporobolus compositus</i>	1	9079040	1998	In production PMC/1999	Released in 2001
	2	9079041	2001	In production PMC/2002	
<b>Tall Tickseed</b> <i>Coreopsis tripteris</i>	1	9079076	2000	In production Green CA/2001	
<b>Tick Trefoil</b> <i>Desmodium canadense</i>	1	9079012	1997	In production PMC/1998	
<b>Virginia Wild Rye</b> <i>Elymus virginicus</i>	1	9079044	1998	In production PMC/1999	Released in 1999
<b>White Prairie Clover</b> <i>Dalea candida</i>	1	9079052	2000	In production PMC/2001	
	1	9079052	2000	In production Green CA/2001	
	2	9079053	2000	In production Green CA/2001	
<b>White Wand Beardtongue</b> <i>Penstemon tubifloris</i>	2	9079101	2001	In production Green CA/2002	

29I142G - Missouri Native Ecotype Collection								Table #2
Missouri Ecotypes at PMC								
Item	Common Name Zone #	Accession No.	Field No.	Plot Size	Disc.	Stand	Date Planted	Clean Seed
				sq. ft. or ft X ft.	no. of plants	Percent		Bulk/no test in pounds
1	big bluestem MO Zone 1	9079000	7f	25x215		85	5/19/98	10.4
2	butterfly milkweed MOZ 1	9079024	1v	7x100	500	80	5/98-99	1.3
3	desmodium MO Zone 1	9079012	14i	25x110	1500	90	5/27/98	2.2
4	grayhead coneflower Z1	9079060	1	20x75	500	95	5/00	3.6
5	grayhead coneflower Z1	9079060	2	12x175	1000	98	5/01	small amount inc. w above
6	little bluestem MO Zone 1	9079004	4b	30x110		95	Apr-97	19.5
6a	little bluestem MO Zone 1	9079004	6a	35x50		90	5/19/98	mixed w above
7	little bluestem MO Zone 2	9079005	12b	5x90		90	4/29/97	1.3
8	little bluestem MO Zone 3	9079006	14e	20x130		90	4/29/97	10.6
9	horsemint MO Zone 1	9079056	6a	9x30	5	<5	5/11/00	27gms
10	pale purple coneflower Z 1	9079032	1f	18x75	700	75	12/17/98	1.8
11	pale purple coneflower Z 2	9079033	14L	22x100	1000	80	5/18/99	5.8
12	Penstemon digitalis Z 1	9079064	6a	10x110	500	85	6/6/00	5.8
13	prairie blazing star Z 1	9079020	2g	10x120	250	60	5/10/99	3.9
14	prairie coreopsis MO Z 1	9079028	1w	15x80	500	95	5/18/98	5.4
15	prairie coreopsis MO Z 2	9079029	14m	25x40	500	90	5/20/99	5.8
16	purple prairie clover MO Z1	9079048	9	25x80	100-15	<20	May-01	0
17	Rough blazing star MO Z1	9079068	14k	10x40	150	75	5/24/00	5.6
18	roundhead bush clover Z+E	9079008	11h	32x100	2000	60	4/00	3.3
19	tall dropseed MO Zone 1	9079040	7f	20x240		80	5/27/99	14.3
20	Virginia wildrye MO Z 1	9079044	7d	15x475		90	3/30/99	43.5
21	White prairie clover MO Z1	9079052	2	12x175	900	90	5/01	1
<b>Not in Production</b>								
22	Indiangrass MO Zone 1	9079036						
23	Indiangrass MO Zone 2+B	9079037						

								Table #2- cont.	
Missouri Ecotypes at Charles Greer Conservation Area									
		Accession	Field			Percent			
Item	Common Name	Zone #	No.	No.	Plot Size	Disc.	Stand	Date Planted	Clean Seed
					sq. ft. or ft X ft.	no. of plants			Bulk/no test in pounds
1	Largeleaf wild indigo 1					620		5/01	
2	largeleaf wild indigo 2					520		5/01	
3	Creamy longbract wild indigo 1					130		5/01	
4	Creamy longbract wild indigo 2					150		6/01	
5	White prairie clover 1				7x90	0		6/01	
6	White prairie clover 2					400		6/01	
7	Purple prairie clover 1				7x135	900		6/01	
8	Purple prairie clover 2					400		6/01	
9	Tall tickseed				7x250	1200	95	5/01	
10	Roundhead bush clover					500		5/01	
11	Rough blazing star				7x200	1400		6/01	
12	Horsemint 1				7x180	1400	100	5/01	
13	Horsemint 2				7x180	1200	100	6/01	
14	Grahead coneflower				7/220	1020	100	6/01	

Study 29I142G - Missouri Ecotype Releases						Table #3
Releases from the Elsberry Plant Materials Center						
Scientific Name	Release Name	Common Name	Accession Number	Cooperating Agency(ies)	Type of Release	Year of Release
<i>Elymus virginicus</i> L.	Northern MO	Virginia wildrye	9079044	MOPMC,UMC,MDC,MODOT	N	1999
<i>Sorghastrum nutans</i> (L) Nash.	Northern MO	indiangrass	9079036	MOPMC,UMC,MDC,MODOT	N	1999
<i>Andropogon gerardii</i> Vitman	Northern MO	big bluestem	9079000	MOPMC,UMC,MDC,MODOT	N	1999
<i>Sorghastrum nutans</i> (L) Nash.	Western MO	indiangrass	9079037	MOPMC,UMC,MDC,MODOT	N	1999
<i>Schizachyrium scoparium</i> , Michx.	Northern MO	little bluestem	9079004	MOPMC,UMC,MDC,MODOT	N	1999
<i>Sporobolus compositus</i> var. <i>compositus</i>	Northern MO	tall dropseed	9079040	MOPMC, MDC, NAS	N	2001
<i>Coreopsis palmata</i>	Northern MO	prairie coreopsis	9079028	MOPMC, MDC, NAS	N	2001
<i>Coreopsis palmata</i>	Western MO	prairie coreopsis	9079029	MOPMC, MDC, NAS	N	2001
<i>Echinacea pallida</i>	Northern MO	pale purple coneflower	9079032	MOPMC, MDC, NAS	N	2001
<i>Echinacea pallida</i>	Western MO	pale purple coneflower	9079033	MOPMC, MDC, NAS	N	2001
Cooperating Agencies: MOPMC=Missouri Plant Materials Center; UMC=University of Missouri at Columbia; MDC=Missouri Department of Conservation; MODOT=Missouri Department of Transportation; NAS=National Audubon Society-Audubon Missouri;. Grow Native.						
N= native releases; collected within the USA, occurring naturally in the USA. Generally refers to a plant which occurs naturally in a particular region, state ecosystem or habitat without direct or indirect human activity.						
Nat.=naturalized releases; collected from a population within the USA, but were originally introduced to the USA sometime in the past.						
I=introduced; means that the original collection from which the release was made was not from within the USA.						

**Study: 29I143G**

**Study Title: Seed Coating/Seeding Rates Study**

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

There is little information available comparing coated seed, versus non-coated seed, and various seeding rates of commonly used forage species used in the Midwest region. Studies done have been short lived (one or two years) and have looked only at emergence, plants at the end of the seedling year, or plants at the end of the first year following seeding.

Evaluations will be made on emergence, stems at the end of the seeding year, and stems at the end of the first through the fourth year following planting. The study will be repeated for five consecutive planting seasons to compensate for changes in yearly weather patterns.

**Problem:**

There is a need to compare coated seed to non-coated seed for selected legumes to determine if a significant difference exists. Disagreement of seeding rates between coated versus non-coated legume seed is quite common. The results of this study could improve on the seeding rate recommendations for legume species being tested.

Can seeding rates of selected legumes and forage grasses be reduced to one-half the current rate or increase to one and a half times the current rate and provide similar results in long term stand density. Selected grass/legume species will be monitored for the emergence date, emergence density, and stand density.

**Objective:**

The objectives of this project is to determine if a significant difference exists between coated versus non-coated seed of selected legume species and determine if the seeding rates of selected legume and forage grasses can be reduced or increased from current rates and provide the same results in stand density.

**Location:**

Selected field on the Freeman Farm at Lincoln University, Jefferson City, Missouri.

- A. Description: Township 44N, Range 10N, and Section 19
- B. MLRA: 115
- C. Soils: Grable silt loam

## Procedure:

### A. Assembly of Materials:

1. A list of species to be tested was developed and approved by cooperators. (See Table #1)
2. Species that were coated, both coated and uncoated lots, were provided by Seedbiotics and CelPril. Grass seed and other non-coated species were provide by USDA-Natural Resources Conservation Service, Plant Materials Center. (See Table #1)

### B. Planting Plan:

1. Plot Design: Randomized split plot design with four replications. (See Table #2 and #3) This study is planned to last a total of ten years.
2. Plot size: (See Table #3)
  - a. Length: 20 feet; 20' between reps.
  - b. Width: 15 feet; 21 rows: 30' between blocks
  - c. Redtop and tall fescue between blocks.
3. Seeding Method and Rate: (See Table #4)

The plots will be seeded with a plot seeder. Seeding rates will be .5, 1.0, and 1.5 X those listed in the current Pasture and Hayland Planting Specifications, NRCS MOFOTG, March, 1997.

### 4. Dates of Establishment:

Plots #1 - #13 (Legume Plots)	Planted	5/5/98	4/13/99	4/19/00
Plots #14-#19 (Cool Season Grass Plots)		4/23/98	4/13/99	4/20/00
Plot #20 (Untr / GT* Eastern Gama Plot)		3/26/98	4/21/99	4/27/00
Plot #21 (Wet Tr Eastern Gama Plot)		4/23/98	4/21/99	4/27/00
Plots #22&#23 (Warm Season Grass Plots)		5/5/98	4/21/99	4/27/00

- \* Plot #20 was a dormant planting using untreated seed in 1998.
- Germtech treated seed was used in 1999 and 2000.

### C. Evaluation Measurement:

1. Climatic Data: The data from the nearest weather station will be used to report precipitation and temperatures. (See Table # 7)

2. Measurements: 1998-2002 (See Tables # 5 & # 6)
  - a. Emergence dates; when 25 plants have emerged.
  - b. Emergence density; conduct emergent density counts by counting the number of plants/foot of row, three counts/plot X number of plots, 3 weeks after planting for legumes, 4 weeks after planting for cool season grasses and 5 weeks after planting for warm season grasses.
  - c. Stem counts; conduct stem counts as same procedure using plants/foot of row, 3 counts/plot X plots at the end of their first, second, third, fourth, and fifth growing seasons (for the three seeding rates and coated versus non-coated seed).

### **Cooperators:**

The following is a listing of cooperators involved with this study: Lincoln University, Jefferson City, Missouri; Seedbiotics, CelPril, and USDA-Natural Resources Conservation Service, Plant Materials Center, Elsberry, Missouri.

### **Discussion:**

#### **1998**

Signatures of all cooperators with the study were received by March of 1998. Seed lots were received for accessions to be planted and new seed tests were secured when necessary.

This study was seeded with a cone type plot planter for all species except eastern gamagrass that was planted with a corn planter using soybean seedcups. Due to a planter malfunction, the legume plots were replanted in the YEAR TWO block and the warm season plots are planted partially in the YEAR ONE block and YEAR TWO block (see Table #2).

The study consists of two comparisons, coated verses non coated seed, and three different seeding rates.

The comparison of coated verses non-coated seed was done by planting equal bulk rates. For example, if a bag of seed has a test of 95% purity and 90% germination, it is 85.5% pure live seed (PLS). If you want to plant 10# PLS per acre you need to plant 11.7# (10 / .855) BULK. A 50# bag of seed with this test has 95% (47.5#) seed and 5% (2.5#) other (dirt, chaff, weed seed, etc.). The 95% seed has a germination of 90% so the seed portion contains 42.75# Pure Live Seed (PLS) and 4.75# non viable seed.

When seed is coated, the coating generally accounts for 25 to 40 percent of the weight according to the seed industry that coats seed. If the above bag of seed was coated and 30% of the total weight was coating, the composition of the coated and uncoated seed would be as follows:

	<b>Coating</b>	<b>Pure Live Seed</b>	<b>Non-viable Seed</b>	<b>Other (Dirt, etc.)</b>
50# coated seed	15# (30%)	29.9# (59.8%)	3.3#	1.8#
50# uncoated seed	0#	42.75# (85.5%)	4.75#	2.5#

When coating is added to seed, the amount of pure live seed goes down and that weight is replaced by coating. This coating is comprised of compounds that are designed to aid in seed germination and seedling development. Discussion from the seed industry suggests that coated seed is equal to or more beneficial than the loss of pure live seed. In a situation where 10.0# PLS is recommended, using the above test of 85.5% PLS, a bulk seeding rate of 11.7# of seed is required. To get 10.0# PLS of the above coated seed you would need 16.7#. The objective of this part of the study is to determine if 11.7# of the coated seed is equal to or better than 11.7# of the uncoated seed.

This study compared bulk weights of coated and uncoated seed. Using the above rates and seed tests, the comparison is as follows:

Uncoated seed    11.7# Bulk Rate    containing 10.0# Pure Live Seed

Compared to:

Coated seed        11.7# Bulk Rate    containing 7.0# Pure Live Seed and 3.5# coating

The seeding rate portion of the study uses a split plot design (see Table #3) to compare different rates of all species in the study including both the coated and uncoated seed. Seeding rates were calculated as both pounds per acre and pure live seeds per square foot. Seed size and seeding rates vary considerably between species (see Table #4). Pure live seed per square foot is not calculated for coated seed because the exact percentage of coating is not known. It is generally about one third. Measurements of emergence density and cover density were done on a row foot basis rather than square foot because the plots were seeded in rows rather than broadcast. Seeding rates can be converted from pure live seed per square foot (100 sq. ft per plot) to row foot (140 row foot per plot) by using a conversion factor of .714 to determine how many seeds it took in correlation to the emergence and cover density evaluations.

Weed control on the plots became somewhat of a problem by mid season due to wet weather. The ladino seed had an incorrect test so both coated and uncoated plots only had about a third of the intended rate but the ratios stayed the same.

The 1998 data from the legume plots indicate most of the coated plots were about the same or slightly better than the uncoated at the lower (.5 full rate) and full seeding rates. The higher seeding rate(1.5 X

full rate) had about the same or slightly lower emergence density. It also varied between species. Treated seed of the eastern gamagrass showed a considerable increase over untreated seed.

Differences in the seeding rates was also quite evident in the data but not always as much as expected. The 1.5 seeding rate was not always a lot better than the half rate. This indicates the amount of seed may not be the problem of a week stand.

## 1999

Data taken in 1998 and 1999 is averaged in Tables #5 and #6. The data showed a significant difference between coated and uncoated for 1999 emergence density that is an important criterion. Coated alfalfa is equal or slightly better at standard rates. Red clover is better at the lower rates but the other rates vary both ways. Coating did not show improvement for birdsfoot trefoil and in some cases was a disadvantage. Coated ladino clover seed was not equal to uncoated and in some cases was a disadvantage. The summary did not show any significant difference between coated and uncoated seed in 1998 indicating that for this year the coating was just as good as having the additional seed.

## 2000

This study was designed for plots to be established for five consecutive years. Local weather patterns are quite variable from year to year and 1998 through 2000 were no exceptions (See Table #7). 1998 was dryer than average in the spring, was well above average during June and July and barely rained at all in August. Weed control became a problem during the summer. 1999 was about the opposite, starting out wetter than average causing ponding on some of the plots and then becoming very dry during the summer. Year 2000 started out extremely dry with the drought of 1999 continuing through April 2000. April 2000 was one of the driest April's on record at 0.84" precipitation. The rest of the growing season in 2000 was wet.

Data for each year was averaged and can be found in Tables #5 and #6.

Table #5 is spring evaluations that are three-year averages of the following evaluation criteria.

1. Days to Emerge: This is the number of days it took for 25 seedlings to emerge in each plot.
2. Emergence Density: This is a count of how many plants per row foot emerged three weeks after planting for legumes, four weeks after planting for cool season grasses, and five weeks after planting for warm season grasses.
3. Percent Stand: This is a visual rating of the percent of the plot that has complete rows of plants and is done at the same time as emergence density.

Table #6 is fall evaluations that are averages of the following evaluation criteria.

4. Cover Density: This is a count of stems per row foot done at the end of the growing season.
5. Percent Cover: This is a visual rating of the percent of the plot that has groundcover from the planted species.

## SUMMARY

The following summary will note differences and relative indifferences between coated and uncoated seed and also seeding rates for each individual species. Differences between species will not be compared.

### **Alfalfa**

(1) Days to Emerge - There was no advantage or disadvantage of coated seed or the different seeding rates.

(2) Emergence Density - The coated seed was less than the uncoated. The higher seeding rate did not always result in many more plants with the coated seed. Higher seeding rates with uncoated seed resulted in more plants but not in proportion to the amount of extra seed planted.

(3) Percent Stand - Percent stand was about equal for the uncoated and Celpril coated but the Seed Biotics was less. Percent stand increased with increase in seeding rate for both the coated and uncoated.

(4) Cover Density - Stems per row foot was less at the end of the first year for the coated seed. By the end of the second year there was very little difference between coated and uncoated except the full standard rate Seed Biotics which was considerably higher. By the end of the third year, there was little difference between the uncoated and the Celpril coated and the Seed Biotics coated was slightly higher. Cover density increased with an increased seeding rate at the end of the first year, but showed no difference by the end of the second year.

(5) Percent Cover - Percent cover was about the same for the uncoated and Seed Biotics coated but Celpril coated was less. The differences became less the second year after planting. The differences in the seeding rates are inconsistent.

Overall the data shows no clear advantage or disadvantage for coated seed after the establishment year. Coated seed plots were thinner the establishment year. The higher seeding rates only showed an advantage the first year.

### **Red Clover**

(1) Days to Emerge - There was no advantage or disadvantage of coated seed or the different seeding rates.

(2) Emergence Density - The Celpril coated seed was about equal to the uncoated but the Seed Biotics coated seed was less. The higher seeding rates did produce more plants but not close in proportion to the extra seed.

(3) Percent Stand - Seed Biotics coated seed was less than the uncoated at all rates and Celpril coated seed was less only at the standard rate. There was no increase when the seeding rate increased from full standard rate to 1.5 rate except the Celpril coated seed, which had no increase from the half rate to the full rate.

(4) Cover Density - Cover density of the coated seed plots at the end of the establishment year was less than the uncoated plots except for the high rate Celpril. By the end of the second year, the coated plots at all seeding rates were less than the uncoated. The higher seeding rates produced more stems per row foot the year of establishment but by the second year, only the uncoated seed indicated an advantage to having more seed.

(5) Percent cover - The uncoated seed provided higher percent cover than the coated except at the highest rate which was about the same. By the end of the second year, the differences were less. Percent cover was not consistently higher with higher seeding rates.

Overall the data indicates the uncoated seed did better than the coated. By the end of the second year, the only advantage to higher seeding rates was with uncoated seed.

### **Birdsfoot trefoil**

(1) Days to Emerge - There was no advantage or disadvantage of coated seed or the different seeding rates.

(2) Emergence Density - Emergence densities were lower for coated seed compared to uncoated except the Seed Biotics coated at the standard rate. Higher seeding rates did result in more plants but not in proportion to the extra seed planted.

(3) Percent Stand - The uncoated and Celpril plots were similar and somewhat higher than the Seed Biotics plots. There was usually only a small increase in percent stand with each increase in seeding rates.

(4) Cover Density - At the end of the establishment year cover density was higher for the uncoated seed. By the end of the second year there was very little difference between the coated and uncoated for the low and full rates. The uncoated was a little higher in the high rate. Increases in cover density was apparent when seeding rates went from low to full. There was very little difference and sometimes a slightly negative response when seeding rates went from the full rate to the high rate.

(5) Percent Cover - At the end of the establishment year there was very little difference between the uncoated and Celpril coated seed. Seed Biotics seed was less. By the end of the second year the Celpril plots were better than the uncoated at the full and high rates. The Celpril high rate plot increased

to equal the uncoated. There was only a small increase in percent cover with increased seeding rates in comparison to the extra seed.

Overall the data indicates the uncoated seed did better the establishment year but by the end of the second year there was very little difference between the uncoated and the Celpril coated. The Seed Biotics coated did not do as well. Response to higher seeding rates was minimal in comparison to the extra seed.

### **Ladino clover**

(1) Days to Emerge - There was no advantage or disadvantage of coated seed or the different seeding rates.

(2) Emergence Density - The coated seed was less than the uncoated. Higher seeding rates resulted in more plants but not close to the same increase as the increased seed applied.

(3) Percent Stand - Seed Biotics coated seed was higher at the full and high rates than the uncoated and Celpril coated. There was good response between the half rate to the full rate and no response between the full rate and the high rate for the uncoated and the Seed Biotics coated. Celpril had more response between the half rate and the full rate.

(4) Cover Density - The uncoated seed had higher cover density at the end of the first year. By the end of the second year the coated seed had higher cover density than the uncoated at the half rate and high rate but not at the standard full rate. Cover density in general had an increase as seeding rates increased but was very small in comparison to the amount of extra seed.

(5) Percent Cover - The uncoated seed was higher than the coated seed at the end of the first year. By the end of the second year there was very little difference. There was slight to moderate increase in percent cover with the increase in seeding rates at the end of the first year. By the end of the second year there was no difference between seeding rates.

Overall the data does not indicate an advantage or disadvantage to coated seed by the end of the second year. Uncoated seed rates were higher at the end of the first year. By the end of the second year, there was little difference between seeding rates.

### **Annual Lespedeza**

(1) Days to Emerge - There was little difference between seeding rates.

(2) Emergence Density - Emergence density was low for all rates when compared to the amount of seed planted. The response from the half rate to the full standard rate was much greater than the response from the full standard rate to the high rate.

(3) Percent Stand - There was a slight increase with increased seeding rates.

(4) Cover Density - Cover Density was extremely low for all seeding rates.

(5) Percent Cover - Percent Cover was extremely low for all seeding rates and lowest for the standard full rate.

Overall the data indicates the amount of seed was not the problem with stand establishment but with other unknown factors.

### **Tall Fescue**

(1) Days to Emerge - The endophyte infected variety emerged approximately two days sooner than Endophyte free varieties.

(2) Emergence Density - The endophyte infected variety had much higher emergence density than the endophyte free varieties. Emergence density increased considerably between the low and standard rates but not nearly as much between the full standard rate and the high rate.

(3) Percent Stand - The endophyte infected was extremely higher than the endophyte free. The difference in seeding rates was not very evident.

(4) Cover Density - Cover density was extremely greater for the endophyte infected variety than the endophyte free variety. The different seeding rates resulted in little difference in cover density in the first or second years.

(5) Percent Cover - The endophyte infected was extremely higher than the endophyte free. There was little difference between seeding rates at the end of the establishment year and none after the second year.

Overall the endophyte infected variety established much better and developed into an excellent stand and the endophyte free varieties did not. The difference in seeding rates was not very evident after the first year.

### **Orchardgrass**

(1) Days to Emerge - There was little difference between seeding rates.

(2) Emergence Density - Emergence density increased along with seeding rates but at a slower rate.

(3) Percent Stand - There was no difference in percent stand between the low and standard full rate and a slight increase in the high rate.

(4) Cover Density - There was a small increase from the low to full standard to high rates the first year. By the end of the second year there was no difference between the low and standard rates and a small decline in the high rate.

(5) Percent Cover - The percent cover data showed small increases as the rates increased at the end of the first and second years.

Overall the data indicates small differences between seeding rates. Even at the high seeding rate, the stand was not filled in a lot more than the low rate.

### **Smooth Brome**

- (1) Days to Emerge - There was no difference between seeding rates.
- (2) Emergence Density - There was very little difference between the three seeding rates.
- (3) Percent Stand - There was moderate increases with increased seeding rates.
- (4) Cover Density - There was little difference between the low and full standard rates and more of an increase between the full standard and high rate the end of the first year. By the end of the second year there was little difference between rates.
- (5) Percent Cover - Percent cover was low at the end of the first year with very little difference between rates. By the end of the second year the percent cover was high but still very little difference between seeding rates.

Overall the data indicates very little difference between seeding rates.

### **Timothy**

- (1) Days to Emerge - The lower seeding rate plots took longer to emerge.
- (2) Emergence Density - There was a small increase between the low and standard full rate and a larger increase between the standard full and high rate.
- (3) Percent Stand - There was very small increases in proportion to the increase of seed.
- (4) Cover Density - There was almost no difference between seeding rates the year of establishment but cover density increased along with seeding rates the year after planting.
- (5) Percent Cover - There was no difference between low and standard full rate and a small increase between standard full and high rate for the end of the year of planting. By the end of the second year the trends were similar only much less. This species is decreasing at all rates.

Overall the data indicates the seeding rate is not the problem with poor stands.

### **Canada Wildrye**

- (1) Days to Emerge - There was no difference between the different seeding rates.
- (2) Emergence Density - There is a large difference between the low and standard full seeding rates. There is little difference between the standard full and the high rate.

(3) Percent Stand - There is twice the difference between the low and standard full rate as there is between the standard full and the high rate.

(4) Cover Density - There was very little difference between rates at the end of the first year. By the end of the second year there was a considerable difference between the low and standard rates but little difference between the standard and high rates.

(5) Percent Cover - Percent cover increased along with seeding rates but much more between the low and standard than the standard and high for both the first and second years..

Overall the data indicates that additional seed above the standard seeding rate did not improve the stand very much.

### **Eastern Gamagrass**

(1) Days to Emerge - Plot 21 was wet treated stratification. Plot 20 was a dormant untreated planting in 1998 and Germtech treated seed was used in 1999 and 2000. Emergence was consistent between rates but quite variable between years. The Germtech seed came up quicker in 1999 but could not be found in 2000.

(2) Emergence Density - Emergence density was similar for Germtech and wet treated seed and higher than the one year data for the untreated dormant planting. There was very little difference between seeding rates.

(3) Percent Stand - The three year average of the wet treated for percent stand was higher than the two year average for the Germtech. The one year data for the untreated dormant was higher than the averages for either treated but lower when compared to the same one year data for the wet treated.

(4) Cover Density - Cover density at the end of the first year averaged about the same for the treated seed except for the standard full rate wet treated which was higher. Treated seed averaged higher than the one year data for the untreated dormant planting. By the second year the high rate Germtech was better and the standard rate wet treated was better.

(5) Percent Cover - At the end of the first year, percent cover is highest for the wet treated seed. By the end of the second year, the one year data for the dormant planting is higher than the Germtech or the two year average of the wet treated, but the dormant planting is lower when just compared to the same year planting data for the wet treated. Seeding rates show little advantage to the higher rates and some disadvantage to the lower rates.

Overall the data is quite variable when comparing one year's data of a dormant seeded planting to two years data of Germtech seed to three years data of wet treated seed. The wet treated seed is doing somewhat better.

## **Switchgrass**

- (1) Days to Emerge - There was no difference between the different seeding rates.
- (2) Emergence Density - The three year average of emergence density increased as the seeding rates increased.
- (3) Percent Stand - There was no difference in percent stand for the different seeding rates.
- (4) Cover Density - There was very little difference between seeding rates for the three year average at the end of the establishment year. By the end of the second and third years, the higher seeding rates actually produced fewer stems.
- (5) Percent Cover - there is a difference between the half rate and the full standard rate at the end of the first second and third years. There is very little difference between the full standard and high rate.

Overall the data indicates there is a difference between rates the first year during emergence. There continued to be a difference between the low and standard full rates during October evaluations but not between the standard full rate and the high rate. Seeding above the Standard full rate indicated no benefit.

## **Caucasian Bluestem**

- (1) Days to Emerge - There was no difference between different seeding rates.
- (2) Emergence Density - There was no difference between the low rate and standard full rate for the three year average. The high rate had a large increase in comparison to the other rates.
- (3) Percent Stand - Percent stand was highest at the low rate and the standard full rate and high rate was the same.
- (4) Cover Density - There was not large differences between rates.
- (5) Percent Cover - Percent cover was similar for the three seeding rates. The biggest difference was between the low and full standard rates at the end of the first year.

Overall the data does not indicate a consistent trend at this time.

## **Big Bluestem**

Big bluestem was only planted two years.

- (1) Days to Emerge - There was no difference in the seeding rates.
- (2) Emergence Density - There was a large difference between rates and the standard full rate was the highest.
- (3) Percent Stand - The standard full rate was the highest although all were low.

(4) Cover Density - Stems per row foot increased slightly with increased seeding rates at the end of the first year. At the end of the second year the results were mixed.

(5) Percent Cover - There was not a big difference in percent cover for the different seeding rates at the end of the first year. The standard full rate was somewhat higher. By the end of the second year the standard full rate was lower.

Overall the data does not indicate a consistent trend at this time.

## 2001

No additional plantings were made in 2001. The plots planted in 1999 and 2000 were evaluated for cover density (stems/row foot) and percent cover (visual observation).

## 2002

No additional plantings were made in 2002. The plots planted in 2000 were evaluated for cover density (stems/row foot) and percent cover (visual observation).

The data from this study is also being used to give an indication of stem density for each species. Stem density of a species at 100% cover density will be estimated by averaging the stem densities of all seeding rates and coatings in relation to the percent cover. Cover density will then be interpolated to reflect a 100% cover. See Table #1.

Study 29I143G – Seed Coat/Seeding Rates Study				Table #1
List of Species Evaluated				
Genus	Species	Common Name	Standard Full Seed Rate (MOFOTG March 1997)	Stems per sq. ft. at 100 % Stand
<i>Medicago</i>	<i>sativa</i>	Alfalfa	9.4# PLS/Ac	35
<i>Trifolium</i>	<i>pratense</i>	Red clover	7.6# PLS/Ac	101
<i>Lotus</i>	<i>corniculatus</i>	Birdsfoot trefoil	6.2# PLS/Ac	105
<i>Lespedeza</i>		Lespedeza (annual)	9.5# PLS/Ac	73
<i>Trifolium</i>	<i>repens</i>	Ladino clover	3.7# PLS/Ac	82
<i>Festuca</i>	<i>arundinacea</i>	Tall fescue	12.0# PLS/Ac	56
<i>Dactylus</i>	<i>glomerata</i>	Orchardgrass	5.2# PLS/Ac	74
<i>Bromus</i>	<i>inermis</i>	Smooth brome grass	10.0# PLS/Ac	46
<i>Phleum</i>	<i>pratense</i>	Timothy	3.9# PLS/Ac	73
<i>Elymus</i>	<i>canadensis</i>	Canada wildrye	10.0# PLS/Ac	55
<i>Tripsacum</i>	<i>dactyloides</i>	Eastern gamagrass	10.0# PLS/Ac	26
<i>Panicum</i>	<i>virgatum</i>	Switchgrass	5.9# PLS/Ac	56
<i>Bothriochloa</i>	<i>ischaemum</i>	Caucasian bluestem	3.1# PLS/Ac	147
<i>Andropogon</i>	<i>gerardii</i>	Big bluestem	10.0# PLS/Ac	89

STUDY 29I143G		PLOT LAYOUT		PLOT SIZE 15' X 20'		Table #2	
		----- county road -----		SUB-PLOT SIZE 5' X 20'			
		30'		445'			
	30'	REP 1	LEGUMES	20'	COOL S. G	20'	WARM S. G
		20'					30'
		REP 2	LEGUMES		COOL S. G		WARM S. G
200'		20'	YEAR	<b>ONE</b>			
		REP 3	LEGUMES		COOL S. G		WARM S. G
		20'					
		REP 4	LEGUMES		COOL S. G		WARM S. G
		30'					
		REP 1	LEGUMES		COOL S. G		WARM S. G
		20'					
		REP 2	LEGUMES		COOL S. G		WARM S. G
170'		20'	YEAR	<b>TWO</b>			
		REP 3	LEGUMES		COOL S. G		WARM S. G
		20'					
		REP 4	LEGUMES		COOL S. G		WARM S. G
880'		30'					
←	North	REP 1	LEGUMES		COOL S. G		WARM S. G
		20'					South →
		REP 2	LEGUMES		COOL S. G		WARM S. G
170'		20'	YEAR	<b>THREE</b>			
		REP 3	LEGUMES		COOL S. G		WARM S. G
		20'					
		REP 4	LEGUMES		COOL S. G		WARM S. G
		30'					
		REP 1	LEGUMES		COOL S. G		WARM S. G
		20'					
		REP 2	LEGUMES		COOL S. G		WARM S. G
170'		20'	YEAR	<b>FOUR</b>			
		REP 3	LEGUMES		COOL S. G		WARM S. G
		20'					
		REP 4	LEGUMES		COOL S. G		WARM S. G
		30'					
		REP 1	LEGUMES		COOL S. G		WARM S. G
		20'					
		REP 2	LEGUMES		COOL S. G		WARM S. G
170'		20'	YEAR	<b>FIVE</b>			
		REP 3	LEGUMES		COOL S. G		WARM S. G
		20'					
		REP 4	LEGUMES		COOL S. G		WARM S. G
		30'					
					* S. G. = SEASON GRASSES		

STUDY 29I143G - SEED COAT/SEEDING RATES STUDY																												Table # 3											
LEGUMES														planted 5/5/98																									
														Y E A R # 1																									
R E P # 1																																							
P#	4		10		7		1		11		2		9		5		13		8		3		12		6														
S#	2	3	1	2	1	3	2	1	3	2	1	3	3	1	2	3	2	1	2	1	3	1	3	2	1	3	2	2	1	3	3	2	1	2	1	3	3	2	1
R E P # 2																																							
P#	2		7		13		8		4		1		11		3		6		12		5		9		10														
S#	1	3	2	3	2	1	1	3	2	1	2	3	3	1	2	1	2	3	2	1	3	2	1	3	1	2	3	3	2	1	2	1	3	3	2	1	3	1	2
R E P # 3																																							
P#	9		12		8		3		10		2		13		7		4		11		5		1		6														
S#	1	3	2	2	1	3	3	1	2	3	1	2	1	2	3	3	2	1	3	2	1	3	2	1	1	2	3	1	3	2	1	3	2	1	2	3	2	1	3
R E P # 4																																							
P#	1		10		9		2		13		6		7		11		8		4		5		12		3														
S#	1	3	2	3	1	2	1	2	3	3	2	1	2	1	3	1	2	3	3	2	1	1	2	3	1	2	3	1	3	2	3	1	2	1	3	2	1	3	2
P# is Plot Number														Plot Size = 5' x 20'																									
S# is SubPlot Number														Subplot Size = 15' x 20'																									

Study 29I143G - Seed Coat/Seeding Rates Study														Table # 3 - continued																	
COOL SEASON GRASSES														WARM SEASON GRASSES																	
Y E A R # 1														Y E A R # 1 & 2																	
R	E	P	#																												
P #	16		18		14		17		15		19			23		21		22		20			99		98		99		98		
S #	2	1	3	2	1	3	1	2	3	2	3	1	3	2	1	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	
planted 4/23/98														planted 5/5/98 and 4/21/99																	
R	E	P	#																												
P #	16		18		17		14		19		15			21		20		23		22			98		98		99		99		
S #	3	2	1	1	2	3	3	1	2	1	2	3	3	1	2	3	2	1	3	2	1	2	3	1	3	1	2	3	1		
R	E	P	#																												
P #	16		19		17		18		14		15			20		22		21		23			98		99		98		99		
S #	2	1	3	3	1	2	2	3	1	1	2	3	2	1	3	2	1	3	3	2	1	3	3	2	1	2	3	1	2		
R	E	P	#																												
P #	19		15		17		18		16		14			20		22		21		23			98		99		98		99		
S #	3	2	1	2	1	3	2	1	3	3	1	2	3	1	2	1	3	2	1	3	3	2	1	2	3	1	1	2	3	2	1
P # is Plot Number														Plot Size = 5' x 20'																	
S # is SubPlot Number														Subplot Size = 15' x 20'																	

STUDY 29I143G - SEED COAT/SEEDING RATES STUDY															Table # 3 - continued																								
LEGUMES															Planted 4/13/99																								
Y E A R # 2																																							
R E P # 1																																							
P#	5		11			2			4			13		10		8		12		6		1		7		9		3											
S#	2	3	1	2	1	3	2	1	3	2	1	3	3	1	2	3	2	1	2	1	3	1	3	2	1	3	3	2	1	2	1	3	3	2	1				
R E P # 2																																							
P#	12		6			10			5			3		1		9		7		2		13		11		8		4											
S#	1	3	2	3	2	1	1	3	2	1	2	3	3	1	2	1	2	3	2	1	3	2	1	3	1	2	3	3	2	1	2	1	3	3	2	1	3	1	2
R E P # 3																																							
P#	6		4			13			1			7		9		5		11		8		12		3		10		2											
S#	1	3	2	2	1	3	3	1	2	3	1	2	1	2	3	3	2	1	3	2	1	3	2	1	1	2	3	1	3	2	1	3	2	1	2	3	2	1	3
R E P # 4																																							
P#	10		3			7			12			5		2		1		9		4		6		8		11		13											
S#	1	3	2	3	1	2	1	2	3	3	2	1	2	1	3	1	2	3	3	2	1	1	2	3	1	2	3	1	3	2	3	1	2	1	3	2	1	3	2
P# is Plot Number															Plot Size = 5' x 20'																								
S# is SubPlot Number															Subplot Size = 15' x 20'																								

STUDY 29I143G - SEED COAT/SEEDING RATES STUDY																	Table # 3 - continued												
COOL SEASON GRASSES										planted 4/13/99							WARM SEASON GRASSES							planted 4/21/99 and 5/5/98					
										Y E A R # 2							Y E A R # 1 & 2												
R E P # 1																													
P #										17 14 19 15 18 16							23 21 22 24 20							98 99 98 99 99					
S #										2 1 3 2 1 3 1 2 3 2 3 1 3 2 1 3 1 2							3 1 2 3 2 1 1 3 2 3 1 2 2 1 3												
R E P # 2																													
P #										19 15 18 14 17 16							24 20 23 22 21							99 99 98 98 99					
S #										3 2 1 1 2 3 3 1 2 1 2 3 3 1 2 3 2 1							3 1 2 3 2 1 3 2 1 2 3 1 1 3 2												
R E P # 3																													
P #										15 19 16 17 18 14							20 22 21 23 24							99 98 99 98 99					
S #										2 1 3 3 1 2 2 3 1 1 2 3 2 1 3 2 1 3							2 3 1 1 3 2 2 1 3 3 2 1 3 1 2												
R E P # 4																													
P #										18 16 15 19 14 17							24 22 21 23 20							99 98 99 98 99					
S #										3 2 1 2 1 3 2 1 3 3 1 2 3 1 2 1 3 2							2 3 1 1 2 3 2 1 3 3 2 1 1 2 3												
P # is Plot Number										Plot Size = 5' x 20'																			
S # is SubPlot Number										Subplot Size = 15' x 20'																			

Study 29I143G - Seed Coat/Seeding Rates Study				Table #4
Plot Number	Sub Plot Number	Forage - Seeds per LB - full seeding rate \4	Sub Plot Seeding Rates	PLS/square foot
1	1	Alfalfa 200,000 seeds/lb	.5 rate	21.6 PLS / Square foot
"	2	Alfalfa 9.4# / ac	1.0 rate	43.2 PLS / Square foot
"	3	Alfalfa	1.5 rate	64.8 PLS / Square foot
2	1	Alfalfa (Cel-coated) \1	.5 rate	\3
"	2	Alfalfa (Cel-coated)	1.0 rate	\3
"	3	Alfalfa (Cel-coated)	1.5 rate	\3
3	1	Alfalfa (S.B.-coated) \2	.5 rate	\3
"	2	Alfalfa (S.B.-coated)	1.0 rate	\3
"	3	Alfalfa (S.B.-coated)	1.5 rate	\3
4	1	Red clover 275,000 seeds/lb	.5 rate	24.0 PLS / Square foot
"	2	Red clover 7.6# / ac	1.0 rate	48.0 PLS / Square foot
"	3	Red clover	1.5 rate	72.0 PLS / Square foot
5	1	Red clover (Cel-coated)	.5 rate	\3
"	2	Red clover (Cel-coated)	1.0 rate	\3
"	3	Red clover (Cel-coated)	1.5 rate	\3
6	1	Red clover (S.B.-coated)	.5 rate	\3
"	2	Red clover (S.B.-coated)	1.0 rate	\3
"	3	Red clover (S.B.-coated)	1.5 rate	\3
7	1	Birdsfoot trefoil 75,000 seeds/lb	.5 rate	26.7 PLS / Square foot
"	2	Birdsfoot trefoil 6.2# / ac	1.0 rate	53.4 PLS / Square foot
"	3	Birdsfoot trefoil	1.5 rate	80.1 PLS / Square foot
8	1	Birdsfoot trefoil (Cel-coated)	.5 rate	\3
"	2	Birdsfoot trefoil (Cel-coated)	1.0 rate	\3
"	3	Birdsfoot trefoil (Cel-coated)	1.5 rate	\3
9	1	Birdsfoot trefoil (S.B.-coated)	.5 rate	\3
"	2	Birdsfoot trefoil (S.B.-coated)	1.0 rate	\3
"	3	Birdsfoot trefoil (S.B.-coated)	1.5 rate	\3
10	1	Ladino clover 871,650 seeds/lb	.5 rate	37.0 PLS / Square foot
"	2	Ladino clover 3.7# PLS/Ac	1.0 rate	74.0 PLS / Square foot
"	3	Ladino clover	1.5 rate	111.1 PLS / Square foot

\1 CelPril coated

\2 Seed Biotics coated

\3 See discussion 1998

\4 Rates as per NRCS MOFOTG March 1997

Study 29I143G - Seed Coat/Seeding Rates Study				Table #4 - continued
Plot	Sub Plot	Forage - Seeds per LB	Sub Plot Seeding	
Number	Number	- full seeding rate \4	Rates	PLS/square foot
11	1	Ladino clover (Cel-coated)	.5 rate	\3
"	2	Ladino clover (Cel-coated)	1.0 rate	\3
"	3	Ladino clover (Cel-coated)	1.5 rate	\3
12	1	Ladino clover (S.B.-coated)	.5 rate	\3
"	2	Ladino clover (S.B.-coated)	1.0 rate	\3
"	3	Ladino Clover (S.B.-coated)	1.5 rate	\3
13	1	Lespedeza (annual)	.5 rate	22.6 PLS / Square foot
"	2	Lespedeza (annual) 9.5# PLS / Ac	1.0 rate	45.3 PLS / Square foot
"	3	Lespedeza (annual)	1.5 rate	67.9 PLS / Square foot
14	1	Tall fescue(end. inf.) 227,000 seeds/lb	.5 rate	31.3 PLS / Square foot
"	2	Tall fescue(end. inf)12.0# PLS / Ac	1.0 rate	62.5 PLS / Square foot
"	3	Tall fescue (endophyte infested)	1.5 rate	93.8 PLS / Square foot
15	1	Tall fescue (endophyte free)	.5 rate	31.3 PLS / Square foot
"	2	Tall fescue (endophyte free)	1.0 rate	62.5 PLS / Square foot
"	3	Tall fescue (endophyte free)	1.5 rate	93.8 PLS / Square foot
16	1	Orchardgrass 654,000 seeds/lb	.5 rate	39.0 PLS / Square foot
"	2	Orchardgrass 5.2# PLS / Ac	1.0 rate	78.1 PLS / Square foot
"	3	Orchardgrass	1.5 rate	117.1 PLS /Square foot
17	1	Smooth bromegrass 136,000 seeds/lb	.5 rate	15.6 PLS / Square foot
"	2	Smooth bromegrass 10.0# PLS / Ac	1.0 rate	31.2 PLS / Square foot
"	3	Smooth bromegrass	1.5 rate	46.8 PLS / Square foot
18	1	Timothy 1,300,000 seeds/lb	.5 rate	58.2 PLS / Square foot
"	2	Timothy 3.9# PLS / Ac	1.0 rate	116.4 PLS /Square foot
"	3	Timothy	1.5 rate	174.6 PLS /Square foot
19	1	Canada wildrye 115,000 seeds/lb	.5 rate	13.2 PLS / Square foot
"	2	Canada wildrye 0.0# PLS / Ac	1.0 rate	26.4 PLS / Square foot
"	3	Canada wildrye	1.5 rate	39.6 PLS / Square foot
20	1	Eastern gamagrass (d. tr) 7,500 seeds/lb	.5 rate	0.9 PLS / Square foot
"	2	Eastern gamagrass (d. tr) 10.0 # PLS seeds/ac	1.0 rate	1.7 PLS / Square foot
"	3	Eastern gamagrass (drytreated)	1.5 rate	2.6 PLS / Square foot

\1 CelPril coated

\2 Seed Biotics coated

\3 See discussion 1998

\4 Rates as per NRCS MOFOTG March 1997

Study 29I143G - Seed Coat/Seeding Rates Study				Table #4-continued	
Plot Number	Sub Plot Number	Forage - Seeds per LB - full seeding rate \4	Sub Plot Seeding Rates	PLS/square foot	
21	1	Eastern gamagrass (wettreated)	.5 rate	0.9 PLS / Square foot	
"	2	Eastern gamagrass (wettreated)	1.0 rate	1.7 PLS / Square foot	
"	3	Eastern gamagrass (wettreated)	1.5 rate	2.6 PLS / Square foot	
22	1	Switchgrass 389,000 seeds/lb	.5 rate	26.3 PLS / Square foot	
"	2	Switchgrass 5.9# PLS / Ac	1.0 rate	52.7 PLS / Square foot	
"	3	Switchgrass	1.5 rate	79.0 PLS / Square foot	
23	1	Caucasian bluestem	.5 rate	38.1 PLS / Square foot	
"	2	Caucasian bluestem 3.1# PLS	1.0 rate	76.3 PLS / Square foot	
"	3	Caucasian bluestem	1.5 rate	114.4 PLS / Square foot	
24	1	Big Bluestem 160,000 seeds/lb	.5 rate	18.4 PLS / Square foot	
"	2	Big Bluestem 10.0# PLS/Ac	1.0 rate	36.7 PLS / Square foot	
"	3	Big Bluestem	1.5 rate	55.1 PLS / Square foot	

\1 CelPril coated

\2 Seed Biotics coated

\3 See discussion 1998

\4 Rates as per NRCS MOFOTG March 1997

Study 29I143G - Seed Coat/Seeding Rates Study										Spring Evaluations			Table #5
Plot													
Sub-plot #	Common Name Source	Days to Emerge *				Emergence Density Plants/Row Foot				Percent Stand **			
Legume Plots #1 - #13		1998	1999	2000	Ave	1998	1999	2000	Ave	1998	1999	2000	Ave
1/1		6.25	9.25	15.50	<b>10.33</b>	4.92	22.08	22.83	<b>16.61</b>	65.00	73.75	26.25	<b>55.00</b>
1/2	Alfalfa	6.25	8.00	14.50	<b>9.58</b>	9.84	19.67	30.17	<b>19.89</b>	83.75	78.75	27.50	<b>63.33</b>
1/3		6.25	9.25	15.00	<b>10.17</b>	17.67	25.58	26.16	<b>23.14</b>	87.50	84.50	37.50	<b>69.83</b>
2/1		6.50	9.25	19.50	<b>11.75</b>	6.92	18.34	10.41	<b>11.89</b>	47.50	88.75	26.25	<b>54.17</b>
2/2	Alfalfa	6.50	8.50	16.75	<b>10.58</b>	8.75	22.34	14.66	<b>15.25</b>	72.50	92.50	18.75	<b>61.25</b>
2/3	Celpril	6.50	9.75	16.75	<b>11.00</b>	15.25	22.50	12.16	<b>16.64</b>	81.25	93.25	23.75	<b>66.08</b>
3/1		6.00	8.50	19.75	<b>11.42</b>	4.42	17.83	6.00	<b>9.42</b>	62.50	70.00	8.75	<b>47.08</b>
3/2	Alfalfa	6.00	8.50	17.50	<b>10.67</b>	11.08	32.00	10.00	<b>17.69</b>	56.25	87.75	16.25	<b>53.42</b>
3/3	Seed Biotics	6.00	9.75	16.75	<b>10.83</b>	14.58	23.58	15.75	<b>17.97</b>	70.00	88.25	23.75	<b>60.67</b>
4/1		7.00	6.00	19.75	<b>10.92</b>	4.33	16.83	13.66	<b>11.61</b>	55.00	66.25	23.75	<b>48.33</b>
4/2	Red Clover	7.00	6.00	16.75	<b>9.92</b>	7.09	22.00	18.75	<b>15.95</b>	77.50	75.00	40.00	<b>64.17</b>
4/3		7.00	7.25	16.75	<b>10.33</b>	13.08	15.25	28.16	<b>18.83</b>	73.75	75.00	42.50	<b>63.75</b>
5/1		7.25	7.25	20.25	<b>11.58</b>	5.08	21.75	4.00	<b>10.28</b>	47.50	78.75	24.00	<b>50.08</b>
5/2	Red Clover	7.25	7.25	16.50	<b>10.33</b>	9.84	19.25	12.91	<b>14.00</b>	53.75	81.25	23.75	<b>52.92</b>
5/3	Celpril	7.25	8.50	16.75	<b>10.83</b>	11.50	19.08	18.25	<b>16.28</b>	76.25	83.75	26.25	<b>62.08</b>
6/1		7.25	8.50	22.50	<b>12.75</b>	6.08	19.92	8.17	<b>11.39</b>	50.00	63.75	16.50	<b>43.42</b>
6/2	Red Clover	7.25	7.25	18.25	<b>10.92</b>	9.83	12.17	12.66	<b>11.55</b>	58.75	72.00	30.25	<b>53.67</b>
6/3	Seed Biotics	7.25	9.75	17.00	<b>11.33</b>	11.42	14.92	15.25	<b>13.86</b>	70.00	71.25	12.50	<b>51.25</b>
7/1		8.50	13.25	21.00	<b>14.25</b>	5.50	14.50	10.33	<b>10.11</b>	47.50	70.00	15.75	<b>44.42</b>
7/2	Birdsfoot trefoil	8.50	11.25	18.75	<b>12.83</b>	7.67	21.68	18.42	<b>15.92</b>	58.75	68.75	26.50	<b>51.33</b>
7/3		8.50	10.50	25.18	<b>14.73</b>	10.67	19.50	29.57	<b>19.91</b>	77.50	72.50	35.00	<b>61.67</b>
8/1		7.75	12.50	22.50	<b>14.25</b>	4.17	11.58	7.08	<b>7.61</b>	43.75	68.75	31.25	<b>47.92</b>
8/2	Birdsfoot trefoil	7.75	12.50	19.50	<b>13.25</b>	10.25	8.75	15.08	<b>11.36</b>	63.75	68.75	26.25	<b>52.92</b>
8/3	Celpril	7.75	12.25	20.00	<b>13.33</b>	12.84	11.09	28.33	<b>17.42</b>	55.00	68.75	23.00	<b>48.92</b>
9/1		8.75	11.75	25.25	<b>15.25</b>	5.00	7.17	8.41	<b>6.86</b>	46.25	58.75	10.75	<b>38.58</b>
9/2	Birdsfoot trefoil	8.75	11.75	21.75	<b>14.08</b>	6.67	20.08	17.58	<b>14.78</b>	50.00	61.25	14.25	<b>41.83</b>
9/3	Seed Biotics	8.75	12.50	18.75	<b>13.33</b>	9.58	18.58	19.75	<b>15.97</b>	58.75	65.00	17.25	<b>47.00</b>
*	Number of days it took, from date planted, for 25 seedlings to emerge in that plot.												
**	Visual rating of percent of plot that has complete rows of plants.												

Study 29I143G - cont.													Spring Evaluations		Table #5		
Plot																	cont.
Sub-plot #	Common Name Source	Days to emerge *				Emergence density Plants/Row Foot				Percent Stand **							
		1998	1999	2000	Ave	1998	1999	2000	Ave	1998	1999	2000	Ave				
10/1		9.00	11.75	27.25	<b>16.00</b>	3.00	25.67	4.41	<b>11.02</b>	30.00	58.75	2.25	<b>30.33</b>				
10/2	Ladino clover	9.00	9.75	21.75	<b>13.50</b>	5.09	27.09	14.57	<b>15.58</b>	38.75	70.00	10.25	<b>39.67</b>				
10/3		9.00	9.75	21.00	<b>13.25</b>	9.50	28.58	20.33	<b>19.47</b>	30.00	72.50	15.25	<b>39.25</b>				
11/1		8.50	12.00	26.75	<b>15.75</b>	2.42	11.08	5.25	<b>6.25</b>	17.50	56.25	3.75	<b>25.83</b>				
11/2	Ladino clover	8.50	9.75	21.50	<b>13.25</b>	3.83	11.42	6.91	<b>7.39</b>	25.00	58.75	3.00	<b>28.92</b>				
11/3	Celpril	8.50	9.75	22.75	<b>13.67</b>	8.00	20.00	10.33	<b>12.78</b>	38.75	66.25	5.50	<b>36.83</b>				
12/1		9.00	11.75	26.25	<b>15.67</b>	4.08	14.33	5.08	<b>7.83</b>	20.00	57.50	12.50	<b>30.00</b>				
12/2	Ladino clover	9.00	11.75	23.00	<b>14.58</b>	9.08	19.67	8.83	<b>12.53</b>	41.25	70.00	21.25	<b>44.17</b>				
12/3	Seed Biotics	9.00	12.25	19.25	<b>13.50</b>	10.84	24.67	10.58	<b>15.36</b>	40.00	70.00	26.25	<b>45.42</b>				
13/1		8.75	18.00	26.75	<b>17.83</b>	5.67	14.67	2.08	<b>7.47</b>	33.75	50.00	14.00	<b>32.58</b>				
13/2	Annual Lespedeza	9.00	14.00	21.50	<b>14.83</b>	14.67	15.33	7.41	<b>12.47</b>	55.00	48.75	11.50	<b>38.42</b>				
13/3		9.00	16.00	21.25	<b>15.42</b>	16.08	18.33	12.25	<b>15.55</b>	56.25	63.75	20.25	<b>46.75</b>				
<b>Cool Season Grasses Plots #14 - #19</b>																	
		<b>1998</b>	<b>1999</b>	<b>2000</b>	<b>Ave</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>	<b>Ave</b>	<b>1998</b>	<b>1999</b>	<b>2000</b>	<b>Ave</b>				
14/1		5.00	18.00	19.00	<b>14.00</b>	14.50	15.67	18.33	<b>16.17</b>	78.75	73.75	16.25	<b>56.25</b>				
14/2	Tall fescue	5.00	18.00	18.50	<b>13.83</b>	28.67	32.08	33.00	<b>31.25</b>	88.75	86.67	20.00	<b>65.14</b>				
14/3	Endophyte infected	5.00	18.00	18.00	<b>13.67</b>	33.59	39.50	36.33	<b>36.47</b>	95.00	82.50	21.25	<b>66.25</b>				
15/1		19.00	0.00	33.25	<b>17.42</b>	1.42	0.00	1.41	<b>0.94</b>	7.50	0.00	2.00	<b>3.17</b>				
15/2	Tall fescue	19.00	0.00	31.50	<b>16.83</b>	0.83	0.00	12.00	<b>4.28</b>	7.50	0.00	2.75	<b>3.42</b>				
15/3	Endophyte free	19.00	0.00	26.75	<b>15.25</b>	3.33	0.00	6.67	<b>3.33</b>	11.25	0.00	1.00	<b>4.08</b>				
16/1		8.00	19.50	23.50	<b>17.00</b>	8.92	22.84	7.83	<b>13.20</b>	77.50	61.25	9.00	<b>49.25</b>				
16/2	Orchardgrass	8.00	18.00	21.25	<b>15.75</b>	19.50	21.67	12.16	<b>17.78</b>	80.00	61.25	7.50	<b>49.58</b>				
16/3		8.00	18.00	20.25	<b>15.42</b>	41.25	16.00	21.91	<b>26.39</b>	87.50	71.25	12.50	<b>57.08</b>				
17/1		8.00	18.75	25.00	<b>17.25</b>	9.33	21.67	5.50	<b>12.17</b>	63.75	38.75	6.50	<b>36.33</b>				
17/2	Smooth brome	8.00	18.00	22.00	<b>16.00</b>	11.09	22.00	8.66	<b>13.92</b>	76.25	52.50	3.00	<b>43.92</b>				
17/3		8.00	18.00	22.00	<b>16.00</b>	20.09	12.25	11.67	<b>14.67</b>	78.75	65.00	16.50	<b>53.42</b>				
*	Number of days it took, from date planted, for 25 seedlings to emerge in that plot.																
**	Visual rating of percent of plot that has complete rows of plants.																

Study 29I143G - cont.													Spring Evaluations		Table #5		
Plot																	cont.
Sub-plot #	Common Name Source	Days to emerge *			Emergence density Plants/Row Foot				Percent Stand **								
		1998	1999	2000	Ave	1998	1999	2000	Ave	1998	1999	2000	Ave				
18/1		8.00	36.00	28.00	<b>24.00</b>	16.67	17.00	3.00	<b>12.22</b>	67.50	23.75	4.75	<b>32.00</b>				
18/2	Timothy	8.00	23.25	24.50	<b>18.58</b>	28.33	8.09	7.33	<b>14.58</b>	73.75	31.25	3.00	<b>36.00</b>				
18/3		8.00	23.25	25.00	<b>18.75</b>	47.58	14.09	7.66	<b>23.11</b>	86.25	31.25	5.50	<b>41.00</b>				
19/1		8.00	22.00	22.50	<b>17.50</b>	7.59	2.42	13.75	<b>7.92</b>	68.75	18.75	2.25	<b>29.92</b>				
19/2	Canada wildrye	8.00	22.00	20.25	<b>16.75</b>	16.50	7.00	17.00	<b>13.50</b>	82.50	27.50	7.50	<b>39.17</b>				
19/3		8.00	22.00	20.25	<b>16.75</b>	20.50	3.59	23.50	<b>15.86</b>	90.00	33.75	8.75	<b>44.17</b>				
<b>Warm Season Grasses Plots #20 - #23</b>						not in											
						average											
20/1		N/A	33.75	0.00		1.17	5.75	1.00	<b>3.38</b>	17.50	23.75	0.00	<b>11.88</b>				
20/2	Eastern gamagrass	N/A	33.75	0.00		0.92	6.08	0.25	<b>3.17</b>	20.00	21.25	0.00	<b>10.63</b>				
20/3	untreated/Germtech	N/A	33.75	0.00		1.67	4.83	1.75	<b>3.29</b>	20.00	12.50	0.25	<b>6.38</b>				
21/1		16.50	46.00	32.75	<b>31.75</b>	1.42	6.67	0.83	<b>2.97</b>	27.50	15.00	3.25	<b>15.25</b>				
21/2	Eastern gamagrass	16.50	46.00	25.00	<b>29.17</b>	2.17	3.17	1.58	<b>2.30</b>	28.75	15.00	4.25	<b>16.00</b>				
21/3	treated	16.50	46.00	26.50	<b>29.67</b>	3.58	6.17	1.42	<b>3.72</b>	38.75	15.00	5.25	<b>19.67</b>				
22/1		20.75	46.00	18.25	<b>28.33</b>	12.17	1.00	6.08	<b>6.42</b>	27.50	10.00	35.00	<b>24.17</b>				
22/2	Switchgrass	20.75	46.00	15.75	<b>27.50</b>	10.50	2.84	14.33	<b>9.22</b>	25.00	10.00	40.00	<b>25.00</b>				
22/3		20.75	46.00	15.00	<b>27.25</b>	13.92	1.00	22.50	<b>12.47</b>	36.25	10.00	38.75	<b>28.33</b>				
23/1		23.00	49.00	32.50	<b>34.83</b>	4.50	5.00	7.17	<b>5.56</b>	11.25	35.00	26.25	<b>24.17</b>				
23/2	Caucasian bluestem	23.00	49.00	27.75	<b>33.25</b>	2.42	4.33	8.25	<b>5.00</b>	10.00	25.00	23.75	<b>19.58</b>				
23/3		23.00	49.00	25.75	<b>32.58</b>	2.17	10.00	20.50	<b>10.89</b>	5.00	30.00	23.75	<b>19.58</b>				
24/1			43.00	19.75	<b>31.38</b>		0.00	9.48	<b>4.74</b>		10.00	36.25	<b>23.13</b>				
24/2			43.00	15.75	<b>29.38</b>		6.00	18.92	<b>12.46</b>		35.00	41.25	<b>38.13</b>				
24/3			43.00	15.75	<b>29.38</b>		0.00	16.50	<b>8.25</b>		10.00	47.50	<b>28.75</b>				
*	Number of days it took, from date planted, for 25 seedlings to emerge in that plot.																
**	Visual rating of percent of plot that has complete rows of plants.																

Study 29I143G - Seed Coat/Seeding Rates Study													Table # 6A
October Evaluations													
Plot /		Cover Density (stems/row foot)				Cover Density (stems/row foot)				Cover Density (stems/row foot)			
Sub-	Common name	1st Year ( Year of Planting)				2nd Year ( Year after Planting)				3rd Year (2 Years after Planting)			
plot #	Source	1998	1999	2000	Average	1999	2000	2001	Average	2000	2001	2002	Average
		98 pltg	99pltg	00pltg	3 Yr	98 pltg	99pltg	00pltg	3 YR	98 pltg	99pltg	00pltg	3 YR
<b>Legume Plots #1 - #13</b>													
1/1		9.75	22.17	35.09	<b>22.33</b>	22.50	28.25	20.50	<b>23.75</b>	17.00	8.50	15.75	<b>13.75</b>
1/2	Alfalfa	9.4925	37.17	40.25	<b>28.97</b>	32.83	30.33	21.75	<b>28.31</b>	20.75	9.50	11.50	<b>13.92</b>
1/3		15.75	37.09	49.42	<b>34.08</b>	24.33	27.75	22.25	<b>24.78</b>	16.42	10.75	17.25	<b>14.81</b>
2/1		7.0025	21.75	19.67	<b>16.14</b>	23.92	31.17	17.25	<b>24.11</b>	19.50	19.75	15.75	<b>18.33</b>
2/2	Alfalfa	7.5	33.17	18.75	<b>19.81</b>	19.67	35.09	14.75	<b>23.17</b>	14.75	12.75	14.00	<b>13.83</b>
2/3	Celpril	9.3325	30.59	25.59	<b>21.83</b>	30.58	28.67	21.75	<b>27.00</b>	21.33	18.75	16.75	<b>18.94</b>
3/1		6.3325	18.08	21.67	<b>15.36</b>	28.83	21.92	18.00	<b>22.92</b>	21.17	13.00	14.25	<b>16.14</b>
3/2	Alfalfa	6.4975	17.25	32.34	<b>18.69</b>	34.42	40.75	17.75	<b>30.97</b>	19.00	12.50	15.75	<b>15.75</b>
3/3	Seed Biotics	10.333	22.84	25.42	<b>19.53</b>	20.42	29.92	18.25	<b>22.86</b>	25.00	8.00	13.25	<b>15.42</b>
4/1		43.25	35.00	39.67	<b>39.31</b>	29.42	39.58	20.00	<b>29.67</b>	25.58	25.00	10.50	<b>20.36</b>
4/2	Red Clover	54.803	36.50	71.42	<b>54.24</b>	85.00	31.25	29.75	<b>48.67</b>	24.67	22.00	16.00	<b>20.89</b>
4/3		54.583	39.67	61.00	<b>51.75</b>	90.00	42.67	31.75	<b>54.81</b>	21.83	22.25	14.75	<b>19.61</b>
5/1		31.333	32.92	22.33	<b>28.86</b>	40.75	44.34	14.50	<b>33.20</b>	20.92	38.75	21.50	<b>27.06</b>
5/2	Red Clover	42	38.50	30.17	<b>36.89</b>	51.75	43.34	11.50	<b>35.53</b>	20.92	24.25	17.00	<b>20.72</b>
5/3	Celpril	51.583	38.33	69.92	<b>53.28</b>	39.50	35.50	32.75	<b>35.92</b>	15.50	15.50	14.25	<b>15.08</b>
6/1		14.665	29.75	15.83	<b>20.08</b>	44.67	37.00	19.00	<b>33.56</b>	28.50	29.50	19.00	<b>25.67</b>
6/2	Red Clover	30.418	40.00	20.75	<b>30.39</b>	47.00	31.83	23.50	<b>34.11</b>	44.83	37.00	11.50	<b>31.11</b>
6/3	Seed Biotics	36.668	48.50	45.75	<b>43.64</b>	52.25	32.00	29.00	<b>37.75</b>	28.34	17.75	18.75	<b>21.61</b>
7/1		11.418	22.00	21.92	<b>18.45</b>	38.25	38.84	30.50	<b>35.86</b>	23.17	10.75	78.00	<b>37.31</b>
7/2	Birdsfoot trefoil	13.665	46.08	34.25	<b>31.33</b>	50.00	50.08	34.25	<b>44.78</b>	27.09	16.75	84.25	<b>42.70</b>
7/3		19.165	45.75	35.42	<b>33.44</b>	40.00	55.75	37.50	<b>44.42</b>	20.34	16.75	84.00	<b>40.36</b>
8/1		7.25	21.25	18.83	<b>15.78</b>	36.17	45.25	25.25	<b>35.56</b>	12.25	3.58	68.50	<b>28.11</b>
8/2	Birdsfoot trefoil	14.25	20.83	34.92	<b>23.33</b>	40.25	49.50	25.25	<b>38.33</b>	14.34	10.75	64.50	<b>29.86</b>
8/3	Celpril	10.918	26.50	38.58	<b>25.33</b>	29.33	45.34	25.75	<b>33.47</b>	17.42	15.00	80.25	<b>37.56</b>
9/1		9.4175	19.67	15.17	<b>14.75</b>	39.00	36.58	30.75	<b>35.44</b>	11.75	9.75	80.25	<b>33.92</b>
9/2	Birdsfoot trefoil	12.668	30.42	26.25	<b>23.11</b>	45.00	50.83	36.25	<b>44.03</b>	18.25	9.75	81.50	<b>36.50</b>
9/3	Seed Biotics	12.75	34.42	22.42	<b>23.20</b>	36.92	52.33	37.25	<b>42.17</b>	19.83	13.75	83.75	<b>39.11</b>

Study 29I143G - continued													Table #6A
													cont.
Plot /		Cover Density (stems/row foot)				Cover Density (stems/row foot)				Cover Density (stems/row foot)			
Sub-	Common name	1st Year ( Year of Planting)				2nd Year ( Year after Planting)				3rd Year (2 Years after Planting)			
plot #	Source	1998	1999	2000	Average	1999	2000	2001	Average	2000	2001	2002	Average
		98 pltg	99pltg	00pltg	3 YR	98 pltg	99pltg	00pltg	3 YR	98 pltg	99pltg	00pltg	3 YR
10/1		15.918	31.33	51.75	<b>33.00</b>	47.09	52.75	46.25	<b>48.70</b>	32.84	35.25	11.00	<b>26.36</b>
10/2	Ladino clover	9.74	26.17	81.83	<b>39.25</b>	54.00	62.17	54.25	<b>56.81</b>	40.17	29.00	19.50	<b>29.56</b>
10/3		27.75	33.83	93.42	<b>51.67</b>	43.08	62.75	48.75	<b>51.53</b>	48.67	26.00	9.75	<b>28.14</b>
11/1		11.5	22.59	31.92	<b>22.00</b>	56.75	72.50	47.25	<b>58.83</b>	36.17	29.25	6.75	<b>24.06</b>
11/2	Ladino clover	3.6675	34.51	29.00	<b>22.39</b>	46.00	61.67	39.00	<b>48.89</b>	43.17	33.00	4.75	<b>26.97</b>
11/3	Celpril	28	36.08	34.67	<b>32.92</b>	72.25	68.58	46.00	<b>62.28</b>	39.08	16.25	5.25	<b>20.19</b>
12/1		9.4175	24.83	31.92	<b>22.06</b>	55.00	58.83	44.75	<b>52.86</b>	38.92	29.00	16.75	<b>28.22</b>
12/2	Ladino clover	20.085	36.75	51.42	<b>36.08</b>	40.50	53.92	47.25	<b>47.22</b>	46.92	27.75	28.75	<b>34.47</b>
12/3	Seed Biotics	18.5	36.75	71.92	<b>42.39</b>	64.17	55.92	52.25	<b>57.45</b>	30.67	33.50	35.25	<b>33.14</b>
13/1		0.915	7.83	4.25	<b>4.33</b>	10.08	5.75	4.25	<b>6.69</b>	3.50	5.00	8.00	<b>5.50</b>
13/2	Annual Lespedeza	3	4.08	4.92	<b>4.00</b>	3.00	2.92	5.75	<b>3.89</b>	6.42	3.25	7.00	<b>5.56</b>
13/3		2.75	9.83	15.17	<b>9.25</b>	19.42	4.08	6.00	<b>9.83</b>	3.50	1.75	4.25	<b>3.17</b>
<b>Cool Season Grasses Plots #14 - #19</b>													
14/1		33	35.58	18.42	<b>29.00</b>	55.09	58.58		<b>56.83</b>	60.17	34.25	24.00	<b>39.47</b>
14/2	Tall fescue	30.25	35.67	23.75	<b>29.89</b>	42.08	68.75		<b>55.42</b>	54.50	30.00	27.75	<b>37.42</b>
14/3	Endophyte infected	34.333	41.50	27.67	<b>34.50</b>	46.34	76.33		<b>61.33</b>	57.58	29.50	24.75	<b>37.28</b>
15/1		4.585	0.67	3.00	<b>2.75</b>	44.42	0.17		<b>22.29</b>	14.83	4.08	9.75	<b>9.55</b>
15/2	Tall fescue	4.1675	4.33	2.17	<b>3.56</b>	44.25	0.92		<b>22.58</b>	16.08	1.25	13.75	<b>10.36</b>
15/3	Endophyte free	9	1.33	0.92	<b>3.75</b>	37.00	4.17		<b>20.58</b>	16.42	1.33	10.50	<b>9.42</b>
16/1		19.665	23.17	13.75	<b>18.86</b>	41.67	54.25		<b>47.96</b>	47.59	32.75	27.50	<b>35.95</b>
16/2	Orchardgrass	23.25	28.17	21.59	<b>24.33</b>	44.50	53.17		<b>48.83</b>	40.17	37.50	25.25	<b>34.31</b>
16/3		25.583	28.42	29.00	<b>27.67</b>	43.17	41.83		<b>42.50</b>	45.33	45.00	24.00	<b>38.11</b>
17/1		12.168	21.34	11.25	<b>14.92</b>	44.33	39.92		<b>42.12</b>	39.25	33.00	16.75	<b>29.67</b>
17/2	Smooth brome	18	15.00	16.84	<b>16.61</b>	64.17	37.00		<b>50.59</b>	38.75	37.50	17.75	<b>31.33</b>
17/3		23.5	26.00	20.42	<b>23.31</b>	45.42	40.00		<b>42.71</b>	48.75	39.75	16.00	<b>34.83</b>

Study 29I143G - continued													Table #6A
													cont.
Plot /		Cover Density (stems/row foot)				Cover Density (stems/row foot)				Cover Density (stems/row foot)			
Sub-	Common name	1st Year ( Year of Planting)				2nd Year ( Year after Planting)				3rd Year (2 Years after Planting)			
plot #	Source	1998	1999	2000	Average	1999	2000	2001	Average	2000	2001	2002	Average
		98 pltg	99pltg	00pltg	3 YR	98 pltg	99pltg	00pltg	3 YR	98 pltg	99pltg	00pltg	3 YR
18/1		12.083	6.42	13.00	<b>10.50</b>	41.17	11.25	33.50	<b>28.64</b>	33.00	19.50	18.25	<b>23.58</b>
18/2	Timothy	22.748	1.83	3.25	<b>9.28</b>	61.00	14.00	45.50	<b>40.17</b>	27.75	16.75	25.00	<b>23.17</b>
18/3		22.253	5.83	3.25	<b>10.45</b>	58.25	33.75	29.00	<b>40.33</b>	23.17	10.25	23.00	<b>18.81</b>
19/1		8.0825	3.25	18.50	<b>9.94</b>	30.59	37.58	21.75	<b>29.97</b>	33.75	4.75	26.50	<b>21.67</b>
19/2	Canada wildrye	12.918	3.42	19.67	<b>12.00</b>	32.92	64.17	17.75	<b>38.28</b>	28.08	3.75	24.25	<b>18.69</b>
19/3		13.25	7.09	15.25	<b>11.86</b>	36.92	57.92	28.75	<b>41.19</b>	38.17	3.75	30.25	<b>24.06</b>
<b>Warm Season Grasses Plots #20 - #23</b>													
20/1		4.165	14.33	6.67	<b>10.50</b>	53.33	33.59	1.75	<b>29.56</b>	26.00	9.00	12.25	<b>15.75</b>
20/2	Eastern gamagrass	7.665	7.44	9.50	<b>8.47</b>	55.33	32.08	7.75	<b>31.72</b>	55.00	17.50	10.75	<b>27.75</b>
20/3	untreated/Germtech	7	4.67	19.84	<b>12.26</b>	24.72	39.00	23.75	<b>29.16</b>	25.42	16.25	17.00	<b>19.56</b>
21/1		12	8.11	12.58	<b>10.90</b>	31.42	31.42	8.00	<b>23.61</b>	32.25	24.25	12.75	<b>23.08</b>
21/2	Eastern gamagrass	18.335	12.78	18.08	<b>16.40</b>	53.08	23.67	17.50	<b>31.42</b>	39.50	24.25	16.00	<b>26.58</b>
21/3	treated - wet	14.833	3.78	19.67	<b>12.76</b>	49.08	7.25	11.25	<b>22.53</b>	44.17	14.75	12.75	<b>23.89</b>
22/1		8.915	14.84	11.00	<b>11.58</b>	16.25	38.59	25.25	<b>26.70</b>	58.67	15.75	33.25	<b>35.89</b>
22/2	Switchgrass	8.165	14.00	19.67	<b>13.94</b>	25.33	25.59	32.25	<b>27.72</b>	40.67	25.75	35.50	<b>33.97</b>
22/3		9.4175	14.50	18.50	<b>14.14</b>	18.17	20.17	26.00	<b>21.44</b>	30.00	21.50	39.75	<b>30.42</b>
23/1		23.665	13.50	88.50	<b>41.89</b>	40.17	66.42	49.25	<b>51.94</b>	75.92	57.00	139.25	<b>90.72</b>
23/2	Caucasian bluestem	26.748	7.33	85.08	<b>39.72</b>	70.92	53.42	43.00	<b>55.78</b>	100.33	43.75	171.75	<b>105.28</b>
23/3		30.918	12.00	92.83	<b>45.25</b>	65.00	42.09	54.00	<b>53.70</b>	91.33	36.00	109.25	<b>78.86</b>
24/1	Big bluestem		5.92	44.17	<b>16.69</b>		44.92	89.75	<b>67.33</b>		23.75	52.75	<b>38.25</b>
24/2			10.42	44.75	<b>18.39</b>		33.83	97.00	<b>65.42</b>		56.50	56.25	<b>56.38</b>
24/3			15.25	58.84	<b>24.70</b>		58.58	75.75	<b>67.17</b>		54.50	62.25	<b>58.38</b>

Study 29I143G - Seed Coat/Seeding Rates Study													Table #6B
October Evaluations													
Plot /	Percent Cover (Visual Observation)					Percent Cover (Visual Observation)				Percent Cover (Visual Observation)			
Sub-	Common name	1st Year (Year Planted)				2nd Year (Year after Planted)				3rd Year (2nd Year after Planted)			
plot #	Source	1998	1999	2000	Average	1999	2000	2001	Average	2000	2001	2002	Average
		98 pltg	99 pltg	00 pltg	3 Yr	98 pltg	99 pltg	00 pltg	3 yr	98 pltg	99 pltg	00 pltg	3 YR
<b>Legume Plots #1 - #13</b>													
1/1		70.00	66.25	39.25	<b>58.50</b>	66.25	78.75	85.75	<b>76.92</b>	65.00	71.25	61.25	<b>65.83</b>
1/2	Alfalfa	78.75	72.50	42.50	<b>64.58</b>	72.50	78.75	80.00	<b>77.08</b>	70.00	75.25	45.00	<b>63.42</b>
1/3		70.00	62.50	50.00	<b>60.83</b>	62.50	81.50	89.25	<b>77.75</b>	61.25	63.25	63.75	<b>62.75</b>
2/1		52.50	63.75	25.00	<b>47.08</b>	63.75	66.25	86.25	<b>72.08</b>	41.25	76.25	72.50	<b>63.33</b>
2/2	Alfalfa	57.50	46.25	13.75	<b>39.17</b>	46.25	73.75	84.50	<b>68.17</b>	40.00	85.00	62.50	<b>62.50</b>
2/3	Celpril	62.50	58.75	20.00	<b>47.08</b>	58.75	77.50	89.00	<b>75.08</b>	45.00	68.75	81.25	<b>65.00</b>
3/1		51.25	76.25	22.75	<b>50.08</b>	76.25	65.00	70.25	<b>70.50</b>	51.25	68.75	48.75	<b>56.25</b>
3/2	Alfalfa	71.25	77.50	40.25	<b>63.00</b>	77.50	68.75	75.25	<b>73.83</b>	62.50	72.50	57.50	<b>64.17</b>
3/3	Seed Biotics	68.75	71.25	37.75	<b>59.25</b>	71.25	66.25	69.50	<b>69.00</b>	46.25	63.75	55.00	<b>55.00</b>
4/1		81.25	77.50	45.00	<b>67.92</b>	77.50	62.50	15.00	<b>51.67</b>	32.50	36.25	20.00	<b>29.58</b>
4/2	Red Clover	93.25	77.50	41.25	<b>70.67</b>	77.50	62.50	23.75	<b>54.58</b>	41.25	37.50	32.50	<b>37.08</b>
4/3		90.25	58.75	42.00	<b>63.67</b>	58.75	63.75	23.00	<b>48.50</b>	25.00	31.25	30.00	<b>28.75</b>
5/1		72.50	51.25	23.75	<b>49.17</b>	51.25	79.50	15.00	<b>48.58</b>	18.00	25.50	30.00	<b>24.50</b>
5/2	Red Clover	82.50	46.25	48.75	<b>59.17</b>	46.25	76.75	9.00	<b>44.00</b>	28.00	13.00	28.75	<b>23.25</b>
5/3	Celpril	91.50	56.25	55.00	<b>67.58</b>	56.25	77.50	21.75	<b>51.83</b>	17.50	25.00	31.25	<b>24.58</b>
6/1		61.25	78.75	25.00	<b>55.00</b>	78.75	51.25	16.25	<b>48.75</b>	38.75	43.25	47.50	<b>43.17</b>
6/2	Red Clover	73.75	73.75	35.00	<b>60.83</b>	73.75	61.25	17.50	<b>50.83</b>	41.25	23.25	29.25	<b>31.25</b>
6/3	Seed Biotics	81.25	61.25	42.50	<b>61.67</b>	61.25	61.25	35.00	<b>52.50</b>	45.00	31.25	38.75	<b>38.33</b>
7/1		37.50	60.00	37.75	<b>45.08</b>	60.00	67.50	53.75	<b>60.42</b>	25.75	22.50	90.00	<b>46.08</b>
7/2	Birdsfoot trefoil	53.75	51.83	40.75	<b>48.78</b>	51.83	62.50	68.75	<b>61.03</b>	32.50	28.75	93.75	<b>51.67</b>
7/3		55.00	67.50	45.00	<b>55.83</b>	67.50	68.25	67.50	<b>67.75</b>	27.50	53.75	91.25	<b>57.50</b>
8/1		36.25	57.50	33.00	<b>42.25</b>	57.50	66.25	46.25	<b>56.67</b>	25.50	10.50	72.50	<b>36.17</b>
8/2	Birdsfoot trefoil	56.25	58.75	40.00	<b>51.67</b>	58.75	72.50	63.75	<b>65.00</b>	31.25	18.75	85.00	<b>45.00</b>
8/3	Celpril	45.00	76.25	45.00	<b>55.42</b>	76.25	72.50	40.00	<b>62.92</b>	22.50	23.75	87.50	<b>44.58</b>
9/1		42.50	37.50	24.25	<b>34.75</b>	37.50	58.75	48.75	<b>48.33</b>	27.50	21.21	91.25	<b>46.65</b>
9/2	Birdsfoot trefoil	46.25	36.25	32.50	<b>38.33</b>	36.25	65.00	68.75	<b>56.67</b>	38.75	27.50	92.00	<b>52.75</b>
9/3	Seed Biotics	53.75	60.00	22.75	<b>45.50</b>	60.00	73.75	52.50	<b>62.08</b>	25.00	45.50	93.25	<b>54.58</b>

Study 29I143G - cont.													Table #6B
													cont.
Plot /		Percent Cover (Visual Observation)				Percent Cover (Visual Observation)				Percent Cover (Visual Observation)			
Sub-	Common name	1st Year (Year Planted)				2nd Year (Year after Planted)				3rd Year (2nd Year after Planted)			
plot #	Source	1998	1999	2000	Average	1999	2000	2001	Average	2000	2001	2002	Average
		98 pltg	99 pltg	00 pltg	3 Yr	98 pltg	99 pltg	00 pltg	3 Yr	98 pltg	99 pltg	00 pltg	3 YR
10/1		36.25	37.50	41.25	<b>38.33</b>	73.75	90.00	65.00	<b>76.25</b>	67.50	61.25	21.25	<b>50.00</b>
10/2	Ladino clover	38.75	31.25	48.75	<b>39.58</b>	75.00	93.75	70.50	<b>79.75</b>	70.00	75.00	30.00	<b>58.33</b>
10/3		53.75	35.00	52.00	<b>46.92</b>	76.25	87.25	55.00	<b>72.83</b>	81.25	60.00	21.25	<b>54.17</b>
11/1		22.50	35.00	17.25	<b>24.92</b>	68.75	94.25	80.00	<b>81.00</b>	50.00	26.75	13.75	<b>30.17</b>
11/2	Ladino clover	30.00	26.25	25.00	<b>27.08</b>	71.25	94.00	75.00	<b>80.08</b>	46.25	31.25	9.25	<b>28.92</b>
11/3	Celpril	35.50	35.00	17.50	<b>29.33</b>	72.50	92.75	77.50	<b>80.92</b>	57.50	34.75	13.00	<b>35.08</b>
12/1		32.50	26.25	28.75	<b>29.17</b>	65.00	92.00	77.50	<b>78.17</b>	60.00	52.50	37.50	<b>50.00</b>
12/2	Ladino clover	55.00	33.75	32.50	<b>40.42</b>	61.25	93.00	81.25	<b>78.50</b>	63.75	53.00	51.25	<b>56.00</b>
12/3	Seed Biotics	53.75	28.75	28.75	<b>37.08</b>	65.00	93.75	83.00	<b>80.58</b>	51.25	56.25	86.25	<b>64.58</b>
13/1		6.25	25.00	9.75	<b>13.67</b>	19.00	4.25	3.50	<b>8.92</b>	3.75	17.50	13.75	<b>11.67</b>
13/2	Annual Lespedeza	10.00	0.00	4.75	<b>4.92</b>	8.75	1.00	1.75	<b>3.83</b>	3.75	6.50	11.25	<b>7.17</b>
13/3		10.00	10.00	14.25	<b>11.42</b>	22.75	2.75	2.50	<b>9.33</b>	4.00	9.00	12.50	<b>8.50</b>
<b>Cool Season Grasses Plots #14 - #19</b>													
14/1		86.25	62.50	31.25	<b>60.00</b>	95.50	96.50	92.50	<b>94.83</b>	98.00	88.75	95.00	<b>93.92</b>
14/2	Tall fescue	90.00	74.75	50.00	<b>53.69</b>	94.25	94.25	94.50	<b>94.33</b>	98.75	90.25	95.00	<b>94.67</b>
14/3	Endophyte infected	95.75	78.00	48.75	<b>55.63</b>	98.50	93.25	93.00	<b>94.92</b>	98.00	95.00	96.25	<b>96.42</b>
15/1		6.50	55.00	1.00		45.00	0.50	26.25	<b>23.92</b>	21.25	8.75	38.75	<b>22.92</b>
15/2	Tall fescue	16.75	50.00	1.50		48.75	1.50	31.25	<b>27.17</b>	22.50	8.00	47.50	<b>26.00</b>
15/3	Endophyte free	21.25	45.00	1.50		60.00	1.75	17.50	<b>26.42</b>	23.75	13.75	50.00	<b>29.17</b>
16/1		70.00	52.50	23.25	<b>48.58</b>	75.00	35.00	73.75	<b>61.25</b>	51.25	87.25	92.50	<b>77.00</b>
16/2	Orchardgrass	78.75	67.50	30.00	<b>58.75</b>	81.00	36.75	78.75	<b>65.50</b>	41.25	81.75	93.75	<b>72.25</b>
16/3		87.50	73.75	32.50	<b>64.58</b>	92.75	45.00	82.50	<b>73.42</b>	60.00	87.00	92.50	<b>79.83</b>
17/1		76.25	27.50	8.50	<b>37.42</b>	91.25	77.50	89.25	<b>86.00</b>	95.25	95.75	96.25	<b>95.75</b>
17/2	Smooth brome	82.50	28.75	20.75	<b>44.00</b>	93.25	87.50	93.75	<b>91.50</b>	96.75	95.00	100.00	<b>97.25</b>
17/3		87.75	42.50	20.00	<b>50.08</b>	94.50	84.75	96.00	<b>91.75</b>	97.75	99.25	100.00	<b>99.00</b>

Study 29I143G - cont.													Table #6B
													cont.
Plot /	Percent Cover (Visual Observation)					Percent Cover (Visual Observation)				Percent Cover (Visual Observation)			
Sub-plot #	Common name	1st Year (Year Planted)			Average	2nd Year (Year after Planted)			Average	3rd Year (2nd Year after Planted)			Average
	Source	1998	1999	2000	3 Yr	1999	2000	2001	3 Yr	2000	2001	2002	3 Yr
		98 pltg	99 pltg	00 pltg		98 pltg	99 pltg	00 pltg		98 pltg	99 pltg	00 pltg	
18/1		43.75	30.25	2.50	<b>25.50</b>	47.50	5.50	37.50	<b>30.17</b>	27.50	19.25	81.25	<b>42.67</b>
18/2	Timothy	55.50	22.50	2.75	<b>26.92</b>	50.00	7.00	36.25	<b>31.08</b>	26.25	8.25	83.75	<b>39.42</b>
18/3		62.50	33.00	4.50	<b>33.33</b>	55.00	13.75	47.50	<b>38.75</b>	35.00	10.75	85.00	<b>43.58</b>
19/1		22.00	31.25	31.25	<b>28.17</b>	48.75	16.25	40.00	<b>35.00</b>	40.00	23.00	95.00	<b>52.67</b>
19/2	Canada wildrye	40.25	42.50	29.50	<b>37.42</b>	57.50	35.00	57.50	<b>50.00</b>	48.75	23.75	95.00	<b>55.83</b>
19/3		47.50	51.25	31.25	<b>43.33</b>	58.75	48.75	60.00	<b>55.83</b>	46.25	22.50	95.00	<b>54.58</b>
<b>Warm Season Grasses Plots #20 - #23</b>					plot #20 - 1998 untreated dormant seeding (UT), 1999 & 2000, germtech seeding (GT)								
		UT	GT	GT	2Yr GT Ave	UT	GT	GT	2Yr GT A	UT		GT	
20/1		15.00	26.67	3.75	<b>15.21</b>	63.75	38.50	13.75	<b>26.13</b>	26.25		47.50	47.50
20/2	Eastern gamagrass	21.25	41.67	5.00	<b>23.34</b>	55.00	39.25	32.50	<b>35.88</b>	52.50		72.50	72.50
20/3	untreated/Germtech	27.50	36.67	4.75	<b>20.71</b>	65.00	18.50	48.75	<b>33.63</b>	42.50		76.25	76.25
21/1		38.75	32.50	8.00	<b>26.42</b>	73.75	4.00	40.00	<b>39.25</b>	50.00		71.25	71.25
21/2	Eastern gamagrass	55.00	25.00	15.00	<b>31.67</b>	85.00	9.50	62.50	<b>52.33</b>	70.00		82.50	82.50
21/3	treated - wet	60.00	23.75	19.50	<b>34.42</b>	77.50	11.50	71.25	<b>53.42</b>	77.50		85.00	85.00
					<b>3 Yr Ave</b>				<b>3 Yr Ave</b>				<b>2 Yr Ave</b>
22/1		28.75	45.00	35.00	<b>36.25</b>	62.50	32.50	88.75	<b>61.25</b>	60.00		97.50	<b>78.75</b>
22/2	Switchgrass	40.00	45.00	55.00	<b>46.67</b>	68.75	52.50	96.75	<b>72.67</b>	68.75		100.00	<b>84.38</b>
22/3		51.25	48.75	47.50	<b>49.17</b>	70.00	35.00	92.00	<b>65.67</b>	72.50		98.75	<b>85.63</b>
23/1		56.25	21.25	57.50	<b>45.00</b>	56.25	82.50	70.75	<b>69.83</b>	80.00		96.25	<b>88.13</b>
23/2	Caucasian bluestem	63.75	28.75	75.00	<b>55.83</b>	78.75	75.00	91.50	<b>81.75</b>	80.00		97.50	<b>88.75</b>
23/3		73.75	36.25	67.50	<b>59.17</b>	75.00	73.75	83.50	<b>77.42</b>	72.50		97.50	<b>85.00</b>
					<b>2 Yr Ave</b>				<b>2 Yr Ave</b>				<b>2 Yr Ave</b>
24/1	Big bluestem		37.50	53.75	<b>45.63</b>		27.50	88.25	<b>57.88</b>		55.00	96.25	<b>75.63</b>
24/2			51.25	61.25	<b>56.25</b>		14.25	91.25	<b>52.75</b>		60.00	98.75	<b>79.38</b>
24/3			32.50	72.50	<b>52.50</b>		28.75	95.50	<b>62.13</b>		75.00	98.75	<b>86.88</b>

**Table #7**

**Study 29I143G - Seed Coating / Seeding Rates Study**

<b>Weather Data</b>	<b>Monthly Precipitation (Inches)</b>		
	<b>Year 1998</b>	<b>Year 1999</b>	<b>Year 2000</b>
<b>April</b>	3.32	4.18	0.84
<b>May</b>	2.21	4.05	7.19
<b>June</b>	5.57	2.00	6.23
<b>July</b>	5.74	2.03	2.91
<b>August</b>	0.31	0.45	6.01
<b>September</b>	4.07	1.15	3.36
<b>October</b>	2.7	1.88	3.3
<b>Total</b>	<b>20.6</b>	<b>11.56</b>	<b>29.84</b>

**Study ID Code:** MOPMC-P-0001-WO, WL, WE

**Study Title:** Assembly, Evaluation and Selection of Bur Oak, *Quercus macrocarpa* Michx.

**Study Leader:** Henry, J.

**Description:**

Bur oak is a large-size tree 60-80 feet tall and 2-3 feet in diameter (max. 170 by 7 feet); crown rounded with large, heavy branches. Leaves are deciduous, oblong to ovate; 6-12 inches long; characteristically 5-9 lobed, with rounded lobes. Fruit matures in one year; acorns are 3/5-2 inches long, ellipsoidal, brown, enclosed for 1/3 to all of its length in a characteristic fringe-margined cup. Twigs are stout; yellow-brown to gray, often with characteristic corky wings. Winter buds; 1/8-1/4 inch long, hairy. Bur oak is one of the largest American Oaks. Commonly distributed throughout Missouri, Iowa and Illinois, bur oak are is important bottomland tree, frequently found in moist flats, wetlands, and undulating flood plains. Important associates of bur oak include red maple, American elm, silver maple, swamp white oak, sycamore and eastern cottonwood.

**Objective:**

The objective of this study is to select a local source, fast growing, high nut producing bur oak.

## **Materials and Methods:**

Field collections were assembled, accessioned, and held in storage until the collection period was ended. The assemblage of collections began at the PMC in October 2000 and ended mid-December 2000. After the collection period was over the seed was stratified and planted in the greenhouse using the Root Pruning Method (RPM) containers. The plants will be transplanted in Field #7 on the PMC in mid to late April 2002. The design will be a randomized complete block with one plant per plot: one block for the Iowa collections, one for the Illinois collections and one block for the Missouri collections.

## **Discussion**

### **2000**

A total of 24 collections were made from the PMC three state service area: seven from Iowa, two from Illinois and 15 from Missouri. As these collections arrived at the PMC they were given accession numbers and placed in stratification for 120 days (cool moist storage 38 degrees Fahrenheit). At the time this report was being developed, these collections were being germinated in the greenhouse.

### **2001**

The 24 collections of bur oaks were taken out of the germination trays and placed in containers (3 5/8" x 6") and allowed to grow to approximately one foot tall. These plants were later transplanted into one-gallon size containers and placed in the portable greenhouse. In early December 2001 the plants were transported to the root cellar for over wintering. The scheduled planting date is April 2002. The plantings will be randomized complete block designs with one block for Iowa's collections, one block for Illinois' collections and one block for Missouri's collections.

Refer to Table #1 for collection information.

### **2002**

Three assemblies of bur oaks were planted in April 2002 representing each state's collections, Iowa, Illinois and Missouri. Iowa's collections were planted in Field #6 on April 18, 2002, Illinois' collections were planted in Field #12 on April 17, 2002, and Missouri's collections were planted in Field #7 on April 18 – 19, 2002. These collections were evaluated for height, spread, vigor, and insect and disease resistance. The evaluation data was not documented in this year's report but will be in the 2003 Annual Technical Report. Table #1 reflects collection information.

**Table # 1**

**Study Title:** Assembly, Evaluation and Selection of Bur Oak *Quercus macrocarpa* Michx.

<b>Temporary No.</b>	<b>State</b>	<b>County</b>	<b>MLRA</b>	<b>Collector</b>
MO-1	Missouri	Calloway	115	Thomas L. Wekenborg
MO-2	Missouri	Chariton	NA	Charles Lewis
MO-3	Missouri	Shannon	053	Randy Misser
MO-4	Missouri	Lincoln	115	Jimmy Henry
MO-5	Missouri	Lincoln	115	Jimmy Henry
MO-6	Missouri	Lincoln		Wayne Lovelace
MO-7	Missouri			
MO-8	Missouri	Pike		Keith Jackson
MO-9	Missouri	Pike		Keith Jackson
MO-10	Missouri	Pike		Keith Jackson
MO-11	Missouri	Pike		Keith Jackson
MO-12	Missouri	Howard	N/A	Robert D. Dewitt
MO-13	Missouri	Boone	N/A	Robert D. Dewitt
MO-14	Missouri	St. Charles	115	Dan Crigler
MO-15	Missouri	Moniteau	115	Douglas Wallace
IL-1	Illinois	Clark	N/a	David E. Hiatt
IL-1	Illinois	Jasper	113	Dennis D. Clency
IA-1	Iowa	Dickinson	103	Tim K. Moran
IA-2	Iowa	Dickinson	103	Tim K. Moran
IA-3	Iowa	Dickinson	103	Tim K. Moran
IA-4	Iowa	Wayne	N/A	Duane Bedford
IA-5	Iowa	Decatur	109	Kevin Reynolds
IA-6	Iowa	Bremer	104	Richard J. Cornes
IA-7	Iowa	Black	104	Rick Cordes

**Study ID Code:** MOPMC-P-0002-WE, WL

**Study Title:** Assembly, Evaluation and Selection of False Indigo Bush, *Amorpha fruticosa*, L.

**Study Leader:** Henry, J.

**Description:**

False indigo bush, *Amorpha fruticosa* L., is a medium sized shrub up to ten feet in height. The general shape is an open canopy with the bulk of foliage and twigs in the upper 1/3 of the crown. The leaves are alternate, pinnately compound. Each leaflet is up to two inches long and just over one inch wide with a small, bristly like point at the rounded tip. The flowers are in dense spikes on the upper part of the plant, often several spikes clustered together. Each flower has dark

indigo-purple petals with yellow tipped stamens. Flowering time: late spring to midsummer. Twigs are rigid, glabrous, red-brown or gray, often with an insect caused, long swelling near the tip. The fruit is a small, warty kidney shaped pod (1/2 inch long), with large glandular dots, in a crowded cylindrical cluster. The fruit persist on the shrub through winter. Found in more open areas along lakes and streams. May be found in upland areas where additional moisture is received.

### **Objective:**

The objective of this study is to select a local source, fast growing, high seed producing false indigo bush.

### **Materials and Methods:**

Field collections were assembled, accessioned and held in storage until the collection period ended. The assemblage of collections began at the PMC in November 2000. After the collection period was over, the seed was planted in the greenhouse using the Root Pruning Method (RPM) containers. The plants will be transplanted in a selected field on the PMC (preferably bottomland site). The design will be a randomized complete block with three plants per plot: one block for the Iowa collections, one for the Illinois collections and one for the Missouri collections.

### **Discussion:**

#### **2000**

A total of 32 collections were made from the PMC three state service area including North Dakota: 19 from Iowa, eight from Missouri, four from Illinois and one from North Dakota. On February 15, 2000, these collections were given accession numbers and placed in the PMC greenhouse for germination. At the time this report was written these collections were continuing germination in the greenhouse. During the period April-May 2001 these collections will be planted in Fields #6, #7 and #10 on the PMC. Collections from each state will be planted in separate fields on the PMC. The planting design will be a randomized complete block with three plants per plot. Refer to Table #1 for collection information.

#### **2001**

Three separate plantings were established in the month of June 2001: Iowa's collections of false indigo bush were planted in Field #10 on the PMC on June 21, Illinois collections were planted in Field # 6 on June 20 and Missouri's collections were planted in Field # 7 on June 21. Each planting reflected a randomized complete block design with four plants per plot. Survival evaluations were conducted in October 2001.

#### **2002**

The three separate plantings representing Iowa, Illinois and Missouri collections were evaluated on several occasions in 2002 to document vigor, height, spread, insect and disease resistance and seed production. Table #1 contains collection information. The evaluation data was not documented in this year's report but will be in the 2003 Annual Technical Report.

Table #1

<b>Temporary No.</b>	<b>State</b>	<b>County</b>	<b>MLRA</b>	<b>Collector</b>
MO-1	Missouri	Audrain	N/A	Mack Ellis
MO-2	Missouri	Knox	N/A	John Keith Doug Rainey
MO-3	Missouri	Marion	N/A	Mack Ellis Jay Lingwall
MO-4	Missouri		N/A	Maurice Davis
MO-5	Missouri		N/A	Maurice Davis
MO-6	Missouri	Lincoln	115	Jerry Kaiser
MO-7	Missouri	Pike	115	Keith Jackson
MO-8	Missouri	Pettis	116B	Shannon Zezula
IL-1	Illinois	Champaign	111	Kenton Macy
IL-2	Illinois	Champaign	110	Graciela Moreno
IL-3	Illinois	Piatt	108	Kenton Macy
IL-4	Illinois	Lawrence	114	Kenton Macy
IA-1	Iowa	Monona	107	Drew Delang
IA-2	Iowa	Adams	108	Mark Palmquist
IA-3	Iowa	Jones	105	Joe Wagner
IA-4	Iowa	Decatur	109	Kevin Reynolds
IA-5	Iowa	Dickinson	103	Carroll Oskvig
IA-6	Iowa	Dickinson	103	Carroll Oskvig
IA-7	Iowa	Dickinson	103	Carroll Oskvig
IA-8	Iowa	Dickinson	103	Carroll Oskvig
IA-9	Iowa	Dickinson	103	Carroll Oskvig
IA-10	Iowa	Dickinson	103	Carroll Oskvig
IA-11	Iowa	Dickinson	103	Carroll Oskvig
IA-12	Iowa	Dickinson	103	Carroll Oskvig
IA-13	Iowa	Dickinson	103	Carroll Oskvig
IA-14	Iowa	Dickinson	103	Carroll-Oskvig
IA-15	Iowa	Iowa	108	Timothy Meyer
IA-16	Iowa	Decatur	109	Melvin Moe
IA-17	Iowa	Henry	108C	Dova Ensminger
IA-18	Iowa	Jefferson	N/A	Shawn Dettmann
IA-19	Iowa	Louisa	108C	Shawn Dettmann

**Study ID Code:** MOPMC-P-0003-PA,WL

**Study Title** - Evaluation and Release of Eastern Gamagrass, *Tripsacum dactyloides*, L.

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

Eastern gamagrass, *Tripsacum dactyloides* L., is a tall warm season perennial grass found from Florida to Texas and Mexico, north and west to Massachusetts, New York, Michigan, Illinois, Missouri, Iowa and Nebraska. Eastern gamagrass grows in large clumps with thick rhizomes, broad flat leaves, the staminate and pistillate flowers in separate parts of the same many-flowered spikes. The pistillate spikelets are solitary and occur in hollowed portions on opposite sides of the thickened hard joints of the lower part of the rachis; this pistillate portion breaks up at maturity into several one-seeded joints. The staminate spikelets are two-flowered and in pairs on one side of a continuous rachis. Eastern gamagrass occurs on prairies, open limestone slopes, borders of woods and thickets, fields, and along roadsides and railroads. Eastern gamagrass is considered by many to be the ice-cream grass of the prairie. It is high in forage production and quality.

**Problem:**

The variety most commonly used in the PMC service area is ‘Pete’ and it performs well although its origin is Oklahoma and Kansas. This species is common in the PMC service area and a more adapted and improved variety should be able to be developed from native collections.

**Objectives:**

To evaluate and compare the variety ‘Pete’ with the best accessions from PMC study 29I107G and accessions developed at Woodward, Oklahoma.

Release an adapted variety and or varieties of eastern gamagrass for forage production and conservation uses in Missouri, Iowa, Illinois, Indiana and Ohio.

**Cooperators:**

Agriculture Research Service (ARS) Southern Plains Range Research Station, Woodward, Oklahoma.

**Procedure:**

Accessions selected previous work (Study 29I107G) at the Elsberry PMC and the Southern Plains Range Research Station at Woodward, Oklahoma will be assembled in 2000. Plants will be started in the greenhouse and planted in a randomized complete block with four replications. Plot size is nine feet by 18 feet consisting of three rows of plants, six plants per row with a three-foot spacing. The accessions will be tested for forage quality and production twice a year for three years.

**Discussion:****2000**

Plants arrived from Oklahoma in May and the study was planted in Field #9, pipeline D and E, June 28, 2000 and July 12, 2000. The plants from Elsberry were not as old so they were allowed to catch up. Plot map can be seen in Table #1.

**2001**

The plants established well in 2000 and only a few border row plants were replaced in 2001. Evaluations were taken on the interior four plants of each plot. Three forage harvests were taken during 2001 to compare yield and quality. Samples were sent to Woodward, Oklahoma for analysis. Evaluations will continue for three years. Results of 2001 harvests can be seen in Table #2.

**2002**

Forage harvests were taken twice in 2002 instead of three times as in 2001 because of the extremely dry summer. The first harvest was June 21, 2002 and the second harvest was delayed until August 6, 2002. These samples were sent to Woodward, Oklahoma for analysis. Evaluations will continue through 2003.

**MOPMC-P-0003-PA,WL**

**Elsberry PMC Field #9**

**Pipeline D and E Table #1**

Rep 4	9061911	FTIV	FTII	9083214		
	FT94-8	9061924	FTG1	Pete	X	X
Rep 3	FTII	9061911	Pete	FTIV	P E	F T
	FTG1	9083214	FO94-8	9061924	T E	II
Rep 2	Pete	FTIV	FTII	FT94-8	X X X	X X X
	9083214	9061924	9061911	FTG1	X 6	X F
Rep 1	9061911+C34	FT94-8 \1	FTIV \1	9061924	1 9 2	T G 1
	FTII	Pete	9083214	FTG1	4 X	 X

Plot Size: 9' x 18'

Planted 6/28/00, 7/12/00

3 rows of plants	X X X X X X
6 plants per row	X X X X X X
3 foot spacing	X X X X X X

\1 Southeast plant in plot was substituted with Pete because proper accession was not available.

\2 Above plots consisted of ten plants each for seed production information.

FTIV – Fertile Triploid OK accession	9061911 – Diploid MO accession
FT II – Fertile Triploid OK accession	9061924 – Diploid MO (North) accession
FGT I – Fertile Gynomonecious Triploid OK accn.	9083214 – Diploid Cross MO accession
FT 94-8 Fertile Triploid OK accession	'Pete' varietal release (Check)

Eastern gamagrass forage yield, crude protein (CP), and in vitro dry matter digestibility (IVOMD) for each harvest at the Elsberry Plant Materials Center (USDA-NRCS) near Elsberry, Missouri in year 2001.

Year 2001		Harvest Date			
Cultivar or Experimental Line		1 June 25	2 August 10	3 September 27	Total or Average *
<b>Pete</b>	<b>Yield (lb/a)</b>	1760 ab	3560 ab	1250 ab	6570 ab
	<b>CP(%)</b>	12.7 bc	8.2 a	11.9 a	9.6 abe
	<b>IVOMD(%)</b>	59.0 a	49.6 ab	54.5 b	52.6 ab
<b>FGT-1</b>	<b>Yield (lb/a)</b>	775 bc	2845 b	1035 ab	4655 b
	<b>CP(%)</b>	17.2 a	9.1 a	10.8 a	10.8 ab
	<b>IVOMD(%)</b>	63.7 a	49.0 b	55.0 b	52.2 b
<b>FT II</b>	<b>Yield (lb/a)</b>	2825 a	5000 a	1490 ab	9315 a
	<b>CP(%)</b>	11.6 c	7.4 a	11.0 a	8.9 bc
	<b>IVOMD(%)</b>	60.8 a	54.1 a	59.3 a	56.5 a
<b>FT IV</b>	<b>Yield (lb/a)</b>	1780 ab	3915 ab	1260 ab	6955 ab
	<b>CP(%)</b>	12.7 bc	7.6 a	11.3 a	9.2 bc
	<b>IVOMD(%)</b>	60.4 a	50.7 ab	55.8 b	53.6 ab
<b>FT 94-8</b>	<b>Yield (lb/a)</b>	185 c	805 c	595 b	1290 c
	<b>CP(%)</b>	15.3 abc	7.0 a	10.5 a	8.6 c
	<b>IVOMD(%)</b>	62.1 a	43.3 c	48.9 c	46.8 c
<b>1911</b>	<b>Yield (lb/a)</b>	1005 bc	2740 b	1030 ab	4770 b
	<b>CP(%)</b>	15.8 ab	9.5 a	11.2 a	11.2 a
	<b>IVOMD(%)</b>	60.6 a	52.5 ab	53.6 b	54.6 ab
<b>61924</b>	<b>Yield (lb/a)</b>	1410 bc	2955 ab	1200 ab	5565 b
	<b>CP(%)</b>	13.7 abc	9.5 a	11.4 a	10.8 ab
	<b>IVOMD(%)</b>	59.9 a	50.7 ab	54.4 b	53.6 ab
<b>83214</b>	<b>Yield (lb/a)</b>	2735 a	4630 ab	1745 a	9110 a
	<b>CP(%)</b>	14.0 abc	8.6 a	9.9 a	10.2 abc
	<b>IVOMD(%)</b>	59.7 a	51.1 ab	53.6 b	54.0 ab

\*Yield is calculated as a total of all harvests.

CP and IVOMD are calculated as weighted averages of all harvests

**Study ID Code: MOPMC-T-0104**

**Study Title: Native Plant Identification**

**Study Leader: Henry, J.**

**Description:**

Plant identification by landowners and NRCS Field Personnel is very challenging in the early seedling stages. As a result, plant identification workshops are being held in several locations in Missouri, Illinois and Iowa. As a result of these sessions, a need has developed which would make available via PowerPoint or on the web, color digital photos illustrating different characteristics of native grasses, legumes, and forbs. These photos could then be used during training workshops or extracted from the web by individuals interested in specific plant identification.

**Objective:**

There are many publications presently available for use regarding plant identification, however the majority of these deal only with identification of matured plants. Not much information (photos) is readily available regarding seedling identification of native grasses, forbs, and legumes. The objective is to make available, particularly over the web, color photos of seedlings of native grasses (cool and warm season), legumes and forbs.

**Materials and Methods:**

Assemble seed and plants of selected native cool and warm season grasses, legumes and forbs and take color photos at different stages of growth.

**Discussion:**

**2001**

The study plan was developed and approved in August 2001. A total of 31 different species of native grasses (cool and warm season), legumes and forbs are involved in this study: five cool season grasses, four warm season grasses, five legumes and 17 forbs. Color photos will be taken of the seed of each species, seven, 14 and 30 days after germination and at flowering and seed set. These photos will then be placed on the plant materials web site and made available to those individuals interested in these plants.

The following is a listing of plant species included in this study.

**2002**

Digital photos and color slides are being taken of New England aster, roundhead lespedeza, oxeye false sunflower, Canada wildrye, Virginia wildrye, stiff goldenrod, prairie blazing star, oxeye false sunflower and rattlesnake master. Photos are being taken of the seed, seven days, 14 days, and 30 days after germination, at flowering, and seed set. Photos will then be placed on the plant materials web site and made available to NRCS employees and other individuals interested in these plants.

<b>Native Cool Season Grasses</b>	<b>Native Warm Season Grasses</b>
<i>Elymus canadensis</i>	<i>Spartina pectinata</i>
<i>Elymus virginicus</i>	<i>Paspalum laeve</i>
<i>Calamagrostic canadensis</i>	<i>Dichanthelium clandestinum</i>
<i>Cinna arundinacea</i>	<i>Sporobolus asper</i>
<i>Uniola latifolia</i>	
<b>Native Forbs</b>	<b>Native Legumes</b>
<i>Liatris pycnostachya</i>	<i>Lespedeza capitata</i>
<i>Eryngium yuccifolium</i>	<i>Desmodium canadense</i>
<i>Coreopsis palmata</i>	<i>Dalea purpureum</i>
<i>Ratibida pinnata</i>	<i>Dalea candide</i>
<i>Aster novae-angliae</i>	<i>Tephrosia virginiana L.</i>
<i>Heliopsis helianthoides</i>	
<i>Echinacea pallida</i>	
<i>Monarda fistulosa</i>	
<i>Zizia aurea</i>	
<i>Asclepias tuberosa</i>	
<i>Solidago rigida</i>	
<i>Silphium laciniatum</i>	
<i>Veronicastrum virginicum</i>	
<i>Penstemon digitalis</i>	
<i>Lobelia siphilitica</i>	
<i>Desmanthus illinoensis</i>	
<i>Liatris aster</i>	

**Study ID Code:** MOPMC-T-0105, PA

**Study Title – Compatibility Study Using Native Warm Season and Cool Season Grasses with Native Legumes and Forbs**

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

Herbaceous plantings using native species are often a single grass species or a mixture of grasses with few legumes or forbs. These types of plantings are typical for forage, conservation cover or even wildlife plantings. Many native forbs and legumes are compatible with native grass species in a native prairie. In a planting using native species it is important to know which ones are most likely to compete with the grasses during the establishment period. Forb and legume seed is more expensive than the grass seed and most plantings lack diversity.

**Problem:**

There is little to no documented information regarding the compatibility of native warm and cool season grasses with native legumes and forbs in a pasture or range seeding. As a result of the lack of this needed information, the PMC Advisory Committee has directed the PMC to initiate this study.

**Objective:**

The objective of this study is to determine which native forbs and legumes will establish the easiest and persist the longest with specific native grasses.

**Procedure:**

Secure seed of the following native cool and warm season grasses, forbs, and legumes.

**Cool Season Grasses:** Virginia wildrye, Western wheatgrass, Junegrass, and Porcupinegrass. **Warm Season Grasses:** Eastern gamagrass, Little bluestem, Big bluestem, Indiangrass, and Switchgrass.

**Forbs:** Oxeye daisy, Prairie coreopsis, and Grayhead coneflower. **Legumes:** Bush clover, *Desmodium canadense*, Purple prairie clover, White prairie clover, Illinois bundleflower, Goat's'rue, Wild senna, and Lead plant.

Plots of a native warm season grass mixture, native cool season grass mixture and warm and cool season grass mixture will be established in four replications. Native legumes and forb mixtures will be planted with the grass mixtures. Plots will be planted in the spring and also as winter dormant plantings. All species will also be planted at the same time in the spring and winter except one warm and cool season grass mixture.

Plots will be mowed for weed control during the establishment year. The forage will be removed two to three times a year from half the plot the following years to assimilate rotational grazing.

**Discussion:****2001**

A site was prepared on the PMC using glyphosate to kill existing vegetation that consisted of mostly annual weedy species. The area was then plowed, disked and planted to an annual covercrop of 80% oats and 20% wheat. Plot composition of species can be seen in Table #1. Seeding rates are 40 pure live seed per square foot with 60% being the grass component and 40% being the forb and legume component.

**2002**

The winter dormant plots were planted January 8 and 9, 2002 using a plot planter. Seeding depth was on fourth inch for all species except the eastern gamagrass which was planted three fourths to one inch deep. The spring plots were planted May 20 and 21. All seed was planted at a depth of one fourth to one half inch with the exception of eastern gamagrass which again was planted at a depth of three fourths to one inch. All species that required treatment were stratified and/or scarified and inoculated. The plot map of the winter dormant planting is Table #2 and the spring planting is Table #3.

Mowing throughout the summer was the weed control method used. The plots were mowed when vegetation reached six to eight inches. Mowing height was three to four inches.

All plots were evaluated toward the end of the growing season for species composition. Most of the grasses were represented in the plots but in very low densities. Only sideoats gramma and Virginia wildrye appeared in plots in densities in the moderate range. The only legumes/forbs that were identified even at low densities were winter dormant planting – Illinois bundleflower, grayheaded coneflower, and prairie coriopsis, and spring planting – Illinois bundleflower and wild senna.



STUDY MOPMC-T-0105 Native Grass/Legume/Forb Compatibility Study										Field #1		Table #2					
										Eastside							
										Dormant planting							
										← Terrace →		North ↓					
REP #1	1	2	3	4	5	9	7	8	← Terrace →								
REP #2	4	7	9	1	6	8	3	2	Rep 1	Rep 2	Rep 3	Rep 4					
REP #3	6	1	4	2	9	3	8	5	2	4	5	9	8	1	6	3	Rep 4
										← Roadway →							
Plot #1	Big bluestem, switchgrass, Legume and Forb Mixture																
Plot #2	Little bluestem, sideoats gramma, Legume and Forb Mixture																
Plot #3	Eastern gamagrass, Legume and Forb Mixture																
Plot #4	Timothy, switchgrass, Legume and Forb Mixture																
Plot #5	Virginia wildrye, Indiangrass, Legume and Forb Mixture																
Plot #6	Virginia wildrye, western wheatgrass, Legume and Forb Mixture																
Plot #7	Junegrass, porcupine grass, Legume and Forb Mixture																
Plot #8	Eastern gamagrass, Kura clover																
Plot #9	Check Legume Mixture Only																

STUDY MOPMC-T-0105 Native Grass/Legume/Forb Compatibility Study														Field #1		Table #3			
														West side					
														spring planting					
														planted 5/20-21/02					
																North ↓			
← Terrace →																			
REP #2	7	3	8	1	6	9	4	5	2	4	7	3	8	6	1	5	9	2	REP #4
REP #1	1	7	9	1	6	8	3	2	9	3	8	1	6	4	9	2	5	7	REP #3
← Roadway →																			
Plot #1 Big bluestem, switchgrass, Legume and Forb Mixture																			
Plot #2 Little bluestem, sideoats gramma, Legume and Forb Mixture																			
Plot #3 Eastern gamagrass, Legume and Forb Mixture																			
Plot #4 Timothy, switchgrass, Legume and Forb Mixture																			
Plot #5 Virginia wildrye, Indiangrass, Legume and Forb Mixture																			
Plot #6 Virginia wildrye, western wheatgrass, Legume and Forb Mixture																			
Plot #7 Junegrass, porcupine grass, Legume and Forb Mixture																			
Plot #8 Eastern gamagrass, Kura clover																			
Plot #9 Check Legume Mixture Only																			

**Study ID Code : MOPMC-T-0106, BU**

**Study Title: Collection and Evaluation of Native Cool Season Grasses and Sedges for Filter Strips**

**Study Leader:** Henry, J.

**Description:**

A need has developed out of a three-state technical review committee and approved by the State Conservationists' Advisory Committee to evaluate different native cool season grasses and sedges for filter strips.

Grasses and sedges to be considered are: Virginia wildrye, *Elymus canadensis*; Canada wildrye, *Elymus canadensis*; Junegrass, *Koeleria cristata*; bluejoint, *Calamagrostis canadensis*; sweet woodreed, *Cinna arundinacea*; river oats, *Uniola latifolia*; longhair sedge, *Carex cosmosa*; Frank sedge, *Carex frankii*; shoreline sedge, *Carex hyalinolepis*; wheat sedge, *Carex atherodes*; and greater straw sedge, *Carex normalis*.

**Objective:**

There is little to no documented information regarding native cool season grasses and sedges being used in filter strip situations. In an attempt to respond to this lack of information, the PMC has been directed to initiate this study. Depending upon the performance of selected native cool season grasses and sedges in filter strip situations, previous recommendations may change to include those native cool season grasses and sedges performing excellently in this situation.

**Discussion:**

**2001**

The study plan for this study was initiated and approved by the State Conservationists' Advisory Committee in August of 2001. Selected field offices in the PMC service area will be contacted in the spring of 2002 requesting their participation in this collection, however everyone is welcomed to participate. One to three collections per state per species are being requested, both seeds and plants. The plants will be grown in the PMC greenhouse and later transplanted in randomized complete blocks. Each block will be one foot wide and five feet long with approximately 30 plants per plot. The spacing of the plants in the blocks will be six inches x six inches.

**2002**

Collections of native sedges and cool season grasses began on July 2, 2002. The following chart reflects a listing of the collections made as of the time this report was developed. The collection period was extended one more year to make collections of

those species that have not been made or those species needing more collections. Field #10 on the PMC has been selected as the site for this study because of the access to water. Two accessions of river oats were planted (vegetatively) on September 9, 2002.

Scientific Name	Common Name	Collector	City, State	Temp. Acc. No
<i>Carex crus-corvi</i>	Ravens foot sedge	Dennis Shirk	Vienna, MO	MO-1
<i>Carex grayii</i>	Gray sedge	Dennis Shirk	Vienna, MO	MO-2
<i>Carex atherodes</i>	Slough Sedge	Dennis Shirk	Vienna, MO	MO-3
<i>Carex vulpinoidea</i> Michx.	Fox sedge	Dennis Shirk	Vienna, MO	MO-4
<i>Carex vulpinoidea</i> Michx.	Fox sedge	Kaiser & Henry	Elsberry, MO	MO-5
<i>Carex hyalinolepis</i> Steud.	Thinscale scale	Kaiser & Henry	Elsberry, MO	MO-6
<i>Carex crus-corvi</i> Shuttlew	Crowfoot sedge	Kaiser & Henry	Elsberry, MO	MO-7
<i>Carex hyalinolepis</i> Steud.	Thinscale sedge	Paul Freese	Albany, MO	MO-8
<i>Carex vulpinoidea</i> Michx	Fox sedge	Kaiser & Henry	Elsberry, MO	MO-9
<i>Scirpus atrovirens</i>	Green bulrush	Kaiser & Henry	Elsberry, MO	MO-10
<i>Scirpus atrovirens</i>	Green bulrush	Kaiser & Henry	Elsberry, MO	MO-11
<i>Carex frankii</i> Kunth.	Franks sedge	Paul Freese	Albany, MO	MO-12
<i>Carex lupulina</i> Muhl.	Hop sedge	Raleigh Redman	Warrensburg, MO	MO-13
<i>Carex grayii</i>	Gray's sedge	Raleigh Redman	Warrensburg, MO	MO-14
<i>Carex hyalinolepis</i> Steud.	Thinscale sedge.	Raleigh Redman	Warrensburg, MO	MO-15
<i>Carex frankii</i> Kunth	Frank's sedge	Lingwall & Ellis	Ralls Co., MO	MO-17
<i>Carex crus-corvi</i>	Crowfoot sedge	Lingwall & Ellis	Ralls Co., MO	MO-18
<i>Carex hyalinolepis</i> Stued.	Thinscale sedge	Lingwall & Ellis	Ralls Co., MO	MO-19

Continued –

Scientific Name	Common Name	Collector	City, State	Temp. Acc. No
<i>Carex frankii</i> Kunth	Frank's sedge	Raleigh Redman	Warrensburg, MO	MO-20
<i>Chasmanthium latifolium</i>	River oats	J. Kaiser	Troy, MO	MO-21
<i>Chasmanthium latifolium</i>	River oats	Travis Dinsdale	Springfield, MO	MO-22
<i>Chasmanthium latifolium</i>	River oats	Rodney Doolen	Puxico, MO	MO-23
<i>Chasmanthium latifolium</i>	River oats	J. Kiaser	Troy, MO	MO-24
<i>Carex hyalinoepis</i> Steud.	Thinscale sedge	Dave Hiatt	Martinsville, IL	IL-1
<i>Carex normalis</i>	Greater straw sedge	Dave Hiatt	Martinsville, IL	IL-2
<i>Carex lupulina</i> Muhl.	Hop sedge	Christine Talige	Fawfield, IA	IA-1

**Study ID Code:** MOPMC-P-0107-PA,WL

**Study Title :** Evaluation and Release of Big Bluestem, *Andropogon gerardii*, L.

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

Big bluestem, *Andropogon gerardi* L., is a tall, native warm season perennial grass with stiff, erect culms; flattened and kneeled sheaths; membranous ligules; and flat or folded leaf blades. Big bluestem has developed a very efficient spreading root system that may reach depths of five-eight feet. Big bluestem reaches a mature height of three-four feet in northern latitudes, and six-eight feet or more in the southern part of its natural range. Although short rhizomes may be present, it usually makes a bunch type growth. Big bluestem is composed of many ecotypes with a wide range of adaptation to soil and climate.

Big bluestem is one of the most widespread important forage grasses of the North American tallgrass prairie region. It is usually associated with one or more of the other three dominant species, Indiangrass, switchgrass, and little bluestem. Big bluestem occurs on subirrigated lowlands, nearly level to gently undulating glacial till plains, overflow sites, level swales and depressions, residual and glacial uplands, and stream terraces and bottomlands along rivers and tributaries. The abundant, leafy forage is palatable to all classes of livestock.

**Problem:**

The variety most commonly used in the PMC service area is Rountree with its origin being Iowa. It performs well although it's performance decreases as it is moved south.

**Objectives:**

This species is common in the PMC service area and a more adapted and improved variety should be able to be developed from native collections from the southern part of the service area.

**Cooperators:**

NRCS Plant Materials Centers in Elsberry, Missouri and Booneville, Arkansas and the Agriculture Research Service (ARS) Southern Plains Range Research Station, Woodward, Oklahoma.

**Procedure:**

Accessions selected from previous work (Study 29I097G) at the Elsberry and Arkansas PMC's and the Southern Plains Range Research Station at Woodward, Oklahoma, will be assembled in 2001. Plants will be started in the greenhouse and planted in replicated plots the spring of 2002. Plot size is nine feet by 18 feet consisting of three rows of plants, six plants per row with a three-foot spacing.

The accessions will be compared to Rountree, Kaw, and Woodward and be tested for forage quality and production twice a year for three years.

**Discussion:****2001**

Seed was not available from all accessions to be included in the study so establishment of plots for this study was postponed.

**Study ID Code: MOPMC-T-0208-PA**

**Study Title: Testing Warm Season Grasses for Forage Quality**

**Study Leader:** Bruckerhoff, S.

**Description:**

Warm season grass species have limited information available in regards to forage quality. Confidence exists in their forage production abilities in relation to beef production. Comparative information on forage quality throughout the growing season is often questioned with little available information to back it up.

## **Objective:**

There is information in regards to forage quality of warm season species but usually it is for a specific variety and generally not throughout the growing season. The objective of this study is to make information available on forage quality throughout the growing season of warm season grass species. This information can be used to compare the quality of warm season grass species during the growing season and also at different vegetative stages.

## **Materials and Methods:**

Commercially available cultivars/selections and accessions in advanced testing of a wide variety of warm season grasses will be assembled for testing. Plants of each accession will be started in the greenhouse from seed and transplanted into an evaluation nursery.

### **Species/cultivars/accessions to be tested**

'Rountree' big bluestem	'Rumsey' Indiangrass
9078831 big bluestem	9083214 eastern gamagrass
'Cave-In-Rock' switchgrass	'Pete' eastern gamagrass
9062244 switchgrass	'El Reno' sideoats gramma
'Aldous' little bluestem	'Osarka' bermudagrass

The experimental design is a randomized complete block with four replications. The plants will be clipped. Original growth and regrowth dry matter will be analyzed for forage quality.

### **Testing times are as follows:**

#### **Original Growth Material**

**Mid-May**  
**Mid-May**  
**Mid-May**  
**Mid-May**

#### **Regrowth Material**

**Mid May-Mid July taken Mid July**  
**Mid May-Mid Aug taken Mid Aug**  
**Mid July-Mid Aug taken Mid Aug**  
**Mid Aug-Mid Sept taken Mid Sept**

## **Discussion:**

**2002**

The plants listed above will be propagated in the greenhouse and transplanted into an evaluation nursery the spring of 2003.

**Study ID Code: MOPMC-P-0209, PA, WL**

**Study Title – Evaluation and Release of Paspalum Species**

**Study Leader:** Bruckerhoff, S. B.

**Introduction:**

There are several species of paspalums that occur within the three state service area of Missouri, Iowa, and Illinois. Most of the species only occur in the southern part of the region but some do extend farther north. Plants growing in wild areas appear to have good characteristics for a late maturing summer grass for pasture and hay. It is consumed by livestock when it appears in a pasture.

Paspalum, *Paspalum L.*, is a tall, native warm season perennial grass with soft leaves and an inflorescence of several racemes arranged digitately, pinnately, or rarely solitarily. The almost round to ovate spikelets, which articulate below the glumes, are 2 or 4 rows on 1 side of the rachis and are flat on one side, curved on the other (plano-convex), with a somewhat flattened appearance. The first glume is usually lacking.

**Problem:**

Paspalum has not been included in warm season pasture and hay plantings because of no available seed source and lack of information on forage production and quality.

**Objective:**

The objective of this study is to determine if paspalum has high enough forage quality to justify development of a cultivator with a local origin. If paspalum samples show high forage quality, plants will be collected from the three-state service area and an evaluation and selection program will be started.

**Procedure:**

This study has two parts.

**Part I**

Identify as many different species of paspalum growing in pastures, right-of-ways, etc., as possible and test a forage sample for percent protein, ADF, and NDF.

Samples will be mailed to the PMC along with collection information including collection location, growth stage and date of collection. Samples will be sent off for analysis.

Determination of starting Part II will be made from the results of the analysis.

## **Part II**

Vegetative samples will be collected from prairie remnants throughout the service area where paspalum naturally exists. A minimum of five collections per MLRA/Administrative Area will be made. Collections will be divided, grown in the greenhouse, and transplanted into a randomized complete block (RCB) with four replications.

These accessions (collections) will be evaluated for forage quality, quantity, vigor, maturity, and disease and insect resistance. Superior selections will continue in a plant-breeding program.

### **Discussion:**

#### **2002**

Forage analyses of samples received in 2002 are as follows:

<b><u>Clipping date</u></b>	<b><u>Notes</u></b>	<b><u>Crude Protein</u></b>	<b><u>ADF</u></b>	<b><u>NDF</u></b>	<b><u>TDN</u></b>
7/22/02	North Missouri	9.0	37.4	64.2	59.8
7/2/02	Original Material	10.2	39.0	68.4	58.5
7/2/02	Regrowth Material	11.6	38.6	62.8	58.9
8/24/02	Cole County	6.2	34.8	66.9	61.8
8/24/02	Gasconade County	7.9	42.7	71.0	55.6
8/24/02	Maries County	6.1	44.7	71.5	54.1

Releases from the Elsberry Plant Materials Center						
Scientific Name	Release Name	Common Name	Accession Number	Secondary Agency(ies)	Type of Release	Year of Release
<i>Elymus virginicus</i> L.	Cuivre River	Virginia wildrye	9803169	MDC	N	2002
<i>Aster novae-angliae</i> L.	Central Iowa	New England Aster	9068682	UNI, IARV, IAT, ICIA	N	2002
<i>Aster novae-angliae</i> L.	Northern Iowa	New England Aster	9068681	UNI, IARV, IAT, ICIA	N	2002
<i>Aster novae-angliae</i> L.	Southern Iowa	New England Aster	9068683	UNI, IARV, IAT, ICIA	N	2002
<i>Echinacea pallida</i> Nutt.	Northern Iowa	pale purple coneflower	9068611	UNI, IARV, IAT, ICIA	N	2002
<i>Echinacea pallida</i> Nutt.	Southern Iowa	pale purple coneflower	9068613	UNI, IARV, IAT, ICIA	N	2002
<i>Sporobolus compositus</i> var. <i>com.</i>	Southern Iowa	tall dropseed	9062315	UNI, IARV, IAT, ICIA	N	2002
<i>Solidago rigida</i> L.	Southern Iowa	rigid goldenrod	9068619	UNI, IARV, IAT, ICIA	N	2002
<i>Solidago rigida</i> L.	Central Iowa	rigid goldenrod	9068618	UNI, IARV, IAT, ICIA	N	2002
<i>Coreopsis palmata</i>	Northern MO	prairie coreopsis	9079028	MDC, NAS	N	2001
<i>Coreopsis Palmuta</i> Nutt.	Western MO	prairie coreopsis	9079029	MDC, NAS	N	2001
<i>Sporobolus compositus</i> var. <i>comp.</i>	Northern MO	tall dropseed	9079040	MDC, NAS	N	2001
<i>Liatris pycnostachya</i> , Michx.	Western MO	blazing star	9079021	MDC, NAS	N	2001
<i>Liatris pycnostachya</i> , Michx	Northern MO	blazing star	9079020	MDC, NAS	N	2001
<i>Sporobolus compositus</i> (Poir.) Merr.	Northern Iowa	tall dropseed	9062313	UNI, IARV, IAT, ICIA	N	2000
<i>Andropogon gerardii</i>	Northern Iowa	big bluestem	9068614	UNI, IARV, IAT, ICIA	N	2000
<i>Liatris pycnostachya</i> , Michx	Southern Iowa	prairie blazing star	9068628	UNI, IARV, IAT, ICIA	N	2000
<i>Lespedeza capitata</i> Michx.	Northern Iowa	roundhead lespedeza	9062284	UNI, IARV, IAT, ICIA	N	2000
<i>Andropogon gerardii</i> Vitman	Southern Iowa	big bluestem	9068616	UNI, IARV, IAT, ICIA	N	1999
<i>Schizachyrium scoparium</i> , Michx.	Northern Iowa	little bluestem	9062319	UNI, IARV, IAT, ICIA	N	1999
<i>Eryngium yuccifolium</i> Michx.	Southern Iowa	rattlesnake master	9068604	UNI, IARV, IAT, ICIA	N	1999
<i>Eryngium yuccifolium</i> Michx.	Central Iowa	rattlesnake master	9068603	UNI, IARV, IAT, ICIA	N	1999
<i>Schizachyrium scoparium</i> , Michx.	Southern Iowa	little bluestem	9962321	UNI, IARV, IAT, ICIA	N	1999
<i>Liatris pycnostachya</i> , Michx	Northern Iowa	prairie blazing star	9068626	UNI, IARV, IAT, ICIA	N	1999
<i>Liatris pycnostachya</i> , Michx	Central Iowa	prairie blazing star	9068627	UNI, IARV, IAT, ICIA	N	1999
<i>Elymus virginicus</i> L.	Northern MO	Virginia wild rye	9079044	UMC, MDC, MODOT	N	1999
<i>Sorghastrum nutans</i> (L) Nash.	Northern MO	indiangrass	9079036	UMC, MDC, MODOT	N	1999
<i>Andropogon gerardii</i> Vitman	Northern MO	big bluestem	9079000	UMC, MDC, MODOT	N	1999
<i>Sorghastrum nutans</i> (L) Nash.	Western MO	indiangrass	9079037	UMC, MDC, MODOT	N	1999
<i>Schizachyrium scoparium</i> , Michx.	Northern MO	little bluestem	9079004	UMC, MDC, MODOT	N	1999
<i>Andropogon gerardii</i> Vitman	Central Iowa	big bluestem	9068615	UNI, IARV, IAT, ICIA	N	1998
<i>Dalea purpurea</i>	Central Iowa	prairie clover	9068609	UNI, IARV, IAT, ICIA	N	1998
<i>Eryngium yuccifolium</i> Michx.	Northern Iowa	rattlesnake master	9068602	UNI, IARV, IAT, ICIA	N	1998
<i>Solidago rigida</i> L.	Northern Iowa	rigid goldenrod	9068617	UNI, IARV, IAT, ICIA	N	1998
<i>Sorghastrum nutans</i> (L.) Nash.	Southern Iowa	indiangrass	9062318	UNI, IARV, IAT, ICIA	N	1998
<i>Andropogon gerardii</i> Vitman.	OH-370	big bluestem	9062323	ARPMC	N	1997
<i>Cornus drummondii</i> C.A. Meyer	Corinth	roughleaf dogwood	9055632		N	1997
<i>Cornus drummondii</i> C.A. Meyer	Jefferson	roughleaf dogwood	9055650		N	1997
<i>Cornus drummondii</i> C.A. Meyer	Tazewell	roughleaf dogwood	9055667		N	1997
<i>Cornus drummondii</i> C.A. Meyer	Nicholson	roughleaf dogwood	9055594		N	1997
<i>Desmodium canadense</i> L.	Alexander	showy tick trefoil	9057110		N	1997
<i>Elymus canadensis</i> L.	Southern Iowa	canada wildrye	9062277	UNI, IARV, IAT, ICIA	N	1997
<i>Heliopsis helianthoides</i> (L.) Sweet	Southern Iowa	oxeye false sunflower	9068607	UNI, IARV, IAT, ICIA	N	1997
<i>Lespedeza capitata</i> Michx.	Southern Iowa	roundhead lespedeza	9062283	UNI, IARV, IAT, ICIA	N	1997
<i>Liriodendron tulipifera</i> L.	Union	tulip poplar	9055584		N	1997
<i>Schizachyrium scoparium</i> (Michx.) Nash	Central Iowa	little bluestem	9062320	UNI, IARV, IAT, ICIA	N	1997
<i>Heliopsis helianthoides</i> (L.) Sweet	Northern Iowa	oxeye false sunflower	9068605	UNI, IARV, IAT, ICIA	N	1996
<i>Lespedeza capitata</i> Michx.	Central Iowa	roundhead lespedeza	9062282	UNI, IARV, IAT, ICIA	N	1996
<i>Sorghastrum nutans</i> (L.) Nash	Central Iowa	Indiangrass	9062317	UNI, IARV, IAT, ICIA	N	1996

Scientific Name	Release Name	Common Name	Accession Number	Secondary Agency(ies)	Type of Release	Year of Release
Sorghastrum nutans (L.) Nash	Northern Iowa	Indiangrass	9062316	UNI,IARV,IAT,ICIA	N	1996
Sporobolus compositus (Poir.) Merr.	Central Iowa	tall dropseed	9062314	UNI,IARV,IAT,ICIA	N	1996
Bouteloua curtipendula (Michx.) Torr.	Central Iowa	sideoats grama	9062279	UNI,IARV,IAT,ICIA	N	1995
Bouteloua curtipendula (Michx.) Torr.	Northern Iowa	sideoats grama	9062278	UNI,IARV,IAT,ICIA	N	1995
Bouteloua curtipendula (Michx.) Torr.	Southern Iowa	sideoats grama	9062280	UNI,IARV,IAT,ICIA	N	1995
Elymus canadensis L.	Central Iowa	Canada wildrye	9062276	UNI,IARV,IAT,ICIA	N	1995
Elymus canadensis L.	Northern Iowa	Canada wildrye	9062275	UNI,IARV,IAT,ICIA	N	1995
Heliopsis helianthoides (L.) Sweet	Central Iowa	oxeye false sunflower	9068606	UNI,IARV,IAT,ICIA	N	1995
Panicum virgatum L. *	Shawnee	switchgrass	591824		N	1995
Cornus mas L.	Redstone	cornelian cherry dogwood	516476		I	1991
Lonicera maackii Maxim	Cling Red	Amur honeysuckle	483450		I	1978
Ulmus parvifolia Jacq.	Elsmo	lace bark elm	9004438		I	1990
Andropogon gerardii Vitman.	Rountree	big bluestem	474216	MOA	N	1983
Sorghastrum nutans (L.) Nash.	Rumsey	Indiangrass	315747	MOA	N	1983
Elaeagnus umbellata Thunb.	Elsberry	autumn olive	476986		I	1979
Acer ginnala Maxim.	Flame	Amur maple	483442		I	1978
Glycine sp. L **	Bobwhite	soybean	421822	MOPMC,ARS, MOA,	I	1975
Panicum virgatum L.	Cave-In-Rock	switchgrass	469228	MOA	N	1974
Bromus inermis Leyss.	Elsberry	smooth brome	469227	MOA	Nat.	1954
* Primary Agencies: ARS=Agricultural Research Service; NEARD=Nebraska Agricultural Research Division; MOPMC=Missouri Plant Materials Center; IAA=Iowa Agricultural Experiment Station at Ames; PARP=Purdue Agricultural Research Program						
** Primary Agency: MDC=Missouri Department of Conservation						
N=Native releases; collected within the USA, occurring naturally in the USA. Generally refers to a plant which occurs naturally in a particular region, state ecosystem or habitat without direct or indirect human activity.						
Nat.=naturalized releases; collected from a population within the USA, but were originally introduced to the USA sometime in the past.						
I=introduced; means that the original collection from which the release was made was not from within the USA.						

<b>Studies/Projects at the Elsberry Plant Materials Center</b>	
<b>Studies 1958 through 2001</b>	
Study/Project Number System: Initially the numbers were assigned numerically plus the year the study/project was initiated. Later a different numbering system was adopted which involved the designated state number, a letter to denote the type of project/study and finally a numerical number.	
<b>Study/Project No.</b>	
<b>Year Started</b>	<b>Title</b>
2-58	Quaker Comphrey Evaluation
3-58	Comparison of Winter Annual Cover Crops
6-62	Fertilizer Rate Study on Midland Bermudagrass, <i>Cynadon dactylon</i>
10-59	Interseeding Cover Crops in Corn
14-61	Evaluation of <i>Lotus corniculatus</i> L. Strains
15-61	Evaluation of Bermudagrass Strains
17-61	Black Locust, <i>Robinia pseudoacacia</i> L. Trials
18-61	The Rate, Date and Method of Seeding <i>Lespedeza daurica schmidae</i>
19-61	Living Fence Trials
20-61	Plants for Bank Stabilization
21-62	Evaluation of Legumes for Wildlife
23-63	Evaluation of <i>Phalaris arundinacea</i> L. 'loreed' Reed Canarygrass Strains
24-62	Method of Seeding Creeping Foxtail
25-63	Advanced Evaluation of Plant Materials for Grass Waterways
26-63	Evaluation of Japanese Pagodatree ( <i>Sophoro japonica</i> ) for Posts
27-63	Direct Seeding vs Transplanting Sawtooth Oak, <i>Quercus acutissima Carruthers</i>
28-63	Effect of Cultural Methods on Crownvetch, <i>Coronilla varia</i> L. Seed Production
31-63	<i>Lespedeza capitata</i> Michx. - Roundhead Lespedeza Ecotype Evaluation
34-63	Cultural Methods for Seeding Grasses in Woodland Pastures
35-63	Effect of Cultural Methods on Seed Production of <i>Phalaris arundinacea</i> L., 'loreed' Reed Canarygrass

Study/Project	Title
37-63	Forage Yields and Season of Production for Several Grasses and Legumes Clipped Bi-Weekly at Three Inches and Six Inches
38-64	Advanced Evaluation of Perennial Grasses for Summer Pasture
42 -65	Establishment of Crownvetch and Trefoil in Dead Litter Mulch
44-65	Grasses and Legumes for Goose Browse on the Clarence Cannon Wildlife Refuge
46-66	Method of Seeding Trials with 'Garrison' Creeping Foxtail
49-69	Seed Yield of Three <i>Panicum virgatum</i> , Switchgrass Selections: Mich 381; Blackwell', M1-5714; and M1-5845, 'Cave-In-Rock'
50-69	Seed Yield and Seed Retention of Four <i>Phalaris arundinacea</i> , Reed Canarygrass Selections: 'Ioreed', 'Rise', 'Frontier', and 'Auburn'
51-A-70	Herbicide Tolerance of Four Waterway Grasses: <i>Alopecurus arundinaceus</i> , 'Garrison' Creeping Foxtail; <i>Bromus inermis</i> , smoothbrome; <i>Phalaris arundinacea</i> , reed canarygrass; and <i>Panicum virgatum</i> , switchgrass
51-B-71	Herbicide Tolerance of New Seeding of <i>Festuca arundinacea</i> , Tall Fescue; <i>Andropogon gerardii</i> , Big Bluestem, <i>Sorghastrum nutans</i> , Indiangrass; and <i>Panicum virgatum</i> , Switchgrass
51-C-71	Herbicide Tolerance of New Seedling of Tall Fescue, Big Bluestem, Indiangrass and Switchgrass
29I052W	Growth Rate Study of European Alder on Deep Alluvial Soil
53-72	Growth Rate Study of Poplar (Cottonwood) On a Deep Alluvial Soil
54-72	Rhizome Development of Two Tall Fescue, <i>Festuca arundinacea</i> , Selections: M1-6161 and M1-6162
29A055	Evaluations of <i>Sorghastrum nutans</i> , Indiangrass (M17073), Poly-Cross Indiangrass for Leafiness, Disease-Free Characteristics and Seed Production
56-71	Comparative Evaluation of New Lotus Accessions With Names and Used Varieties to Determine Potential as a Long Lived Legume in Three State Area Saved
29I057-72	Growth Rate Study of Poplars (Cottonwood) On a Deep Alluvial Soil Deep Alluvial Soil
29A058-72	Evaluation for Naming and Releasing of Elsberry Developed Big Bluestem and Indiangrass

Study/Project	Title
59-72	Sorghum Evaluation as Wildlife Game Feed
29I060-69	Replacement of the American Elm Tree
61-72	Advanced Evaluation of Meadow Foxtail, <i>Alopecurus pratensis</i> , PI-305495, as a Waterway Grass as Compared to 'Garrison' Creeping Foxtail, <i>Alopecurus arundinaceus</i> the Standard for Comparison
29I062J	Trees and Shrubs for Use as Wildlife Food and Cover Plants
29I063	Plants for Use in Critical Area Stabilization
29I064W	Plants for Wood Products
65-78	Plants for Use in Landscape and Beautification
29I066W-72	Developing Winterhardy Nut Bearing Trees and Shrubs for Planting in Parks, Wildlife Areas and Natural Areas
29I067K	Trees for Windbreaks
68-72	Response of Yellow Poplar to Thinning
69-72	Black Cherry Demonstration
70-73	Desmodium for Wildlife Food and Cover
71-73	Evaluation for Naming and Releasing of Elsberry Developed Autumn Olive, M1-6369
72-73	Evaluation of M1-4701, <i>Lonicera maackii</i> , Amur Honeysuckle for Naming and Releasing
73-73	Establishment of Warm-Season Grasses with Herbicides for Weed Control. Herbicides are Not Tested or Have Label Clearance for Warm-Season Grasses
29A074M	Cover Crops in Soybeans
-----	NJ-927, <i>Eleagnus umbellata</i> , Autumn Olive for Wildlife Food and Cover
29A075F	Plants for Shoreline and Wetland Stabilization
29I076G-78	Establishment of Warm Season Grasses
-----	Evaluation of Cold Hardy <i>Paspalum notatum</i> Selections
29I077P	Evaluation of Plants for Vegetating Salt Damaged Areas

Study/Project	Name
29I078D	Field Evaluation Planting to Evaluate Plants for Use on Alkali Bearing Soils in Southern Illinois
29I079D	Field Evaluation Planting to Evaluate Species of Plants for Use on Revegetating Acid Coal Mine Spoil in Illinois
29I081D	Field Evaluation Planting to Evaluate Species of Plants for use in Revegetating Acid Coal Mine Spoil in Iowa
29I082D	Field Evaluation Planting to Evaluate Species of Plants for Use in Revegetating Acid Coal Mine Spoil in Illinois
29I083M	Legume Cover Crop for No-Till Corn Production
29I084G	Legumes to Enhance Fescue Pastures
29A085S	Debearding Fluffy Native Grass Seed, (Big Bluestem and Indiangrass)
29I086L	Use of an Absorbant Polymer in Coating Native Grass Seed
29I087D	Plants with Increased Tolerance to Aluminum and Manganese
29A088W	Cooperative Screening Study of Native and Introduced Sources of Eastern Cottonwood
29I089V	Multiple Use Legume Assembly and Evaluation
29I090G	No-Till Establishment of Warm-Season Grasses in Cool Season Grass Sod
29I091G	Weed Control Treatments for Warm Season Grass Establishment
29I092G	Perennial Grasses as Cover Crops for Use in No-Till Systems
29I093R	Miscellaneous Grass Evaluation
29A094M	Cover Crops in Corn, Soybeans and Milo
29A095M	Field Evaluation Planting to Evaluate Cover Crops - Rochester, Minnesota
29I097G	Assembly and Evaluation of Big Bluestem, <i>Andropogon gerardii</i> , Vitman.
29I099J	Assembly and Evaluation of Roughleaf Dogwood, <i>Cornus drummondii</i>
29I100J	Assembly and Evaluation of Blackhaw, <i>Viburnum prunifolium</i> L.
29I101J	Assembly and Evaluation of Arrowwood, <i>Viburnum dentatum</i> L.

Study/Project	Name
29A105M	Evaluation of Winter Annual Grass for Cover Crops in No-Till Soybeans
29I107G	Assembly and Evaluation of Eastern Gamagrass, <i>Tripsacum dactyloides</i> L.
29I108G	Assembly and Evaluation of Low Growing Rhizomatous Switchgrass, <i>Panicum virgatum</i> L., for Use in Waterways, Filter Strips and Other Conservation Uses
29I109W	Direct Seeding Methods of <i>Quercus</i> sp., Oaks
29I110J	Assembly and Evaluation of Chokecherry, <i>Prunus virginiana</i> L.
29A111G	Field Evaluation of Selected Perennial Grasses for Pasture Wildlife Habitat and Erosion Control (Varietal Study)
29I112J	Assembly and Evaluation of Nannyberry, <i>Viburnum lentago</i> L.
29I113J	Assembly and Evaluation of Serviceberry, <i>Amelanchier arborea</i> (Michx. F.) Fern.
29I114K	Field Evaluation of Woody Plant Materials in Cooperation with Mineral Area College
29A116W	Evaluation of Miscellaneous Trees and Shrub Species
29A117H	Intercenter Strain Trial of <i>Tripsacum dactyloides</i> L., Eastern Gamagrass
29A118G	Field Evaluation of Selected Perennial Grasses for Pasture, Wildlife Habitat and Erosion Control (Varietal Study)
29A121W	Conifer Evaluation for Windbreak Plantings
29A122G	Evaluation of Perennial Warm-Season Grasses as Windbarriers in Southeast Missouri
29A123M	Winter Cover Crop Study for No-Till Soybeans
29I124G	Production of Native Iowa Ecotypes of Grasses and Forbs for Roadside, Critical Areas, and All Other Vegetative Plantings Where Native Grasses and Forbs are Now Being Planted
29A125G	Fertility and Harvest Management of Eastern Gamagrass for Forage Production
29I126W	Woody Columnar Collection

Study/Project	Title
29A127G	Field Evaluation of Selected Perennial Grasses for Pasture, Wildlife Habitat and Erosion Control
29A128J	<i>Cornus florida</i> L., Flowering Dogwood, Interagency Study Between Department of Interior, National parks Service, National Capital Region and the Department of Agriculture
29A130G	Grass Hedges for Control of Runoff and Erosion
29A131O	Treatment of Animal Wastewaters by Constructed Wetlands
29I132O	Miscellaneous Wetland Plant Evaluation
29I133J	Assembly and Evaluation of Gray Dogwood, <i>Cornus racemosa</i>
29I134J	Assembly and Evaluation of Eastern Redcedar, <i>Juniper virginiana</i> L.
29I135J	Assembly and Evaluation of Hazelnut, <i>Corylus americana</i> , Marsh.
29I136J	Assembly and Evaluation of Wild Plum, <i>Prunus americana</i> , Marsh.
29A137O	Wetland Riparian Propagation, Establishment and Demonstration
29I138G	Residue Decomposition Trial
29A139G	Field Evaluation of Establishment of Herbaceous Plant Materials on Sand Covered Flooded Areas in Missouri
29A140W	Yellow Poplar Evaluation
29I141G	Assembly and Evaluation of Little Bluestem, <i>Schizachyrium scoparium</i> , Michx.
29I142G	Production of Native Missouri Ecotypes of Grasses, Legumes and Forbs for Roadside, Critical Areas, and All Other Vegetative Plantings Where Native Plants are Now Being Planted
29I143G	Seed Coat/Seeding Rates Study
29A144G	Biofuel Study of Different Strains/Varieties of Switchgrass
29A145	Wear Tolerance Demonstration of Vegetation in High Traffic Areas
MOPMC-P-0001 WO,WL,WE	Assembly, Evaluation and Selection of Bur Oak, <i>Quercus macrocarpa</i> , Michx.
MOPMC-P-0002 WE, WL	Assembly, Evaluation and Selection of False Indigo Bush,. <i>Amorpha fruticosa</i> , L.
MOPMC-P-0003 PA, WL	Evaluation and Release of Eastern Gamagrass, <i>Tripsacum dactyloides</i> , L.

Study/Project	Title
MOPMC-T-0104	Native Plant Identification
MOPMC-PA-0105	Compatibility Study Using Warm Season and Cool Season Native Grasses with Native Legumes and Forbs
MOPMC-BC-0106	Collection and Evaluation of Native Cool Season Grasses and Sedges for Filter Strips
MOPMC-P-0107	Evaluation and Release of Big Bluestem, <i>Andropogon gerardii</i> , L
MOPMC-T-0208-PA	Testing Warm Season Grasses for Forage Quality
MOPMC-0209-PA, WL	Evaluation and Release of <i>Paspalum</i> Species

## Herbaceous and Woody Seed and Plant Production at the Elsberry PMC 2002

The plant and seed inventory at the Elsberry PMC is used for field plantings, special plantings, demonstration plantings, research studies and commercial release. The 2002 production of grass, legume, forb, and woody seed reflected a below average year.

Name	Seed Inventory as of December 2002 PLS (Pounds)
<b>Herbaceous</b>	
<u>'Rountree' big bluestem</u> <i>Andropogon gerardii</i>	125 Foundation
<u>'Rumsey' indiangrass</u> <i>Sorghastrum nutans</i>	1,095 Foundation
<u>'Pete' eastern gamagrass</u> <i>Tripsicum dactyloides</i> L.	551 Foundation
<u>'Cave-In-Rock' switchgrass</u> <i>Panicum virgatum</i>	1,273 Foundation
<u>'Svalofs' field brome</u> <i>Bromus arvensis</i>	230
<u>'Elsberry' smoothbrome</u> <i>Bromus inermis</i>	41
<u>OH-370 big bluestem</u> <i>Andropogon gerardii</i>	82 Foundation
<u>'Niagara' big bluestem</u> <i>Andropogon gerardii</i>	56
<u>'Bobwhite' soybean</u> <i>Glycine species</i>	50
<u>'Aroostook' rye</u> <i>Secale cereale</i>	1,000

**Herbaceous and Woody Seed and Plant Production – continued**

Name:	Seed & Plant Inventory as of December 2002 Bulk (Pounds)	
	Plants	Seed
<u>Union tulip tree</u> <i>Liriodendron tulipifera</i>	18	0.6
<u>Nicholson Germplasm roughleaf dogwood</u> <i>Cornus drummondii</i>	139	1.2
<u>Corinth Germplasm roughleaf dogwood</u> <i>Cornus drummondii</i>	74	1.4
<u>Tazewell Germplasm roughleaf dogwood</u> <i>Cornus drummondii</i>	80	0.2
<u>Jefferson Germplasm roughleaf dogwood</u> <i>Cornus drummondii</i>	148	1.4
<u>American hazelnut (9057168) (Illinois)</u> <i>Corylus americana</i>	260	4.4
<u>American hazelnut (9057169) (Illinois)</u> <i>Corylus americana</i>	85	3.4
<u>American hazelnut (9068562) (Illinois)</u> <i>Corylus americana</i>	63	5.3
<u>American hazelnut (9057188) (Illinois)</u> <i>Corylus americana</i>	176	14.2
<u>American hazelnut (9068528) (Illinois)</u> <i>Corylus americana</i>	102	25.5
<u>American hazelnut (9068573) (Missouri)</u> <i>Corylus americana</i>		
<u>American hazelnut (9068574) (Missouri)</u> <i>Corylus americana</i>	249	4.8
<u>American plum (9068546) (Missouri)</u> <i>Prunus americana</i>	276	3.6
<u>American plum (9068580) (Missouri)</u> <i>Prunus americana</i>	141	1.6
<u>American plum (9057088) (Illinois)</u> <i>Prunus americana</i>	96	8.2
<u>American plum (9062309) (North Dakota)</u> <i>Prunus americana</i>	230	0.6
<u>American plum (9068545) (Missouri)</u> <i>Prunus americana</i>	276	1.0
<u>Arrowwood (9062310) (Iowa)</u> <i>Viburnum dentatum</i>	192	2.3



New England Aster  
*Aster novae-angliae* L.

For more information about this and other conservation plants, contact your local NRCS field office or Conservation District, or browse the Web at <http://Plant-Materials.nrcs.usda.gov> (Plant Materials) or <http://plants.usda.gov> (PLANTS database).

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audio tape, etc.) should contact the USDA's TARGET Center at 1-202-720-2600 (Voice and TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or 1-202-720-127 (TDD).  
USDA is an equal opportunity employer.